

J. S. Gupta; D. K. Bhola

Maximum modulus function of derivatives of entire functions defined by Dirichlet series

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 14 (1973), No. 3, 507--518

Persistent URL: <http://dml.cz/dmlcz/105505>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1973

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

MAXIMUM MODULUS FUNCTION OF DERIVATIVES OF ENTIRE FUNCTIONS  
DEFINED BY DIRICHLET SERIES

J.S. GUPTA, D.K. BHOLA, Jammu

Abstract: Let  $E$  be the set of mappings  $f: C \rightarrow C$  ( $C$  is the complex field) such that the image under  $f$  of a point  $s \in C$  is

$$f(s) = \sum_{n \in N} a_n e^{s \lambda_n} \quad \text{with } \limsup_{n \rightarrow +\infty} \frac{\log n}{\lambda_n} = D \in R_+ \cup \{0\}$$

( $R_+$  is the set of positive reals), and  $\sigma_c^f = +\infty$ . Then  $f$  is an entire function and is bounded on each vertical line  $\operatorname{Re}(s) = \sigma_0$ .

Denoting the maximum modulus function of the  $\mu$ -th derivative  $f^{(\mu)}$  of  $f$  for any  $\mu \in Z_+$  ( $Z_+$  is the set of positive integers) by  $M_\mu$ , we have investigated into some of its properties. In particular, we have shown that, for the members of a certain subset of  $E$ , the functions  $M_\mu$  and  $M_{\mu+1}$  are separated from each other by the derivative of the former for sufficiently large values of  $\sigma$ .

Key words: Entire function, Dirichlet series, maximum modulus function, Ritt order, lower order, convex function, ordinary proximate linear order.

AMS: 30A66

Ref. Ž. 7.549.2

1. Let  $E$  be the set of mappings  $f: C \rightarrow C$  ( $C$  is the complex field) such that the image under  $f$  of an element  $s \in C$  is

$$f(s) = \sum_{n \in N} a_n e^{s \lambda_n} \quad \text{with } \limsup_{n \rightarrow +\infty} \frac{\log n}{\lambda_n} = D \in R_+ \cup \{0\}$$

( $\mathbb{R}_+$  is the set of positive reals), and  $\sigma_c^f = +\infty$  ( $\sigma_c^f$  is the abscissa of convergence of the Dirichlet series defining  $f$ );  $\mathbb{N}$  is the set of natural numbers  $0, 1, 2, \dots$ ,  $\langle a_m \mid m \in \mathbb{N} \rangle$  is a sequence in  $\mathbb{C}$ ,  $s = \sigma + it$ ,  $\sigma, t \in \mathbb{R}$  ( $\mathbb{R}$  is the field of reals), and  $\langle \lambda_m \mid m \in \mathbb{N} \rangle$  is a strictly increasing unbounded sequence of nonnegative reals. Since the Dirichlet series defining  $f$  converges for each  $s \in \mathbb{C}$ ,  $f$  is an entire function. Also, since  $D \in \mathbb{R}_+ \cup \{0\}$ , we have ([1], p. 168)  $\sigma_a^f = +\infty$  ( $\sigma_a^f$  is the abscissa of absolute convergence of the Dirichlet series defining  $f$ ), and that  $f$  is bounded on each vertical line  $\operatorname{Re}(s) = \sigma_0$ .

For any  $n \in \mathbb{N}$ , the maximum modulus function  $M_n$  of the  $n$ -th derivative  $f^{(n)}$  of an entire function  $f \in E$ , on any vertical line  $\operatorname{Re}(s) = \sigma$ , is defined as

$$(1.1) \quad M_n(\sigma, f^{(n)}) = \sup_{t \in \mathbb{R}} \{|f^{(n)}(\sigma + it)|\}, \forall \sigma < \sigma_c^f.$$

We denote the function  $M_0$  by  $M$  and study a few properties of the function  $M_n$  in this paper.

2. For every entire function  $f \in E$ , Döetsch has shown ([2], p. 240) that  $\log M_n$  is a downward convex function of  $\sigma$ . We may, therefore, write, for any  $\sigma$ ,  $\sigma_0 \in \mathbb{R}$  such that  $\sigma > \sigma_0$ ,

$$(2.1) \quad \log M_p(\sigma, f^{(p)}) = O(1) + \int_{\sigma_0}^{\sigma} V(x, f^{(p)}) dx,$$

where  $V$  is a real valued indefinitely increasing function of  $\sigma$ , and establish

Theorem 1. If the function  $M_p$  assumes the value unity at the origin, and  $\sigma_1, \sigma_2, \dots, \sigma_m \in \mathbb{R}$  are such that  $0 \leq \sigma_1 \leq \sigma_2 \leq \dots \leq \sigma_m$ , then

$$(2.2) \quad \frac{1}{n} \sum_{1 \leq k_0 \leq m} \sigma_{k_0} V(\sigma_{k_0}, f^{(p)}) \geq \sigma_1 V(0, f^{(p)}) + \sum_{2 \leq k_0 \leq m} [\lambda_k \sigma_{k_0} - (\sigma_1 + \dots + \sigma_{k_0})] V\left(\frac{\sigma_1 + \dots + \sigma_{k_0-1}}{k_0 - 1}, f^{(p)}\right).$$

Proof. Using the convexity property of  $\log M_p$ , we get

$$\log M_p\left(\frac{m_1 \sigma_1 + \dots + m_m \sigma_m}{m_1 + \dots + m_m}, f^{(p)}\right) \leq \frac{m_1 \log M_p(\sigma_1, f^{(p)}) + \dots + m_m \log M_p(\sigma_m, f^{(p)})}{m_1 + \dots + m_m}$$

or

$$M_p\left(\frac{m_1 \sigma_1 + \dots + m_m \sigma_m}{m_1 + \dots + m_m}, f^{(p)}\right) \leq (M_p(\sigma_1, f^{(p)}))^{m_1} \dots (M_p(\sigma_m, f^{(p)}))^{m_m}^{1/(m_1 + \dots + m_m)},$$

or, supposing  $m_1 = \dots = m_m = 1$ ,

$$(2.3) \quad M_p\left(\frac{\sigma_1 + \dots + \sigma_m}{m}, f^{(p)}\right) \leq (M_p(\sigma_1, f^{(p)})) \dots \dots M_p(\sigma_m, f^{(p)})^{1/m}.$$

Since (2.1) is true for any  $\sigma, \sigma_0 \in \mathbb{R}$ , we take  $\sigma_0 = 0$

and  $\sigma_0 = \sigma_m$  getting

$$(2.4) \quad \log M_p(\sigma_m, f^{(n)}) = \int_0^{\sigma_m} V(x, f^{(n)}) dx \\ \leq \sigma_m V(\sigma_m, f^{(n)}),$$

and so  $M_p(\sigma_m, f^{(n)}) \leq \exp(\sigma_m V(\sigma_m, f^{(n)}))$ . Therefore,

$$(2.5) \quad (M_p(\sigma_1, f^{(n)}) \dots M_p(\sigma_m, f^{(n)}))^{1/n} \leq \\ \leq \exp(n^{-1}(\sigma_1 V(\sigma_1, f^{(n)}) + \dots + \sigma_m V(\sigma_m, f^{(n)}))).$$

Again, from (2.4),

$$\log M_p\left(\frac{\sigma_1 + \dots + \sigma_m}{m}, f^{(n)}\right) = \int_0^{(\sigma_1 + \dots + \sigma_m)m^{-1}} V(x, f^{(n)}) dx \\ = \left( \int_0^{\sigma_1} + \int_{\sigma_1}^{(\sigma_1 + \sigma_2)2^{-1}} + \dots + \int_{(\sigma_1 + \dots + \sigma_{m-1})(m-1)^{-1}}^{(\sigma_1 + \dots + \sigma_m)m^{-1}} \right) V(x, f^{(n)}) dx \\ (2.6) \quad \geq \sigma_1 V(0, f^{(n)}) + \frac{2\sigma_2 - (\sigma_1 + \sigma_2)}{2 \cdot 1} V(\sigma_1, f^{(n)}) + \dots + \\ + \frac{m\sigma_m - (\sigma_1 + \dots + \sigma_m)}{m(m-1)} V\left(\frac{\sigma_1 + \dots + \sigma_{m-1}}{m-1}, f^{(n)}\right).$$

But, from (2.4), we have

$$(2.7) \quad \log M_p(\sigma_1, f^{(n)}) \geq \sigma_1 V(0, f^{(n)}).$$

The theorem, therefore, follows from (2.3), (2.5), (2.6) and (2.7).

3. We next study two theorems concerning the derivative of

Theorem 2. If  $f \in E$  is an entire function of Ritt order  $\rho \in \mathbb{R}_+$  and lower order  $\lambda \in \mathbb{R}_+$  such that  $\lambda \geq \rho > 0$ , then for any  $\varepsilon \in \mathbb{R}_+$ , such that  $\varepsilon = \varepsilon(\sigma, f)$  tends to zero as  $\sigma$  tends to plus infinity, and sufficiently large  $\sigma$ ,

$$(3.1) \quad M'_\rho(\sigma, f^{(\rho)}) > M'(\sigma, f) \left( \frac{\log M'(\sigma, f)}{\left(\frac{1}{\lambda} - \frac{1}{\rho} + \varepsilon\right) \log \lambda_{\rho}(\sigma, f^{(\rho)})} \right)^\rho,$$

and

$$(3.2) \quad M''_\rho(\sigma, f^{(\rho)}) > M'(\sigma, f) \left( \frac{\log M'(\sigma, f)}{\left(1 - \frac{\lambda}{\rho} + \varepsilon\right)} \right)^\rho;$$

$\rho(\sigma, f^{(\rho)})$  is the rank of the maximum term  $\mu(\sigma, f^{(\rho)})$  in the Dirichlet series defining  $f^{(\rho)}$ .

The proof of this theorem is based on the following lemmas.

Lemma 1. For every entire function  $f \in E$  of Ritt order  $\rho \in \mathbb{R}_+$  and lower order  $\lambda \in \mathbb{R}_+$ ,

$$(3.3) \quad \limsup_{\sigma \rightarrow +\infty} \frac{\log M(\sigma, f)}{\lambda_{\rho}(\sigma, f) \log \lambda_{\rho}(\sigma, f)} \leq \frac{1}{\lambda} - \frac{1}{\rho},$$

and

$$(3.4) \quad \limsup_{\sigma \rightarrow +\infty} \frac{\log M(\sigma, f)}{\sigma \lambda_{\rho}(\sigma, f)} \leq 1 - \frac{\lambda}{\rho} .$$

The proof follows from the facts ([3], Theorem 5) that, as  $\sigma \rightarrow +\infty$

$$(3.5) \quad \log \mu(\sigma, f) \sim \log M(\sigma, f) ,$$

and ([4], p.57) that

$$(3.6) \quad \limsup_{\sigma \rightarrow +\infty} \frac{\log \mu(\sigma, f)}{\lambda_{\rho}(\sigma, f) \log \lambda_{\rho}(\sigma, f)} \leq \frac{1}{\lambda} - \frac{1}{\rho} ,$$

and

$$(3.7) \quad \limsup_{\sigma \rightarrow +\infty} \frac{\log \mu(\sigma, f)}{\sigma \lambda_{\rho}(\sigma, f)} \leq 1 - \frac{\lambda}{\rho} .$$

Lemma 2 ([5], p.254). For every entire function  $f \in E$  ,

$$(3.8) \quad M_{\rho}(\sigma, f^{(\rho)}) \geq M_{\rho-1}(\sigma, f^{(\rho-1)}) .$$

Proof of Theorem 2. We know ([6], p.67) that  $\log \mu$  and  $\log M$  are convex functions and (3.5) holds. It, therefore, follows that, as  $\sigma \rightarrow +\infty$  ,

$$\frac{\mu'(\sigma, f)}{\mu(\sigma, f)} \sim \frac{M'(\sigma, f)}{M(\sigma, f)} .$$

But ([7], p.241), for almost all values of  $\sigma$ ,

$$(3.9) \quad \frac{\mu'(\sigma, f)}{\mu(\sigma, f)} = \lambda_{\nu(\sigma, f)}.$$

Hence, for almost all sufficiently large  $\sigma$ ,

$$(3.10) \quad \frac{M'(\sigma, f)}{M(\sigma, f)} \sim \lambda_{\nu(\sigma, f)}.$$

From (3.3) and (3.10) we get

$$\limsup_{\sigma \rightarrow +\infty} \frac{\log M(\sigma, f)}{M(\sigma, f) \log \lambda_{\nu(\sigma, f)}} \leq \frac{1}{\lambda} - \frac{1}{\varphi}.$$

Hence, for any  $\varepsilon \in \mathbb{R}_+$  and sufficiently large  $\sigma$ ,

$$(3.11) \quad \frac{M'(\sigma, f)}{M(\sigma, f)} > \frac{\log M(\sigma, f)}{(\frac{1}{\lambda} - \frac{1}{\varphi} + \varepsilon) \log \lambda_{\nu(\sigma, f)}}.$$

Writing (3.11) for  $f^{(k)}$  we get

$$\frac{M'_{\nu_k}(\sigma, f^{(k)})}{M_{\nu_k}(\sigma, f^{(k)})} > \frac{\log M_{\nu_k}(\sigma, f^{(k)})}{(\frac{1}{\lambda} - \frac{1}{\varphi} + \varepsilon) \log \lambda_{\nu(\sigma, f^{(k)})}},$$

whence, in view of Lemma 2,

$$(3.12) \quad \frac{M'_{\nu_k}(\sigma, f^{(k)})}{M'_{\nu_{k-1}}(\sigma, f^{(k-1)})} > \frac{\log M'_{\nu_{k-1}}(\sigma, f^{(k-1)})}{(\frac{1}{\lambda} - \frac{1}{\varphi} + \varepsilon) \log \lambda_{\nu(\sigma, f^{(k)})}}.$$



Giving  $k$  the values  $1, 2, \dots, n$  in (3.12) and multiplying the resulting  $n$  inequalities, we get

$$(3.13) \quad M'_n(\sigma, f^{(n)}) > \frac{M'(\sigma, f) \prod_{1 \leq k \leq n} \log M'_{k-1}(\sigma, f^{(k-1)})}{\left(\frac{1}{\lambda} - \frac{1}{\rho} + \varepsilon\right)^n \prod_{1 \leq k \leq n} \log \lambda_{>}(\sigma, f^{(k)})}.$$

But ([5], p.254), for sufficiently large  $\sigma$ ,

$$M(\sigma, f) < M'(\sigma, f) < M'_1(\sigma, f^{(1)}) < M'_2(\sigma, f^{(2)}) < \dots,$$

and ([8], p.708),

$$\lambda_{>}(\sigma, f) \leq \lambda_{>}(\sigma, f^{(1)}) \leq \lambda_{>}(\sigma, f^{(2)}) \leq \dots.$$

Making use of these facts in (3.13), we get (3.1).

The proof of (3.2) is similar to that of (3.1) except that instead of (3.3) we have to use (3.4).

Corollary. Under the hypothesis of Theorem 2 and for sufficiently large  $\sigma$ ,

$$M'_n(\sigma, f^{(n)}) > M'(\sigma, f) \left( \frac{\log M'(\sigma, f)}{\left(\frac{1}{\lambda} - \frac{1}{\rho} + \varepsilon\right)(\rho + \varepsilon)\sigma} \right)^n.$$

This follows from (3.1) and the following result in [9]:

$$\limsup_{\sigma \rightarrow +\infty} \frac{\log \lambda_{>}(\sigma, f)}{\sigma} = \rho.$$

The next theorem is interesting since it shows that for a certain class of entire Dirichlet series, the functions  $M_n$  and  $M_{n+1}$  are separated from each other by the

derivative of the former for sufficiently large values of  $\sigma$ .

Theorem 3. Under the hypothesis of Theorem 2 and for sufficiently large

$$(3.14) \quad M_{\mu}(\sigma, f^{(n)}) < M'_{\mu}(\sigma, f^{(n)}) \leq M_{\mu+1}(\sigma, f^{(n+1)}), \quad \forall \mu \in \mathbb{N}.$$

Proof. It is known ([10], Lemma 2) that for every entire function  $f \in E$  of Ritt order  $\rho \in \mathbb{R}_+^* \cup \{0\}$  ( $\mathbb{R}_+^*$  is the set of extended positive reals) and lower order  $\lambda \in \mathbb{R}_+^* \cup \{0\}$ ,

$$(3.15) \quad = \lim_{\sigma \rightarrow +\infty} \frac{\sup \log(M'(\sigma, f^{(n)}))}{\inf \sigma} \bigg/ \frac{M(\sigma, f^{(n)})}{\sigma},$$

$$\forall \mu \in \mathbb{N}.$$

Hence, for any  $\varepsilon \in \mathbb{R}_+$  and sufficiently large  $\sigma$ ,

$$M_{\mu}(\sigma, f^{(n)}) e^{\sigma(\lambda-\varepsilon)} < M'_{\mu}(\sigma, f^{(n)}).$$

Since  $\varepsilon$  is arbitrary and  $\lambda \geq \sigma > 0$ , it follows that, for sufficiently large  $\sigma$ ,

$$(3.16) \quad M_{\mu}(\sigma, f^{(n)}) < M'_{\mu}(\sigma, f^{(n)}).$$

Combining (3.16) with (3.8), we get (3.14).

4. Finally, we establish a result regarding ordinary proximate linear order of entire functions in  $E$  of irregular growth. We first recall its definition.

Definition ([11], p.112): A nonnegative extended real valued function  $\Phi$  of reals  $\sigma$  is called an ordinary proximate linear order of an entire function  $f \in E$  of Ritt order  $\rho \in \mathbb{R}_+$  provided

(a)  $\Phi$  is eventually a continuous function,

(b)  $\Phi$  is differentiable almost everywhere except at the isolated points at which the left and right derivatives exist,

$$(c) \quad \lim_{\sigma \rightarrow +\infty} \sigma \Phi'(\sigma) = 0 ,$$

$$(d) \quad \lim_{\sigma \rightarrow +\infty} \sup \Phi(\sigma) = \rho , \quad \text{and}$$

$$(e) \quad \lim_{\sigma \rightarrow +\infty} \sup \frac{\log M(\sigma, f)}{e^{\sigma \Phi(\sigma)}} = 1 .$$

Theorem 4. For every entire function  $f \in E$  of irregular growth of Ritt order  $\rho \in \mathbb{R}_+$  and lower order  $\lambda \in \mathbb{R}_+$  and ordinary proximate linear order  $\Phi$  ,

$$(4.1) \quad \lim_{\sigma \rightarrow +\infty} \inf \frac{\log M_{\rho}(\sigma, f^{(\rho)})}{e^{\sigma \Phi(\sigma)}} = 0 .$$

Proof. We know that

$$(4.2) \quad \lim_{\sigma \rightarrow +\infty} \sup \inf \frac{\log M_{\rho}(\sigma, f^{(\rho)})}{\sigma} = \rho .$$

From (4.2), we get, for any  $\epsilon \in \mathbb{R}_+$  and sufficiently

large values of  $\sigma$  ,

$$(4.3) \quad \log M_n(\sigma, f^{(n)}) > e^{(\lambda-\epsilon)\sigma} ,$$

and, for an infinite sequence of  $\sigma$ 's,

$$(4.4) \quad \log M_n(\sigma, f^{(n)}) < e^{(\lambda+\epsilon)\sigma}$$

Dividing (4.3) and (4.4) by  $e^{\sigma\phi(\sigma)}$  and proceeding to limit we get (4.1) in view of the condition (d) of the definition of ordinary proximate linear order.

#### R e f e r e n c e s

- [1] MANDELBROJT S.: The Rice Institute Pamphlet (Dirichlet series), No.4, Vol.31, Houston, 1944.
- [2] DÖETSCH G.: Über die obere Grenze des absoluten Betrages einer analytischen Funktion auf Geraden, Math.Z.18(1920), 237-240.
- [3] SUGIMURA K.: Übertragung einiger Sätze aus der Theorie der ganzen Funktionen auf Dirichletsche Reihen, Math.Z.29(1929), 264-267.
- [4] GUPTA J.S.: A note on the maximum term of an entire Dirichlet series, Publ.Math.Debrecen 10(1963), 53-57.
- [5] SRIVASTAVA S.N.: A note on the derivatives of an integral function represented by Dirichlet series, Rev.Mat.Hisp.-Amer.22(1962), 246-259.
- [6] Yu C.Y.: Sur les droites de Borel de certaines fonctions entières, Ann.Sci.École Norm.Sup.68(1951), 65-104.
- [7] SRIVASTAVA R.S.L. and GUPTA J.S.: On the maximum term of an integral function defined by Dirichlet

series, *Math. Ann.* 174(1967), 240-246.

- [8] SRIVASTAVA R.S.L.: On the order of integral functions defined by Dirichlet series, *Proc. Amer. Math. Soc.* 12(1961), 702-708.
- [9] RAHMAN Q.I.: On the maximum modulus and the coefficients of an entire Dirichlet series, *Tohoku Math. J.* (2), 8(1956), 108-113.
- [10] SRIVASTAV R.P. and GHOSH R.K.: On the entire functions represented by Dirichlet series, *Ann. Polon. Math.* 13(1963), 93-100.
- [11] GUPTA J.S.: On the ordinary proximate linear order of integral functions defined by Dirichlet series, *Monatsh. Math.* 70(1963), 111-117.

University of Jammu  
Jammu, J & K  
India

(Oblatum 6.4.1973)