

E. A. Kalita

On the Hölder continuity of solutions of nonlinear parabolic systems

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 35 (1994), No. 4, 675--680

Persistent URL: <http://dml.cz/dmlcz/118708>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## On the Hölder continuity of solutions of nonlinear parabolic systems

E. KALITA

*Abstract.* Non-linear second order parabolic systems in the divergent form are considered. It is proved that under some restrictions on the modulus of ellipticity, all weak solutions are continuous.

*Keywords:* non-linear parabolic systems, weak solutions, regularity

*Classification:* 35D10, 35K55

We consider the parabolic system

$$(1) \quad \frac{\partial u}{\partial t} - \sum_{k=1}^n D_k A_k(x, t, u, Du) = 0$$

of  $N$  equations ( $N \geq 1$ ) in the domain  $Q \subset \mathbf{R}^{n+1}$ , ( $n \geq 2$ ).

Here  $u$  and  $A_k$  are vector functions of dimension  $N$ ,  $D = (D_1, D_2, \dots, D_n)$ ,  $D_k = \frac{\partial}{\partial x_k}$ . The coefficients  $A_k(x, t, u, \xi)$  are assumed to be differentiable over all arguments except  $t$ .

Further, it is supposed that

$$(2) \quad \sum_{i,k} (\eta_k^i - \kappa \sum_{j,l} A_{kl}^{ij}(x, t, u, \xi) \eta_l^j)^2 \leq (1 - \nu^2) |\eta|^2, \quad \nu > 0,$$

$$(3) \quad \begin{cases} \left| \frac{\partial A}{\partial u} \right| \leq c |\xi|^{\frac{2}{n+2}} + c |u|^{\frac{2}{n}} + f(x, t), \\ \left| \frac{\partial A}{\partial x} \right| \leq c |\xi| + c |u|^{1+\frac{2}{n}} + f^2(x, t), \end{cases}$$

where  $A_{kl}^{ij} = \frac{\partial A_k^i}{\partial \xi_l^j}$  with  $A = (A_1, A_2, \dots, A_n)$ ,  $\kappa > 0$  is a norming factor,  $f \in L_a$  with  $a > n + 2$ .

By  $c$  we denote any positive constant the value of which is of no importance.

Observe that the condition (2) follows from the pair of inequalities usually written in the form

$$(4) \quad \begin{cases} \sum_{i,j=1}^N \sum_{k,l=1}^n A_{kl}^{ij}(x, t, u, \xi) \eta_k^i \eta_l^j \geq \lambda |\eta|^2, \\ \sum_{i,k} (\sum_{j,l} A_{kl}^{ij}(x, t, u, \xi) \eta_l^j)^2 \leq \Lambda^2 |\eta|^2, \end{cases}$$

where  $\nu \geq \frac{\lambda}{\Lambda}$ . Evidently, if (4) holds, we obtain easily that

$$\sum_{i,k} (\eta_k^i - \kappa \sum_{j,l} A_{kl}^{ij} \eta_l^j)^2 = |\eta|^2 - 2\kappa \sum_{i,j,k,l} A_{kl}^{ij} \eta_l^j \eta_k^i + \kappa^2 \sum_{i,k} (\sum_{j,l} A_{kl}^{ij} \eta_l^j)^2 \leq (1 - 2\kappa\lambda + \kappa^2\Lambda^2) |\eta|^2;$$

setting  $\kappa = \frac{\lambda}{\Lambda^2}$  we obtain (2) with  $\nu = \frac{\lambda}{\Lambda}$ .

**Theorem.** *In the condition (2), let*

$$(5) \quad \nu > 1 - \frac{2}{n}.$$

*Then each generalized solution  $u \in V_{2,loc}(Q)$  of the system (1) belongs to  $C_{loc}^\alpha(Q)$  for some  $\alpha > 0$ .*

(Here  $V_{2,loc}(Q)$  is the space of measurable functions on  $Q$  with the expression

$$\sup_t \int |u(x, t)|^2 dx + \int \int |Du(x, t)|^2 dx dt$$

finite over each compact subcylinder contained in  $Q$ .)

It is a well known fact that weak solutions of nonlinear parabolic and elliptic systems can be discontinuous even if the coefficients are analytic. It is also known that this cannot take place in the case of one equation of the second order. In the course of the long-lasting effort to find conditions under which weak solutions of elliptic systems are regular it was Cordes' idea which appeared to be particularly fruitful. Cordes observed that the regularity properties of weak solutions can be substantially influenced by the modulus of ellipticity of the system. (For the further development see e.g. [1], [2].)

The success of Cordes' idea obtained in the elliptic case was not so far fully repeated in case of parabolic systems. The Hölder continuity is proved in the paper [3] by means of a modification of the technique of integral estimates with weights.

In my paper I prove this result under less restrictive assumption (5). The proof is based on Moser's iterative method.

Set

$$DuDv = \sum_{i,k} D_k u^i D_k v^i, \quad |Du|^2 = DuDu.$$

In the following lemma we give the Moser-type modification of natural energy space.

**Lemma.** Let  $u = (u^1, u^2, \dots, u^N)$  be a function which is differentiable almost everywhere. Put  $v = u|u|^s$  with  $s \in (-1, \infty)$ .

Then for  $\mu(s) = 1 - (\frac{s}{2+s})^2$  we have

$$DuDv \geq \mu^{\frac{1}{2}}(s)|Du||Dv|.$$

PROOF: Denote

$$I = |u|^s|Du|^2, \quad J = |u|^{s-2} \sum_k (\sum_i u^i D_k u^i)^2.$$

Then

$$DuDv = I + sJ, \quad |u|^{-s}|Dv|^2 = I + (2s + s^2)J.$$

As  $s > -1$  and  $I \geq J$  (which follows from Hölder's inequality), we obtain that  $DuDv \geq 0$ .

The assertion of the lemma now follows from the relation

$$(DuDv)^2 - \mu(s)|Du|^2|Dv|^2 = s^2(\frac{I}{2+s} - J)^2 \geq 0.$$

□

**Remark.** In the proof of Theorem we use this result only with  $s \in (0, \infty)$ . Its other part ( $s \in (-1, 0]$ ) is helpful when we consider the removable singularities of solutions.

PROOF OF THE THEOREM: Using finite difference operators in a standard way leads us to the fact that

$$Du \in V_{2,loc}(Q), \quad \frac{\partial u}{\partial t} \in L_{2,loc}(Q).$$

(In what follows, all the spaces are local in  $Q$ .) Further, the local variant of Gehring's lemma (see [4], appendix) yields that  $D^2u$  and  $\frac{\partial u}{\partial t}$  belong to  $L_p$  for some  $p > 2$ .

Let now  $z_o = (x_o, t_o)$  be an arbitrary point in  $Q$ ,  $\varrho \leq \frac{1}{3} \text{dist}(z_o, \partial Q)$ . Define the cut-off function

$$\varphi(z) = \psi(\frac{|x - x_o|}{\varrho} - 1)\psi(\frac{t_o - t}{\varrho})$$

where  $z = (x, t)$  and  $\psi$  is a smooth decreasing function on  $\mathbf{R}$  with  $\psi(\tau) = 1$  for  $\tau < 0$  and  $\psi(\tau) = 0$  for  $\tau > 1$ .

Set  $v_m = u_m|u_m|^s$  for  $s > 0$  where  $u_m = D_m u$ , ( $m \in (1, 2, \dots, n)$ ). Rewrite the system (1) in the form

$$\kappa \frac{\partial u^i}{\partial t} - \Delta u^i = \sum_k D_k(\kappa A_k^i - D_k u^i),$$

where  $\kappa$  is the norming factor from the condition (2). Multiplying the system by  $D_m(v_m\varphi^2)$  we integrate it over the domain

$$Q_1 = \{z = (x, t); |x - x_o| < 2\varrho \text{ and } |t - t_o| < \varrho\}.$$

As  $u \in W_p^{2,1}$  with  $p > 2$  the integral is finite for sufficiently small positive  $s$ . Integrating by parts with respect to the variable  $x$  we obtain

$$(6) \quad \int_{Q_1} \left( \frac{\kappa}{2+s} \frac{\partial}{\partial t} |u_m|^{2+s} + Du_m Dv_m \right) \varphi^2 dz = \\ \int_{Q_1} \sum_{i,k} \left\{ \left[ \kappa \sum_{j,l} A_{kl}^{ij} D_l u_m^j - D_k u_m^i \right] + \sum_j \frac{\partial A_k^i}{\partial u^j} u_m^j + \frac{\partial A_k^i}{\partial x_m} \right\} \varphi^2 D_k v_m^i + \\ \kappa D_m A_k^i v_m^i D_k \varphi^2 \} dz$$

The first member in the right-hand side of the formula (6) can be estimated by means of Hölder’s inequality and of the assumption (2) as

$$\left| \sum_{i,k} \left( \kappa \sum_{j,l} A_{kl}^{ij} D_l u_m^j - D_k u_m^i \right) D_k v_m^i \right| \leq (1 - \nu^2)^{\frac{1}{2}} |Du_m| |Dv_m|.$$

Using now the Lemma, we obtain from (6) the estimate

$$(7) \quad \max_{|t-t_o| < \varrho} \frac{\kappa}{2+s} \int_{\Omega_t} |u_m|^{2+s} \varphi^2 dx + \\ \left( \mu(s)^{\frac{1}{2}} - (1 - \nu^2)^{\frac{1}{2}} \right) \int_{Q_1} |Du_m|^2 |u_m|^s \varphi^2 dz \leq \\ c \int_{Q_1} \left[ \left( \left| \frac{\partial A}{\partial u} \right| |u_m| + \left| \frac{\partial A}{\partial x} \right| \right) |Dv_m| \varphi^2 + (|D_m A| |v_m| + |u_m|^{2+s}) \varphi \right] dz,$$

where

$$\Omega_t = \{(x, t); |x - x_o| < 2\varrho\}.$$

In the case that  $\frac{s}{2+s} < \nu$ , we obtain the same kind of estimate as (7) for the left-hand side

$$\max_{|t-t_o| < \varrho} \int_{\Omega_t} |u_m|^{2+s} \varphi^2 dx + \int_{Q_1} |Du_m|^2 |u_m|^s \varphi^2 dz.$$

(The eventual increase of the constant  $c$  does not play any important role.)

Now we apply Young’s inequality to the right-hand side of (7). Taking account of the fact that from the condition (3) and from  $Du \in V_2$  it follows  $\frac{\partial A}{\partial u} \in L_b$  with  $b = \min\{\frac{(n+2)^2}{n}, a\}$  we get

$$(8) \quad \max_{|t-t_o| < \varrho} \int_{\Omega_t} |u_m|^{2+s} \varphi^2 dx + \int_{Q_1} |Du_m|^2 |u_m|^s \varphi^2 dz \\ \leq c \int_{Q_1} \left[ (1 + |Du|)^q + |u|^{2+4/n} |Du|^s + \left| \frac{\partial A}{\partial u} \right|^b + f^a \right] dz,$$

where  $q = \max\{b\frac{2+s}{b-2}, \frac{as}{a-4}\}$ . By the restrictions  $a < \frac{(n+2)^2}{n}$  and  $s < a - 4$ , we obtain the value of  $q$  as  $q = a\frac{2+s}{a-2}$ .

As for  $u$  and  $Du$ , the condition  $Du \in V_2$  yields that  $Du \in L_{2\frac{n+2}{n}}$  and  $u \in L_{2\frac{n+2}{n-2}}$ . (In the case of  $n = 2$  the solution  $u$  is integrable with any positive power.)

Take  $s_1 \in (0, \frac{2\nu}{1-\nu})$  so small that all the considerations that we have made until now are correct for each  $s, s \leq s_1$ . As  $z_o$  was an arbitrary point of  $Q$ , the estimate (8) yields

$$|Du|^{1+s_1/2} \in V_2.$$

The imbedding theorem for the space  $V_2$  gives then

$$Du \in L_{(2+s_1)\frac{n+2}{n}}, \quad u \in L_{(2+s_1)\frac{n+2}{n-2}}$$

So all the reasonings leading to the estimate (8) can be repeated for  $s \leq s_2$ , where  $s_2 \in (0, \frac{2\nu}{1-\nu})$  satisfies the inequalities

$$\begin{aligned} s_2 &\leq (2 + s_1)\frac{n + 2}{n}\left(1 - \frac{2}{p}\right), \\ 2 + s_2 &\leq (2 + s_1)\frac{n + 2}{n} \frac{a - 2}{a}, \\ s_2 &\leq \left(s_1 + \frac{4}{n}\right)\frac{n + 2}{n}. \end{aligned}$$

As  $p > 2$  and  $a > n + 2$ , the value of  $s_2$  can be chosen in a way that  $s_2 > s_1$ . From (8) we have  $|Du|^{1+s_2/2} \in V_2$ .

Analogously, we can determine  $s_3$  starting from  $s_2$  and so on. It is evident now that after a finite number of steps we obtain that  $|Du|^{1+s/2} \in V_2$  for any  $s < s^* = \min\{\frac{2\nu}{1-\nu}, a - 4\}$ .

It follows now from (5) and from  $a > n + 2$  that  $s^* > n - 2$ . Thus for each  $t$  the function  $Du(x, t)$  is integrable with respect to  $x$  to power bigger than  $n$  and so (by the imbedding theorem)  $u$  is locally Hölder continuous with respect to  $x$  in  $Q$ .

As  $\frac{\partial u}{\partial t} \in L_2$ , it follows in the standard way that the Hölder continuity with respect to  $t$  with the exponent  $\alpha = \frac{\beta}{n+2\beta}$ . Namely, we have that

$$\begin{aligned} |u(x, t) - u(x, \tau)| &\leq |u(x, t) - u(y, t)| + |u(x, \tau) - u(y, \tau)| + |u(y, t) - u(y, \tau)| \\ &\leq c|x - y|^\beta + |u(y, t) - u(y, \tau)| \end{aligned}$$

where  $y$  is an arbitrary point of the ball  $B(x, \varrho) = \{y; |y - x| < \varrho\}$ .

Fubini's theorem now gives that

$$\int \left| \frac{\partial u}{\partial t}(y, t) \right|^2 dt < \infty$$

for almost all  $y$ , and for such  $y$  the function  $u(y, t)$  is Hölder continuous with respect to  $t$  with the exponent  $\frac{1}{2}$ :

$$|u(y, t) - u(y, \tau)| \leq c|t - \tau|^{\frac{1}{2}} \left( \int_{\tau}^t \left| \frac{\partial u}{\partial t}(y, h) \right|^2 dh \right)^{\frac{1}{2}}.$$

Evidently,

$$\min_{y \in B(x, \varrho)} \int_{\tau}^t \left| \frac{\partial u}{\partial t}(y, h) \right|^2 dh \leq |B(x, \varrho)|^{-1} \int_{B(x, \varrho)} \int_{\tau}^t \left| \frac{\partial u}{\partial t} \right|^2 dh dy \leq c\varrho^{-n}.$$

Taking  $\varrho = |t - \tau|^{\frac{1}{n+2\beta}}$ , we obtain finally  $|u(x, t) - u(x, \tau)| \leq c|t - \tau|^{\alpha}$  with  $\alpha = \frac{\beta}{n+2\beta}$ .  $\square$

**Acknowledgement.** This work was written during my stay in the Department of Mathematical Analysis of Charles University, Prague, and was supported by the grant n. 201/93/2171 of Grant Agency of Czech Republic.

#### REFERENCES

- [1] Kalita E.A., *On sharp exponent of smoothness of solutions to the second order elliptic systems* (in Russian), *Izvestija Vyshyh Uchebnyh Zavedenij. Matematika* **3** (1992), 10–17.
- [2] Koshelev A.I., Chelkak S.I., *Regularity of solutions of quasilinear elliptic systems*, Teubner, Leipzig, 1985.
- [3] Koshelev A.I., *Regularity of solutions for some quasilinear parabolic systems*, *Math. Nachr.* **162** (1993), 59–88.
- [4] Giaquinta M., Modica G., *Regularity results for some classes of higher order non linear elliptic systems*, *J. Reine Angew. Math.* **311/312**, 145–169.

INSTITUTE OF APPLIED MATHEMATICS AND MECHANICS OF THE ACADEMY OF SCIENCES,  
DONETSK, UKRAINE

(Received February 22, 1994)