

Acta Universitatis Palackianae Olomucensis. Facultas Rerum
Naturalium. Mathematica

Mihail Banaru

On six-dimensional G_1 -submanifolds of Cayley algebra

Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica, Vol. 40 (2001), No. 1, 17--21

Persistent URL: <http://dml.cz/dmlcz/120433>

Terms of use:

© Palacký University Olomouc, Faculty of Science, 2001

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>



On Six-Dimensional G_1 -Submanifolds of Cayley Algebra

MIHAIL BANARU

*Smolensk University of Humanities,
Gertsen str. 2, Smolensk, 214 025, Russia
e-mail: banaru@keytown.com*

(Received October 20, 2000)

Abstract

It is proved that six-dimensional G_1 -submanifolds of the octave algebra are $W_1 \oplus W_3$ -manifolds.

Key words: Almost Hermitian structure, G_1 -manifold, Gray–Hervella classification.

2000 Mathematics Subject Classification: 53C40

1.

The classification of the almost Hermitian structures on first order differential-geometrical invariants can be rightfully attributed to the most significant results obtained by the outstanding American mathematician Alfred Gray in collaboration with his Spanish colleague Luis M. Hervella. According to this classification all the almost Hermitian structures are divided into 16 classes. Analytical criteria for every concrete structure to belong to one or another class have been received [1].

One of the least studied Gray–Hervella classes is the G_1 (or $W_1 \oplus W_3 \oplus W_4$) class. Almost the only significant work devoted to the G_1 -manifolds is the original and interesting article by L. M. Hervella and E. Vidal [2]. This situation is at least strange because the G_1 -manifolds class includes the Kählerian (K), the nearly-Kählerian (NK), the Hermitian (H), the special Hermitian (SH) and the locally conformal Kählerian (LCK) manifolds as well as the Vaisman–Gray manifolds. And what is more, the G_1 -manifolds class forms a minimal class that includes two of the most interesting classes of almost Hermitian manifolds—the

nearly-Kählerian and Hermitian manifolds classes, to the study of which a great number of publications have been devoted. Without going into details on such an extensive subject, we can mark out the classical works [3]–[8].

In the present article we give some results obtained in this direction by using the Cartan structure equations of the octave algebra submanifold.

2.

Let $\mathbf{O} \equiv \mathbf{R}^8$ be the Cayley algebra. As it is well-known [3], two non-isomorphic 3-vector cross products are defined on it by

$$\begin{aligned} P_1(X, Y, Z) &= -X(\overline{Y}Z) + \langle X, Y \rangle Z + \langle Y, Z \rangle X - \langle Z, X \rangle Y, \\ P_2(X, Y, Z) &= -(X\overline{Y})Z + \langle X, Y \rangle Z + \langle Y, Z \rangle X - \langle Z, X \rangle Y, \end{aligned}$$

where $X, Y, Z \in \mathbf{O}$, $\langle \cdot, \cdot \rangle$ is the scalar product in \mathbf{O} , $X \rightarrow \overline{X}$ is the operator of conjugation [9]. Moreover, any other 3-vector cross product in the octave algebra is isomorphic to one of the above-mentioned [3].

If $M^6 \subset \mathbf{O}$ is a six-dimensional oriented submanifold, then the induced almost Hermitian structure $\{J_\alpha, g = \langle \cdot, \cdot \rangle\}$ is determined by the relation

$$J_\alpha(X) = P_\alpha(X, e_1, e_2), \quad \alpha = 1, 2,$$

where $\{e_1, e_2\}$ is an arbitrary orthonormal basis of the normal space of M^6 at a point p , $X \in T_p(M^6)$. The submanifold $M^6 \subset \mathbf{O}$ is called G1-submanifold if for arbitrary vector fields $X, Y \in \mathfrak{N}(M^6)$ the following condition is fulfilled

$$\nabla_x(F)(X, Y) - \nabla_{JX}(F)(JX, Y) = 0. \quad (1)$$

Here $F(X, Y) = \langle X, JY \rangle$ is the fundamental (or Kählerian [8]) form of M^6 , ∇ is the Riemannian connection of the metric. We note that A. Gray and L. M. Hervella have proved that the condition (1) is equivalent to the following relation imposed on the Nijenhuis tensor S [1]:

$$\langle S(X, Y), X \rangle = 0.$$

We recall that the point $p \in M^6$ is called general [6], if

$$e_0 \notin T_p(M^6) \quad \text{and} \quad T_p(M^6) \subseteq L(e_0)^\perp,$$

where e_0 is the unit of Cayley algebra and $L(e_0)^\perp$ is its orthogonal supplement. A submanifold $M^6 \subset \mathbf{O}$ consisting only of general points is called a general-type submanifold. In what follows all the considered M^6 are meant as general-type submanifolds.

3.

We use the Cartan structure equations of an almost Hermitian $M^6 \subset \mathbf{O}$ obtained in [7]:

$$d\omega^a = \omega_b^a \wedge \omega^b + \frac{1}{\sqrt{2}} \varepsilon^{ah[b} D_h^c] \omega_b \wedge \omega_c + \frac{1}{\sqrt{2}} \varepsilon^{abh} D_{hc} \omega^c \wedge \omega_b;$$

$$d\omega_a = -\omega_a^b \wedge \omega_b + \frac{1}{\sqrt{2}} \varepsilon_{ah[b} D^h_{c]} \omega^b \wedge \omega^c + \frac{1}{\sqrt{2}} \varepsilon_{abh} D^{hc} \omega_c \wedge \omega^b;$$

$$d\omega_b^a = \omega_c^a \wedge \omega_b^c - \left(\frac{1}{2} \delta_{bg}^{ah} D_{h[k} D^g_{j]} + \sum_{\varphi} T_{\hat{a}[k}^{\varphi} T_{j]b}^{\varphi} \right) \omega^k \wedge \omega^j.$$

Here $\varepsilon_{abc} = \varepsilon_{abc}^{123}$, $\varepsilon^{abc} = \varepsilon_{123}^{abc}$ are components of the Kronecker tensor of the third order [10], $\delta_{bg}^{ah} = \delta_b^a \delta_g^h - \delta_g^a \delta_b^h$,

$$D^{ch} = D_{\hat{c}\hat{h}}, \quad D_h^c = D_{h\hat{c}}, \quad D^h_c = D_{\hat{h}c},$$

$$D_{cj} = \pm T_{cj}^8 + iT_{cj}^7, \quad D_{\hat{c}\hat{j}} = \pm T_{\hat{c}\hat{j}}^8 - iT_{\hat{c}\hat{j}}^7,$$

where T_{kj}^{φ} are components of the configuration tensor (in A. Gray's notation [4], or the Euler curvature tensor [11]), $i = \sqrt{-1}$. Here and further $a, b, c, d, g, h = 1, 2, 3$; $\hat{a} = a + 3$; $\varphi = 7, 8$; $k, j = 1, 2, 3, 4, 5, 6$.

The Gray–Hervella condition (1) on almost Hermitian structures belonging to the $G1$ ($= W_1 \oplus W_3 \oplus W_4$) class is equivalent to the skew-symmetry of the Kirichenko structure tensor on the first pair of indexes [12]:

$$B^{abc} = -B^{bac}, \quad B_{abc} = -B_{bac}.$$

We use the Kirichenko tensor relations for six-dimensional almost Hermitian submanifolds of Cayley algebra [12], [13]:

$$B^{abc} = \frac{1}{\sqrt{2}} \varepsilon^{ah[b} D_h^c], \quad B_{abc} = \frac{1}{\sqrt{2}} \varepsilon_{ah[b} D^h_{c]}.$$

Hence, the following equalities are a criterion for an arbitrary six-dimensional submanifold of Cayley algebra to belong to the $G1$ -class.

$$\varepsilon^{ah[b} D_h^c] = \varepsilon^{bh[a} D_h^c], \quad \varepsilon_{ah[b} D^h_{c]} = \varepsilon_{bh[a} D^h_{c]}. \quad (2)$$

Lemma *The condition (2) is equivalent to the fact that the matrix $(D_{d\hat{a}})$ is scalar, i.e.*

$$D_{d\hat{a}} = \frac{1}{3} \delta_d^a \operatorname{tr} D_{d\hat{a}}.$$

Proof First we note that every equality from (2) can be obtained by the complex conjugation of another one. It is enough to adduce the following arguments.

1. Let (2) be fulfilled:

$$\varepsilon^{ah[b} D_h^c] = \varepsilon^{bh[a} D_h^c] \Leftrightarrow$$

$$\varepsilon^{ah[b} D_h^c] - \varepsilon^{bh[a} D_h^c] = 0 \Leftrightarrow$$

$$\varepsilon^{ahb} D_h^c - \varepsilon^{ahc} D_h^b + \varepsilon^{bha} D_h^c - \varepsilon^{bhc} D_h^a = 0 \Leftrightarrow$$

$$\varepsilon^{ahc} D_{h\hat{b}} + \varepsilon^{bhc} D_{h\hat{a}} = 0.$$

We reduce this equality (ε_{cfd}):

$$\begin{aligned} \delta_{fd}^{ah} D_{hb} + \delta_{fd}^{bh} D_{ha} &= 0 \Leftrightarrow \\ (\delta_f^a \delta_d^h - \delta_d^a \delta_f^h) D_{hb} + (\delta_f^b \delta_d^h - \delta_d^b \delta_f^h) D_{ha} &= 0 \Leftrightarrow \\ \delta_f^a D_{db} - \delta_d^a D_{fb} + \delta_f^b D_{da} - \delta_d^b D_{fa} &= 0, \\ (bf) : D_{da} - \delta_d^a \operatorname{tr} D_{da} + 3D_{da} - D_{da} &= 0 \Leftrightarrow \\ D_{da} &= \frac{1}{3} \delta_d^a \operatorname{tr} D_{da}. \end{aligned}$$

Thus, (2) $\Rightarrow D_{da} = \frac{1}{3} \delta_d^a \operatorname{tr} D_{da}$.

2. We prove the inverse statement. Let $D_{da} = \frac{1}{3} \delta_d^a \operatorname{tr} D_{da}$. Then

$$\begin{aligned} \varepsilon^{ah[b} D_h^{c]} - \varepsilon^{bh[a} D_h^{c]} &= \varepsilon^{ahc} D_{hb} + \varepsilon^{bhc} D_{ha} \\ &= \frac{1}{3} (\varepsilon^{ahc} \delta_h^b \operatorname{tr} D_{da} + \varepsilon^{bhc} \delta_h^a \operatorname{tr} D_{da}) = \frac{1}{3} (\varepsilon^{abc} \operatorname{tr} D_{da} + \varepsilon^{bac} \operatorname{tr} D_{da}) \\ &= \frac{1}{3} \operatorname{tr} D_{da} (\varepsilon^{abc} + \varepsilon^{bac}) = \frac{1}{3} \operatorname{tr} D_{da} (\varepsilon^{abc} - \varepsilon^{abc}) = 0. \end{aligned}$$

Taking into account the remark at the beginning of the proof, we come to the conclusion

$$D_{da} = \frac{1}{3} \delta_d^a \operatorname{tr} D_{da} \Rightarrow (2).$$

As a result we have that the conditions (2) and $D_{da} = \frac{1}{3} \delta_d^a \operatorname{tr} D_{da}$ are equivalent. \square

It is interesting to note that the condition for the matrix (D_{da}) to be scalar is a criterion for an almost Hermitian submanifold of Cayley algebra to belong to the $W_1 \oplus W_3$ -manifolds class [12], [13].

Thus is proved

Theorem *Every six-dimensional G_1 -submanifold of Cayley algebra is a manifold of the $W_1 \oplus W_3$ -class.*

As the almost Hermitian structures of the $W_1 \oplus W_3$ -class are semi-Kählerian (SK -structures) [1], [12] we get an important

Corollary 1 *Every six-dimensional G_1 -submanifold of Cayley algebra is a semi-Kählerian manifold.*

We remark that the Hermitian (H , or $W_3 \oplus W_4$) structures on the six-dimensional submanifolds of Cayley algebra can be represented only by the structures belonging to the $H \cap SK$ -class:

$$H = W_3 \oplus W_4, \quad SK = W_1 \oplus W_2 \oplus W_3,$$

$$H \cap SK = W_3.$$

Corollary 2 *Every six-dimensional Hermitian submanifold of Cayley algebra is a special Hermitian (W_3) manifold.*

Similar considerations will lead us to the correctness of two other statements.

Corollary 3 *Every six-dimensional Vaisman-Gray submanifold of Cayley algebra is a nearly-Kählerian manifold.*

Corollary 4 *Every six-dimensional locally conformal Kählerian submanifold of Cayley algebra is a Kählerian manifold.*

References

- [1] Gray, A., Hervella, L. M.: *The sixteen classes of almost Hermitian manifolds and their linear invariant.* Ann. Mat., Pure ed Appl. **123**, 4 (1980), 35–58.
- [2] Hervella, L. M., Vidal, E.: *Novelles géométries pseudo-Kählériennes G_1 et G_2 .* C. R. Acad. Sci. Paris **283** (1976), 115–118.
- [3] Gray, A.: *Vector cross products on manifolds.* Trans. Amer. Math. Soc. **141** (1969), 465–504.
- [4] Gray, A.: *Some examples of almost Hermitian manifolds.* Ill. J. Math. **10**, 2 (1966), 353–366.
- [5] Vaisman, I.: *On locally conformal almost Kähler manifolds.* Israel J. Math. **24** (1976), 338–351.
- [6] Kirichenko, V. F.: *On nearly-Kählerian structures induced by means of 3-vector cross products on six-dimensional submanifolds of Cayley algebra.* Vestnik MGU **3** (1973), 70–75.
- [7] Kirichenko, V. F.: *Classification of Kählerian structures induced by means of 3-vector cross products on six-dimensional submanifolds of Cayley algebra.* Izvestia Vuzov, Kazan, **8** (1980), 32–38.
- [8] Kirichenko, V. F.: *Rigidity of almost Hermitian structures induced by means of 3-vector cross products on six-dimensional submanifolds of Cayley algebra.* Ukrain Geom. Coll. **25** (1982), 60–68.
- [9] Freudenthal, H.: *Octaves, singular groups and octaves geometry.* Coll. Math., Moscow, 1957, 117–153.
- [10] Lichnerovicz, A.: *Théorie globale des connexions et des groupes d’holonomie.* Cremonese, Roma, 1955.
- [11] Cartan, E.: *Riemannian geometry in an orthogonal frame.* MGU, Moscow, 1960.
- [12] Banaru, M.: *Hermitian geometry of six-dimensional submanifolds of Cayley algebra.* MSPU, Moscow, 1993.
- [13] Banaru, M.: *On almost Hermitian structures induced by 3-vector cross products on six-dimensional submanifolds of Cayley algebra.* Polyanalytical Functions, Smolensk, 1997, 113–117.