

Aplikace matematiky

Summaries of Papers Appearing in this Issue

Aplikace matematiky, Vol. 18 (1973), No. 5, (384a)

Persistent URL: <http://dml.cz/dmlcz/103492>

Terms of use:

© Institute of Mathematics AS CR, 1973

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SUMMARIES OF PAPERS APPEARING IN THIS ISSUE

(These summaries may be reproduced)

JÁN LOVIŠEK, Bratislava: *Operator of a thin plate reinforced with thin-walled ribs.* Apl. mat. 18 (1973), 305—314. (Original paper.)

In the paper the solution of a thin plate reinforced with stiffening ribs is defined. A differential operator $\tilde{\Delta}^2$ (operator of a thin plate reinforced with thin-walled ribs) is introduced. For this operator properties of symmetry as well as of positive definiteness are proved on a linear subset $M \subset W_2^{(2)}(\Omega)$.

MIROSLAV ŠÍSLER, Praha: *Bemerkungen zur Optimierung eines gewissen Iterationsverfahrens.* Apl. mat. 18 (1973), 315—324. (Originalartikel.)

Die Arbeit bafasst sich mit der Optimierung der Konvergenzgeschwindigkeit eines gewissen einparametrischen Iterationsverfahrens für die Lösung eines linearen Gleichungssystems mit der zyklischen Matrix.

MIROSLAV ŠÍSLER, Praha: *Über ein zweiparametrisches Iterationsverfahren.* Apl. mat. 18 (1973), 325—332. (Originalartikel.)

Die Arbeit befassst sich mit einem gewissen allgemeinen Iterationsverfahren für die Lösung eines linearen Gleichungssystems mit der zyklischen Matrix. Die untersuchte Methode verallgemeinert einige bekannte Iterationsmethoden, wie z.B. Jacobi-Verfahren, Gauss-Seidelschen Verfahren, Oberrelaxationsverfahren usw.

Zbyněk ŠIDÁK, Praha: *Tables for two normal-scores rank tests for the two-sample location problem.* Apl. mat. 18 (1973), 333—345. (Original paper.)

The paper presents the tables of scores and of critical values for the Fisher - Yates - Terry - Hoeffding test and for the van der Waerden test for cases when the pooled sample size $m + n$ lies within the bounds $6 \leq m + n \leq 20$ and the one-sided significance level lies near 0.5%, 1%, 2.5%, 5%.

Zbyněk ŠIDÁK, Praha: *Tables for two normal-scores rank tests for the two-sample scale problem.* Apl. mat. 18 (1973), 346—363. (Original paper.)

The paper presents the tables of scores and of upper-tail and lower-tail critical values for the Capon test and for the Klotz test for cases when the pooled sample size $m + n$ lies within the bounds $6 \leq m + n \leq 20$ and the one-sided significance levels lie near 0.5%, 1%, 2.5%, 5%.

Zbyněk ŠIDÁK, Praha: *Tables for the two-sample Savage rank test optimal for exponential densities.* Apl. mat. 18 (1973), 364—374. (Original paper.)

The paper presents the tables of scores and of upper-tail and lower-tail critical values for the so called Savage test for cases when the pooled sample size $m + n$ satisfies $6 \leq m + n \leq 20$ and the one-sided significance levels lie near 0.5%, 1%, 2.5%, 5%.