

Juhani Nieminen

On normality relation and its generalization on lattices

Commentationes Mathematicae Universitatis Carolinae, Vol. 17 (1976), No. 4, 615--631

Persistent URL: <http://dml.cz/dmlcz/105723>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1976

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON NORMALITY RELATION AND ITS GENERALIZATION ON LATTICES

Juhani NIEMINEN, Helsinki

Abstract: Normality relation and its generalization are on a lattice L binary, unsymmetric and reflexive relations with restricted substitution properties. The lattices of these relations are considered in the case where L is a finite lattice, and a decomposition theorem is proved.

Key words: Finite lattices, normality relations, generalizations, the lattice of relations, decomposition.

AMS: Primary 06235

Ref. Ž.: 2.724.31

1. Preliminaries and introduction. A binary relation N on a lattice L is called a normality relation on L , if it satisfies the following conditions of Dean and Kruse (see Beran [1]):

(DK0) aNa for each $a \in L$.(DK1) $aNb \implies a \leq b$.(DK2) $(aNb \text{ and } cNd) \implies a \wedge cNb \wedge d$.(DK3) $(aNb \text{ and } aNc) \implies aNb \vee c$.(DK4) $(aNb \text{ and } cNd) \implies a \vee cNa \vee c \vee (b \wedge d)$.(DK5) $\{a \leq b \text{ and } (aNa \vee c \text{ or } cNa \vee c)\} \implies a \vee (b \wedge c) = b \wedge (a \vee c)$.

We shall call a binary relation on L satisfying the conditions (DK0) - (DK3) a generalized normality relation.

As one can easily see, normality and generalized normality relations on a lattice are unsymmetric generalizations

of lattice congruences and lattice tolerances (see e.g. Zelinka and Chajda [2]). The purpose of this paper is to determine a few properties of the lattice $N(L)$ of all normality relations and of the lattice $GN(L)$ of all generalized normality relations on a finite lattice L . It will be shown that in a class of finite distributive lattices, a lattice of this class is directly decomposable if and only if there are two non-trivial generalized normality relations GK and GM on L such that $GK \vee GM = 1$ and $GK \wedge GM = 0$ in the lattice $GN(L)$.

The condition (DK5) is a restricted modularity condition, and hence it is valid in each modular lattice.

As a general reference in lattice theory we have used the monograph [4] of G. Szász. The few terms of graph theory of this paper can be found in the book [3] of F. Harary.

2. Joins and meets of relations. At first we give a characterization of normality relations in terms of sublattices of a finite modular lattice.

Let L be a finite lattice. We denote by $\mathcal{A} = \{A_t \mid t \in T\}$ a family of convex sublattices of L , where T is a set of indices, and by 0_t and 1_t the least and greatest elements of A_t , respectively. Further, we assume that for each $x \in L$ there is a sublattice $A_t \in \mathcal{A}$ such that $x = 0_t$.

Theorem 1. Let L be a finite modular lattice. Each family \mathcal{A} of convex sublattices of L determine a normality relation on L and conversely, each N determines such a family if and only if for any two indices $s, u \in T$ there exist

indices $p, r \in T$ such that

$$(i) \quad 0_s \wedge 0_u = 0_p \text{ and } 1_s \wedge 1_u \leq 1_p,$$

$$(ii) \quad 0_s \vee 0_u = 0_r \text{ and } 0_s \vee 0_u \vee (1_s \wedge 1_u) \leq 1_r.$$

Proof. 1° : Let \mathcal{A} be a family with properties given in the theorem. We define a binary antisymmetric relation on L given by \mathcal{A} as follows:

$$0_s R x \iff x \in A_s \in \mathcal{A}.$$

We show that R is a normality relation on L .

aRa for each $a \in L$, as for each $a \in L$ there was a sublattice $A_t \in \mathcal{A}$ such that $0_t = a$, and so (DK0) holds. (DK1) follows directly from the definition of R .

(DK2): Let aRb and cRd . According to the definition $a = 0_s$ and $c = 0_u$ for some indices $u, s \in T$. Further, $a \wedge c = 0_s \wedge 0_u = 0_p$ and $0_p \leq b \wedge d \leq 1_s \wedge 1_u \leq 1_p$ for some $p \in T$, and thus the definition of R implies $0_p R b \wedge d$.

(DK3): Let aRb and aRc , i.e. $a, b, c \in A_t$ for some $t \in T$. As A_t is a sublattice of L , $b \vee c \in A_t$, and so $aRb \vee c$. The proof of (DK4) is similar to that of (DK2), and (DK5) holds, as L is modular.

2° : Let N be a given normality relation on L . We shall show that N generates a family \mathcal{F} of convex sublattices of L having the same properties as \mathcal{A} in the theorem. Let $F_x = \{y \mid xNy, y \in L\}$ for each $x \in L$, and we denote $\mathcal{F} = \{F_x \mid x \in L\}$.

As xNx holds for each $x \in L$, there is, according to (DK1), for each $x \in L$ a set $F_x \in \mathcal{F}$ such that x is the least element of F_x . As F_x is finite, there exists an element $w = \bigvee \{y \mid y \in F_x\}$, and according to (DK3), xNw . For each

$v \in [x, w] \subseteq L$ it holds vNv . By applying (DK2) to xNw and vNv , we obtain xNv . Hence $F_x = [x, w]$, which is a convex sublattice of L .

Let xNy and zNv . According to (DK2), $x \wedge zNy \wedge v$, and on the other hand $F_{x \wedge y} \in \mathcal{F}$. As $x \wedge zNy \wedge v$, then $y \wedge v \leq 1_{x \wedge z}$, and so (i) holds. (ii) follows similarly from (DK4), and (DK5) holds, as L is modular. This completes the proof.

The following corollary follows immediately from the proof above.

Corollary. Let L be a finite lattice. Each family \mathcal{A} of convex sublattices of L determines a generalized normality relation GN on L and conversely, GN determines such a family if and only if for any two indices $s, u \in T$ there exists an index $p \in T$ such that (i) of Theorem 1 holds.

In the following we look for meets and joins of two generalized normality relations (normality relations). The assertion of the following lemma is obviously valid.

Lemma 1. Let L be a finite lattice and GN and GR two generalized normality relations on L . The relation K , where $aKb \iff \{aGNb \text{ and } aGRb\}$ is a generalized normality relation on L and $K = GN \wedge GR$.

Analogous lemma holds also for normality relations.

If GM is a generalized normality relation on a finite lattice L we denote the corresponding family of intervals of L by $\mathcal{A}(GM)$, an interval of $\mathcal{A}(GM)$ with the least element $x \in L$ by A_{GMx} and the greatest element of A_{GMx} by l_{GMx} . The following theorem gives the most simple join of two generalized normality relations.

Theorem 2. Let GM and GN be two generalized normality relations on a finite distributive lattice L. The family

\mathcal{A} (GH), where $A_{GMx} = [x, l_{GMx} \vee l_{GNx}]$, determines a generalized normality relation on L and $GH = GM \vee GN$ if and only if

(i) $L = L_1 \times L_2 \times \dots \times L_m$, where L_i is a chain, $i = 1, \dots, m$,
or

(ii) L can be divided into two convex sublattices L^* and L^{**} such that $L^* \wedge L^{**}$ contains only one element, which is 0 of L^* and 1 of L^{**} , L^{**} is a chain and L^* satisfies the condition (i) above.

Proof. 1^0 : Let L satisfy (i) of the theorem; it is sufficient to show the validity of (DK2) - the conditions (DK0), (DK1) and (DK3) hold obviously.

Let $aGhb$ and $cGHd$; we shall show that $d \wedge b \leq (l_{GMa} \vee l_{GNa}) \wedge (l_{GMc} \vee l_{GNc}) \leq l_{GMa \wedge c} \vee l_{GNa \wedge c}$. At first, by applying the distributivity, $(l_{GMa} \vee l_{GNa}) \wedge (l_{GMc} \vee l_{GNc}) = (l_{GMa} \wedge l_{GMc}) \vee (l_{GNa} \wedge l_{GNc}) \vee (l_{GMa} \wedge l_{GNc}) \vee (l_{GNa} \wedge l_{GMc})$, where $l_{GMa} \wedge l_{GMc} \leq l_{GMa \wedge c}$ and $l_{GNa} \wedge l_{GNc} \leq l_{GNa \wedge c}$, as GM and GN are generalized normality relations on L. In the following we consider the term $l_{GMa} \wedge l_{GNc}$ and show that it is equal to or less than $l_{GNa \wedge c} \vee l_{GMa \wedge c}$; the proof is similar for $l_{GMc} \wedge l_{GNa}$.

As $L = L_1 \times \dots \times L_m$, $a = (a_1, a_2, \dots, a_m)$, $c = (c_1, \dots, c_m)$, $l_{GMa} = (x_1, \dots, x_m)$ and $l_{GNc} = (y_1, \dots, y_m)$, where $a_i, c_i, x_i, y_i \in L_i$. As $aGML_{GMa}$ and $cGNL_{GNc}$, we obtain $(a_1, \dots, a_i, \dots, a_m)GM(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_m)$ and $(c_1, \dots, c_i, \dots, c_m)GN(c_1, \dots, c_{i-1}, y_i, c_{i+1}, \dots, c_m)$. Furthermore, as L_i is a chain, $a_i \leq c_i$ or $c_i \leq a_i$, and we assume that

$a_i \leq c_i$, i.e. $a_i \wedge c_i = a_i$, and $x_i \wedge y_i \leq x_i$ holds always. But then $(a_1, \dots, a_m)GM(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_m)$ implies $(a_1, \dots, a_{i-1}, a_i \wedge c_i, a_{i+1}, \dots, a_m)GM(a_1, \dots, a_{i-1}, x_{i-1}, x_i \wedge y_i, a_{i+1}, \dots, a_m)$. According to the properties (DKO) and (DK2) of GM, we can now form the meet of both sides with $(c_1, \dots, c_{i-1}, y_i, c_{i+1}, \dots, c_m)$, and we obtain $(a_1 \wedge c_1, \dots, a_m \wedge c_m)GM(a_1 \wedge c_1, \dots, a_{i-1} \wedge c_{i-1}, x_i \wedge y_i, a_{i+1} \wedge c_{i+1}, \dots, a_m \wedge c_m)$ as $c_i \leq y_i$. So, in general, for each i , $(a_1 \wedge c_1, \dots, a_m \wedge c_m)GT(a_1 \wedge c_1, \dots, a_{i-1} \wedge c_{i-1}, x_i \wedge y_i, a_{i+1} \wedge c_{i+1}, \dots, a_m \wedge c_m)$, where GT is GM or GN, $i = 1, \dots, m$. Let z be the join of all elements $(a_1 \wedge c_1, \dots, a_{i-1} \wedge c_{i-1}, x_i \wedge y_i, a_{i+1} \wedge c_{i+1}, \dots, a_m \wedge c_m)$ which are in the relation GN with $(a_1 \wedge c_1, \dots, a_m \wedge c_m)$ for some value of i , and let the corresponding join be w in the case of GM; these joins exist according to (DK3). As GM and GN are generalized normality relations and $a \wedge cGMw$ and $a \wedge cGNz$, $w \leq l_{GMa \wedge c}$ and $z \leq l_{GNa \wedge c}$, and trivially, $w \vee z = (x_1 \wedge y_1, \dots, x_m \wedge y_m) = l_{GMa} \wedge l_{GNc}$, where $w \vee z \leq l_{GMa \wedge c} \vee l_{GNa \wedge c}$. As mentioned above, we can similarly see that $l_{GMc} \wedge l_{GNa} \leq l_{GMa \wedge c} \vee l_{GNa \wedge c}$.

As each term of the join $(l_{GMa} \wedge l_{GMc}) \vee (l_{GNa} \wedge l_{GNc}) \vee (l_{GMa} \wedge l_{GNc}) \vee (l_{GMc} \wedge l_{GNa})$ is less or equal to $l_{GMa \wedge c} \vee l_{GNa \wedge c}$, the join satisfies this relation as well. Hence $(l_{GMa} \vee l_{GNa}) \wedge (l_{GMc} \wedge l_{GNc}) \leq l_{GMa \wedge c} \vee l_{GNa \wedge c}$.

The proof for the lattice L satisfying (ii) is a repetition of the proof above, and hence we will omit it. For completing the proof of necessity we must show that $GH = GM \vee GN$. Let $GK \geq GM, GN$, and so for each $x \in L$, $xGKl_{GMx}$ and $xGKl_{GNx}$. According to (DK3), $xGK(l_{GMx} \vee l_{GNx})$, whence $GK \geq GH$,

and thus $GH = GM \vee GN$.

2°: Let GH be the join of relations GM and GN on L , and $A_{GHx} = [x, l_{GMx} \vee l_{GNx}]$. Let us remove from the Hasse diagram of L all the points and the lines incident to those points, which are meet-reducible in L . Remove further the chain C_0 containing the zero element of L , if such a chain exists. If the diagram graph thus obtained is empty, L was the chain C_0 , and the theorem holds. If not, let us consider the graph D obtained. If it is a tree, where the degree of point 1 only can be 3 or greater, then there is nothing to prove: the chains of this tree are the factors L_1, \dots, L_m in (i), as the elements of a finite distributive lattice can be uniquely represented as meets of meet-irreducibles.

Assume that D is a tree and there is a point $a \neq 1$ with the degree at least 3. Then there are in D two points x and y which are meet-irreducible in L . Let us consider the sublattice of elements $\{x \wedge y, x, y, a, z\}$ of L , where $z \in D$, and $a < q < z$ holds for no $q \in L$ (e.g. $a < z$); such an element z exists in L as D is a tree and $a \neq 1$ (see Fig. 1(a)). We define




Figure 1

a generalized normality relation GM as follows: $rGMs \iff r = s$ or $\exists q \in L$ such that $r = y \wedge q$ and $s \leq z \wedge q$; obviously GM is a generalized normality relation on L. We define another relation FN analogously: $tGNu \iff t = u$ or $\exists p \in L$ such that $t = p \wedge x$ and $u \leq z \wedge p$. One can easily see that $[x \wedge y, l_{GMx \wedge y} \vee l_{GNx \wedge y}] = [x \wedge y, a]$, but it holds for each $GK \geq GM, GN$ that $xGKz$ and $yGKz$, whence $x \wedge yGKz$, as well. But $z \notin [x \wedge y, a]$, which is a contradiction. So in the tree D only the point 1 can have degree 3 or greater.

Assume that D is unconnected graph. Let x be the point of D such that $x \neq 1$, but all the points h_1, \dots, h_{n_x} which are joined by a line to x in D are less than x in L. As the chain C_0 has been removed, there are in L also elements that are less than x. On the other hand, as $x \neq 1$, there is also a meet-reducible element a in L satisfying $x \prec a$, and let the shortest meet-representation of a in terms of meet-irreducibles contain an element $z \in L$. As the chain C_0 has been removed, there is in L an element y such that $y \vee x = a$, or there are two non-comparable elements $u, y \not\leq x$ such that $x = u \vee y$ (see Figures 1(b) and 1(c)).

In the case of Figure 1(b) we define two generalized normality relations GM and GN as in the case above. There are not two non-comparable elements $b \geq x$ and $c \geq y$ such that $b \vee c = z$ and $b \wedge c = x \wedge y$, as in the other case $b \wedge a = x$, because $b \wedge c = x \wedge y$, $a \succ x$, $a \geq y$ and $c \geq y$. Hence $z \notin [x \wedge y, l_{GMx \wedge y} \vee l_{GNx \wedge y}]$, and we get the desired contradiction.

In the case of Figure 1(c), the relations GM and GN can be defined as follows: $rGMs \iff r = s$ or $\exists p \in L$ such

that $u \wedge p = r$ and $a \wedge p \geq s$, and $tGNv \iff t = v$ or $\exists f \in L$ such that $f \wedge y = t$ and $f \wedge a \geq v$. The assumption in the case of Figure 1(c) says that there are not two non-comparable elements $b \geq u$ and $c \geq y$ such that $b \vee c = a$ and $b \wedge c = u \wedge y$, as in the other case $b \vee x = a$ or $c \vee x = a$. Hence $a \notin [u \wedge y, l_{GMu \wedge y} \vee l_{GNu \wedge y}]$. So D must be a connected tree, where only the point 1 can have the degree 3 or greater. This completes the proof.

The following lemma gives a join construction for generalized normality relations in the general case.

Lemma 2. Let GM and GN be two generalized normality relations on a finite lattice L . Then the family $\mathcal{A}(GH) = \{[a, l_{GMa} \vee l_{GNa} \vee U_a] \mid a \in L\}$, where $U_a = \bigvee_{S_a} \{(l_{GMx} \vee l_{GNx} \vee U_x) \wedge (l_{GM_y} \vee l_{GN_y} \vee U_y) \mid S_a \text{ is the set of all pairs } x, y \in L \text{ for which } x \wedge y = a\}$, generates a generalized normality relation GH on L and $GH = GM \vee GN$.

Proof. As $U_{a \wedge c}$ contains at least the term $(l_{GMa} \vee l_{GNa} \vee U_a) \wedge (l_{GMc} \vee l_{GNc} \vee U_c)$, then $b \wedge d \in [a \wedge c, l_{GMa \wedge c} \vee l_{GNa \wedge c} \vee U_{a \wedge c}]$ and (DK2) holds for $aGHb$ and $cGHd$. The other conditions hold obviously.

Let GP be a generalized normality relation on L such that $GP \geq GM, GN$. Then $xGP l_{GMx}$ and $xGP l_{GNx}$ for each $x \in L$, and so $xGP(l_{GMx} \vee l_{GNx})$, as well. According to the property (DK2) and to the finiteness of L , also $xGPU_x$. Hence $xGP(l_{GMx} \vee l_{GNx} \vee U_x)$ for each $x \in L$, and thus $GP \geq GH$. Consequently, $GH = GM \vee GN$, and the lemma follows.

The following lemma gives a construction for the join of normality relations analogous to the results in Theorem 2.

Lemma 3. Let M and N be two normality relations on a finite distributive lattice L . The family $\mathcal{A}(H) = \{[a, l_{Ma} \vee l_{Na} \vee w_a] \mid a \in L\}$, where $w_a = \bigvee_{S_a} \{ (l_{Mx_1} \vee l_{Nx_1}) \wedge (l_{Mx_2} \vee l_{Nx_2}) \wedge \dots \wedge (l_{Mx_n} \vee l_{Nx_n}) \mid S_a \text{ is the set of all sequences } x_1, \dots, x_n \text{ for which } a = x_1 \vee x_2 \vee \dots \vee x_n, n \geq 2 \}$, generates a normality relation H on L and $H = N \vee M$, if $L = I_1 \times I_2 \times \dots \times I_n$, where I_i is a chain for each value of $i = 1, \dots, n$.

Proof. Let us consider first the condition (DK4). Let aHb and cHd ; we must show that $a \vee c \vee (b \wedge d) \leq a \vee c \vee \{ (l_{Ma} \vee l_{Na} \vee w_a) \wedge (l_{Mc} \vee l_{Nc} \vee w_c) \} \in [a \vee c, l_{Mave} \vee l_{Nave} \vee w_{a \vee c}]$. By applying the distributivity we see that $(l_{Ma} \vee l_{Na} \vee w_a) \wedge (l_{Mc} \vee l_{Nc} \vee w_c) = \{ (l_{Ma} \vee l_{Na}) \wedge (l_{Mc} \vee l_{Nc}) \} \vee \{ w_a \wedge (l_{Mc} \vee l_{Nc}) \} \vee \{ w_c \wedge (l_{Ma} \vee l_{Na}) \} \vee \{ w_a \wedge w_c \} \leq w_{a \vee c}$ according to the definition of $w_{a \vee c}$. As $a \vee c \leq l_{Mave} \vee l_{Nave}$, the assertion follows by combining these two observations.

(DK0), (DK1) and (DK3) hold obviously, and so we shall consider the condition (DK2) only. Let aHb and cHd . The relation H satisfies (DK2), if $b \wedge d \leq (l_{Ma} \vee l_{Na} \vee w_a) \wedge (l_{Mc} \vee l_{Nc} \vee w_c) \in [a \wedge c, l_{Ma \wedge c} \vee l_{Na \wedge c} \vee w_{a \wedge c}]$. As above, we consider the term $\{ (l_{Ma} \vee l_{Na}) \wedge (l_{Mc} \vee l_{Nc}) \} \vee \{ w_a \wedge (l_{Mc} \vee l_{Nc}) \} \vee \{ w_c \wedge (l_{Ma} \vee l_{Na}) \} \vee \{ w_a \wedge w_c \} = (l_{Ma} \vee l_{Na} \vee w_a) \wedge (l_{Mc} \vee l_{Nc} \vee w_c)$. Similarly as in the proof of Theorem 1, we can show that

$$(1) \quad (l_{Ma} \vee l_{Na}) \wedge (l_{Mc} \vee l_{Nc}) \leq l_{Ma \wedge c} \vee l_{Na \wedge c}.$$

As $a \wedge c = (x_1 \wedge c) \vee (x_2 \wedge c) \vee \dots \vee (x_n \wedge c)$ for each sequence

x_1, x_2, \dots, x_n with the property $x_1 \vee \dots \vee x_n = a$, $W_{a \wedge c} \geq$
 $\geq (l_{Mx_1 \wedge c} \vee l_{Nx_1 \wedge c}) \wedge \dots \wedge (l_{Mx_n \wedge c} \vee l_{Nx_n \wedge c}) \geq \{ (l_{Mx_1} \vee l_{Nx_1}) \wedge$
 $\wedge (l_{Mc} \vee l_{Nc}) \} \wedge \{ (l_{Mx_2} \vee l_{Nx_2}) \wedge (l_{Mc} \vee l_{Nc}) \} \wedge \dots \wedge \{ (l_{Mx_n} \vee$
 $\vee l_{Nx_n}) \wedge (l_{Mc} \vee l_{Nc}) \} = \{ (l_{Mx_1} \vee l_{Nx_1}) \wedge \dots \wedge (l_{Mx_n} \vee l_{Nx_n}) \} \wedge$
 $\wedge (l_{Mc} \vee l_{Nc})$. By forming the join of all terms $\{ (l_{Mx_1} \vee l_{Nx_1}) \wedge$
 $\wedge \dots \wedge (l_{Mx_n} \vee l_{Nx_n}) \} \wedge (l_{Mc} \vee l_{Nc})$, where $x_1 \vee \dots \vee x_n = a$,
 we obtain the term $W_a \wedge (l_{Mc} \vee l_{Nc})$, and as each member of
 the join was less or equal to $W_{a \wedge c}$, then

$$(2) \quad W_{a \wedge c} \geq W_a \wedge (l_{Mc} \vee l_{Nc}).$$

Similarly we see that

$$(3) \quad W_{a \wedge c} \geq W_c \wedge (l_{Ma} \vee l_{Na}).$$

Consider finally the term $W_a \wedge W_c$. Let $a = x_1 \vee \dots \vee x_n$ and
 $c = y_1 \vee \dots \vee y_m$, then $a \wedge c = (x_1 \wedge y_1) \vee (x_2 \wedge y_1) \vee \dots$
 $\vee (x_n \wedge y_1) \vee (x_1 \wedge y_2) \vee (x_2 \wedge y_2) \vee \dots \vee (x_n \wedge y_2) \vee (x_1 \wedge y_3) \vee$
 $\vee \dots \vee (x_n \wedge y_m)$. According to the definition of $W_{a \wedge c} \geq$
 $\geq (l_{Mx_1 \wedge y_1} \vee l_{Nx_1 \wedge y_1}) \wedge (l_{Mx_2 \wedge y_1} \vee l_{Nx_2 \wedge y_1}) \wedge \dots \wedge (l_{Mx_n \wedge y_m} \vee$
 $\vee l_{Nx_n \wedge y_m})$. On the other hand,

$$(l_{Mx_1 \wedge y_1} \vee l_{Nx_1 \wedge y_1}) \geq (l_{Mx_1} \vee l_{Nx_1}) \wedge (l_{My_1} \vee l_{Ny_1}),$$

$$(l_{Mx_2 \wedge y_1} \vee l_{Nx_2 \wedge y_1}) \geq (l_{Mx_2} \vee l_{Nx_2}) \wedge (l_{My_1} \vee l_{Ny_1}),$$

\vdots
 \vdots

$$(l_{Mx_n \wedge y_1} \vee l_{Nx_n \wedge y_1}) \geq (l_{Mx_n} \vee l_{Nx_n}) \wedge (l_{My_1} \vee l_{Ny_1}),$$

\vdots
 \vdots

$$(l_{Mx_n \wedge y_m} \vee l_{Nx_n \wedge y_m}) \geq (l_{Mx_n} \vee l_{Nx_n}) \wedge (l_{My_m} \vee l_{Ny_m}),$$

and by forming the meets of both sides and by ordering the

terms in the right side, we see that $W_{a \wedge c} \geq (l_{Mx_1 \wedge y_1} \vee \vee l_{Nx_1 \wedge y_1}) \wedge \dots \wedge (l_{Mx_n \wedge y_n} \vee l_{Nx_n \wedge y_n}) \geq (l_{Mx_1} \vee l_{Nx_1}) \wedge \wedge (l_{Mx_2} \vee l_{Nx_2}) \wedge \dots \wedge (l_{Mx_n} \vee l_{Nx_n}) \wedge (l_{My_1} \vee l_{Ny_1}) \wedge (l_{My_2} \vee \vee l_{Ny_2}) \wedge \dots \wedge (l_{My_n} \vee l_{Ny_n})$.

By forming the join over all pairs (x_1, \dots, x_n) and (y_1, \dots, \dots, y_n) , where $x_1 \vee \dots \vee x_n = a$ and $y_1 \vee \dots \vee y_n = c$, we see that

$$(4) \quad W_{a \wedge c} \geq W_a \wedge W_c.$$

By combining now the results (1), (2), (3) and (4) obtained above, we see that $(l_{Ma} \vee l_{Na} \vee W_a) \wedge (l_{Mc} \vee l_{Nc} \vee W_c) \leq (l_{Ma \wedge c} \vee \vee l_{Na \wedge c} \vee W_{a \wedge c})$. Obviously $a \wedge c \leq (l_{Ma} \vee l_{Na} \vee W_a) \wedge (l_{Mc} \vee l_{Nc} \vee \vee W_c)$, and the assertion follows. So H satisfies also (DK2), and hence H is a normality relation on L.

Let K be a normality relation on L such that $K \geq N, M$. According to (DK3), $xK(l_{Nx} \vee l_{Mx})$ for each $x \in L$, and according to (DK4) and (DK3), $xK(x \vee W_x)$ for each $x \in L$. By applying (DK3) once again, we see that $xK(l_{Nx} \vee l_{Mx} \vee W_x)$ for each $x \in L$, and hence $K \geq H$. Thus $H = N \vee M$, and the lemma follows.



Now we can prove a theorem on the distributivity of the lattice $GN(L)$.

Theorem 3. The lattice $GN(L)$ of all generalized normality relations on a finite lattice is distributive if and only if L is distributive and $GH = GN \vee GM$ is determined by the family $\mathcal{A}(GH) = \{ [x, l_{GNx} \vee l_{GMx}] \mid x \in L \}$.

Proof. Let L be a finite distributive lattice satisfying the condition of the theorem, and GK, GN and GM three generalized normality relations on L. It is sufficient to show that $GK \wedge (GN \vee GM) \leq (GK \wedge GN) \vee (GK \wedge GM)$, from which the

distributivity of $GN(L)$ follows. Let $a \{GK \wedge (GN \vee GM)\} b \iff \iff aGKb$ and $a(GN \vee GM)b$. Furthermore, $a(GN \vee GM)b \implies b \in \in [a, l_{GNa} \vee l_{GMa}]$, and so $b = b \wedge (l_{GNa} \vee l_{GMa}) = (b \wedge l_{GNa}) \vee \vee (b \wedge l_{GMa})$. Trivially, $a(GK \wedge GN)(b \vee l_{GNa})$ and $a(GK \wedge GM)(b \vee \vee l_{GMa})$, which imply according to (DK3) that $a \{ (GK \wedge GM) \vee \vee (GK \wedge GN) \} b$. Thus $GK \wedge (GN \vee GM) = (GK \wedge GN) \vee (GK \wedge GM)$.

In the converse part we shall first show that L is necessarily distributive. If L is non-distributive, it contains as a sublattice at least one of the lattices L' and L'' of Figure 2. Consider first the case of sublattice L' .



As L is finite, we can construct five normality relations such that the only nontrivial interval in the family \mathcal{A} generat-

ing the relations is $[0, q]$, $[0, a]$, $[0, b]$, $[0, c]$ or $[0, e]$; we denote the corresponding relations by $G[0, q]$, $G[0, a]$, $G[0, b]$, $G[0, c]$ and $G[0, e]$. Clearly these relations form a non-distributive sublattice of the lattice $GN(L)$ as $U_0 \leq q$. Similarly we see that the lattice $GN(L)$ of a lattice L containing L'' as sublattice, contains a non-distributive sublattice. Hence L is distributive.

If the join $GH = GN \vee GM$ cannot be generated by the family $\mathcal{A}(GH) = \{ [x, l_{GMx} \vee l_{GNx}] \mid x \in L \}$, we obtain the cases of the proof of Theorem 2 given in Figure 1. In the cases of Figure 1(a) and 1(b), we define GK as follows: $sGKu \iff \iff s = u$ or $\exists t \in L$ such that $t \wedge (x \wedge y) = s$ and $t \wedge z \geq u$.

As L is distributive, GK is a generalized normality relation on L ; GN and GM are defined similarly as in the proof of Theorem 2. So $(x \wedge y) \{ GK \wedge (GN \vee GM) \} z$. According to the definition of GK , for each $d \geq x \wedge y$, $A_{KGd} = [d, d]$, and hence $U_{x \wedge y} = x \wedge y$ for $(GK \wedge GM) \vee (GK \wedge GN)$. On the other hand, the proof of Theorem 2 shows that there are not in L two non-comparable elements $b \geq x$ and $c \geq y$ such that $b \vee c = z$ and $b \wedge c = x \wedge y$, whence the relation $(x \wedge y) \{ (GK \wedge GM) \vee (GK \wedge GN) \} z$ does not hold. The proof is similar in the case of Figure 1(c). This completes the proof.

3. On direct decompositions. At first we prove a theorem on direct decompositions by means of generalized normality relations.

Theorem 4. Let L be a finite lattice such that $L = L'_1 \times L'_2 \times \dots \times L'_n$, where L'_i is a chain. L has a direct decomposition if and only if there are two nontrivial generalized normality relations $GM, GK \in GN(L)$ such that $GM \wedge GK = 0$ and $GM \vee GK = 1$ in $GN(L)$.

Proof. 1° : Let $L = L_1 \times L_2$. We define two relations as follows: $aGMb \iff a = (x_1, x_2), b = (x_1, y_2)$ and $x_2 \leq y_2$; $cGKd \iff c = (z_1, z_2), d = (w_1, z_2)$ and $z_1 \leq w_1$. It is an exercise to show that GM and GK are generalized normality relations on L ; we shall only show that GM and GK are complements in $GN(L)$. Let $t \leq u$ in L , where $u = (u_1, u_2)$ and $t = (t_1, t_2)$. Then $(u_1, u_2)GM(u_1, t_2)$ and $(u_1, u_2)GK(t_1, u_2)$. Furthermore, $(t_1, u_2) \vee (u_1, t_2) = (u_1 \vee t_1, u_2 \vee t_2) = (t_1, t_2)$, and so the relations above imply $a(GK \vee GM)t$. Hence $GM \vee GK = 1$. If $h(GM \wedge GK)f$, then according to the definition of GM ,

$h_1 = f_1$ in $h = (h_1, h_2)$ and $f = (f_1, f_2)$. Similarly GK implies that $h_2 = f_2$, whence $(h_1, h_2) = (f_1, f_2) = h = f$. Thus $GK \wedge GM = 0$.

2°: Let $GM \wedge GK = 0$ and $GM \vee GK = 1$ in $GN(L)$. We shall show that $L = [0, l_{GKO}] \times [0, l_{GMO}]$. Each join-irreducible element of L belongs to one of the sets $[0, l_{GKO}]$, $[0, l_{GMO}]$. Indeed, assume that x is join-irreducible and $x \notin [0, l_{GKO}]$, $[0, l_{GMO}]$. Then $x \in [0, l_{GKO} \vee l_{GMO}]$, as $GM \vee GK = 1$. So $x \wedge (l_{GKO} \vee l_{GMO}) = (x \wedge l_{GKO}) \vee (x \wedge l_{GMO})$, from which it follows that x is join-reducible, or $l_{GKO} = 0$, or $l_{GMO} = 0$, and $x \in [0, l_{GMO}]$, or $x \in [0, l_{GKO}]$, respectively; a contradiction in each case. Furthermore, $GM \wedge GK = 0$, and so $[0, l_{GMO}] \cap [0, l_{GKO}] = \{0\}$. As L is finite and distributive, for each $z \in L$, z is the join of suitable join-irreducibles, i.e. $z = (\bigvee_i (q_{GK}^z)_i) \vee (\bigvee_j (p_{GM}^z)_j)$, where $(q_{GK}^z)_i$ is a join-irreducible of $[0, l_{GKO}]$ and $(p_{GM}^z)_j$ a join-irreducible of $[0, l_{GMO}]$. Clearly $\bigvee_i (q_{GK}^z)_i = q_{GK}^z \in [0, l_{GKO}]$ and $\bigvee_j (p_{GM}^z)_j = p_{GM}^z \in [0, l_{GMO}]$. We map z onto (q_{GK}^z, p_{GM}^z) . According to the uniqueness of the joinrepresentation by means of join-irreducibles in a distributive lattice, the mapping is a lattice morphism. If z has the figures (q_{GK}^z, p_{GM}^z) and $(q_{GK}^{z1}, p_{GM}^{z1})$, then the uniqueness of the joinrepresentation implies that $p_{GM}^z = p_{GM}^{z1}$ and $q_{GK}^z = q_{GK}^{z1}$. Similarly we see that each element of $[0, l_{GKO}] \times [0, l_{GMO}]$ has an image in L , and hence $L = [0, l_{GKO}] \times [0, l_{GMO}]$. This completes the proof.

As in the case of the preceding theorem $GN(L)$ is distributive, one can prove the following generalization by an

analogous way.

Corollary. Let L be a finite lattice, $L = L'_1 \times \dots \times L'_m$, where L'_1, \dots, L'_m are chains. L has a direct decomposition with n factors if and only if there are n nontrivial generalized normality relations GM_1, GM_2, \dots, GM_n such that $GM_k \wedge GM_j = 0$ for each pair k, j , $k \neq j$, and $GM_1 \vee GM_2 \vee \dots \vee GM_n = 1$ in $GN(L)$.

The following theorem gives the corresponding result in the case of normality relations.

Theorem 5. Let L be a finite lattice such that $L = L'_1 \times \dots \times L'_m$, where L'_1, \dots, L'_m are chains. L has a direct decomposition if and only if there are two nontrivial normality relations $K, M \in N(L)$ such that $K \wedge M = 0$ and $K \vee M = 1$ in $N(L)$.

Proof. 1^0 : Let $L = L_1 \times L_2$. We define K and M similarly as the generalized normality relations of Theorem 4: $aKb \iff a = (a_1, a_2)$, $b = (a_1, b_2)$ and $a_2 \leq b_2$; $cMd \iff c = (c_1, c_2)$, $d = (d_1, d_2)$ and $c_1 \leq d_1$. We shall show that (DK4) holds for K ; the proof is similar for M . Let aKb and fKh . Then $a \vee f = (a_1 \vee f_1, a_2 \vee f_2)$ and $h \wedge b = (a_1 \wedge f_1, b_2 \wedge h_2)$. Further, $a \vee f \vee (h \wedge b) = (a_1 \vee f_1 \vee (a_1 \wedge f_1), a_2 \vee f_2 \vee (b_2 \wedge h_2)) = (a_1 \vee f_1, a_2 \vee f_2 \vee (b_2 \wedge h_2))$. The first components of $a \vee f$ and $a \vee f \vee (h \wedge b)$ are the same and $a_2 \vee f_2 \leq a_2 \vee f_2 \vee (b_2 \wedge h_2)$, whence $(a \vee f)K(a \vee f \vee (h \wedge b))$. The other conditions hold obviously, and hence K and M are normality relations. The latter part of 1^0 is a repetition of 1^0 in the proof of Theorem 4, and hence we omit it.

2^0 : We shall show that the construction of the proof

2° of Theorem 4 holds. We must only show that each join-irreducible element x of L belongs to $[0, l_{KO}]$ or to $[0, l_{MO}]$; in fact, we show that $l_{KO} \vee l_{MO} = 1$ in L . Let us consider the normality relation $K \vee M$. $A_{K \vee M, 0} = [0, l_{KO} \vee l_{MO} \vee w_0]$, and as the only join-expression for 0 is $0 = 0 \vee 0$, $w_0 = (l_{KO} \vee l_{MO}) \wedge (l_{KO} \vee l_{MO})$, we see that $A_{K \vee M, 0} = [0, l_{KO} \vee l_{MO}]$. Furthermore, as $K \vee M = 1$ in $N(L)$, then $A_{K \vee M, 0} = L$, and hence $l_{KO} \vee l_{MO} = 1$ in L . The rest is a repetition of the proof 2° in Theorem 4.

As we have not shown the distributivity of $N(L)$, the corollary of Theorem 4 need not hold in the case of normality relations.

R e f e r e n c e s

- [1] L. BERAN: Note on a normality relation in lattices, Acta Univ. Carolinae Math. Phys. 16(1975), 59-62.
- [2] I. CHAJDA and B. ZELINKA: Tolerance relations on lattices, Časopis pěst. mat. 99(1974), 394-399.
- [3] F. HARARY: Graph theory, Addison-Wesley, Reading Mass. (1969).
- [4] G. SZÁSZ: Théorie des treillis, Akad. Kiadó, Budapest (1971).

(Oblatum 25.5. 1976)

Dept. of Technical Sciences
 Finnish Academy
 Lauttasaarentie 1
 00200 Helsinki 20, Finland