

Josef Dalík

Stability of quadratic interpolation polynomials in vertices of triangles without obtuse angles

Archivum Mathematicum, Vol. 35 (1999), No. 4, 285--297

Persistent URL: <http://dml.cz/dmlcz/107703>

Terms of use:

© Masaryk University, 1999

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

STABILITY OF QUADRATIC INTERPOLATION POLYNOMIALS IN VERTICES OF TRIANGLES WITHOUT OBTUSE ANGLES

JOSEF DALÍK

ABSTRACT. An explicit description of the basic Lagrange polynomials in two variables related to a six-tuple a^1, \dots, a^6 of nodes is presented. Stability of the related Lagrange interpolation is proved under the following assumption: a^1, \dots, a^6 are the vertices of triangles T_1, \dots, T_4 without obtuse inner angles such that T_1 has one side common with T_j for $j = 2, 3, 4$.

1. INTRODUCTION AND NOTATIONS

We say that the nodes a^1, \dots, a^6 are *regular* if the following conditions (a), (b) are satisfied.

- (a) $T_1 = \overline{a^1 a^3 a^5}$ is a central triangle and $T_2 = \overline{a^1 a^2 a^3}$, $T_3 = \overline{a^3 a^4 a^5}$, $T_4 = \overline{a^5 a^6 a^1}$ are its neighbours.
(b) $\alpha \leq \pi/2$ for all inner angles α of T_1, \dots, T_4 .

We denote by h the maximal length of sides of T_1, \dots, T_4 , by α_0 the minimal inner angle of T_1, \dots, T_4 and by T the (closed) convex hull of $T_1 \cup \dots \cup T_4$ in R^2 . We adopt the convention that a^{i+j} denotes the node a^{i+j-6} for any $i, j \in \{1, \dots, 6\}$ such that $i + j > 6$.

For $i = 1, \dots, 6$, we denote by l_i a certain quadratic polynomial such that $l_i(a^j) = 0$ whenever $j \neq i$ and describe a positive lower bound of $|l_i(a^i)|$ explicitly.

The fact that $l_i(a^i) \neq 0$ for $i = 1, \dots, 6$ guaranties existence and unicity of the solution of Lagrange interpolation problem in the nodes a^1, \dots, a^6 by second-degree polynomials in two variables. In Sauer, Xu [2], such nodes are called *poised* and are studied in a general setting. The value of lower estimate of $|l_i(a^i)|$ from our Main Theorem assures stability of the interpolation polynomial in the following sense: In Dalík [1], it is derived from this estimate that interpolation errors of values of functions as well as of their first and second partial derivatives are of optimal order.

1991 *Mathematics Subject Classification*: 65D05.

Key words and phrases: quadratic Lagrange interpolation in 2D, stability.

This work was supported by the grant No. 201/95/0095 of the Grant Agency of Czech Republic.

Received July 10, 1997.

We denote by x_1, x_2 the coordinates of any point $x \in R^2$. Let $a, b, c, d \in R^2$. We define the halflines

$$\overrightarrow{ab} = \{a + t \cdot \vec{ab}; 0 \leq t\}, \quad \overleftarrow{ab} = \{b + t \cdot \vec{ba}; 0 \leq t\}$$

and put $\overleftrightarrow{ab} = \overrightarrow{ab} \cap \overleftarrow{ab}, \overline{ab} = \overrightarrow{ab} \cup \overleftarrow{ab}$. We denote by $|ab|$ the length of the segment \overleftrightarrow{ab} and by $d(a, \overline{bc})$ the distance of a from the line \overline{bc} . Further, we put

$$D(abc) = \frac{1}{2} \begin{vmatrix} a_1 - c_1 & a_2 - c_2 \\ b_1 - c_1 & b_2 - c_2 \end{vmatrix}, \quad P(abc) = |D(abc)|.$$

Then $P(abc)$ is the area of the triangle \overline{abc} and $D(abc) > 0, D(abc) < 0$ whenever the orientation of a, b, c is positive, negative respectively. We denote by $P(abcd)$ the area of the tetragon \overline{abcd} and abbreviate $P(T_i)$ by P_i for $i = 1, \dots, 4$.

2. MAIN THEOREM

Definition. We relate the polynomial

$$l_i(x) = D(xa^{4+i}a^{5+i})D(xa^{1+i}a^{2+i})D(a^{3+i}a^{4+i}a^{2+i})D(a^{3+i}a^{5+i}a^{1+i}) \\ - D(xa^{2+i}a^{4+i})D(xa^{1+i}a^{5+i})D(a^{3+i}a^{4+i}a^{5+i})D(a^{3+i}a^{1+i}a^{2+i})$$

to $i = 1, \dots, 6$.

Lemma 1. *If $i, j \in \{1, \dots, 6\}$ then we have*

$$l_i(a^j) = \begin{cases} 0 & j \neq i, \\ (-1)^{i+1} [D(a^1a^5a^6)D(a^1a^2a^3)D(a^4a^5a^3)D(a^4a^6a^2) \\ - D(a^1a^3a^5)D(a^1a^2a^6)D(a^4a^5a^6)D(a^4a^2a^3)] & j = i. \end{cases}$$

Proof. See [1]. □

Notations. We denote by $\alpha_i, \beta_i, \gamma_i, \delta_i$ the inner angle at a^i in the triangle T_1, T_2, T_3, T_4 respectively and put $\varphi_1 = \alpha_1 + \beta_1 + \delta_1, \varphi_2 = \beta_2, \varphi_3 = \alpha_3 + \beta_3 + \gamma_3, \varphi_4 = \gamma_4, \varphi_5 = \alpha_5 + \gamma_5 + \delta_5, \varphi_6 = \delta_6$. See Fig.1. By definition,

$$(1) \quad l_1(a^1) = D(a^1a^5a^6)D(a^1a^2a^3)D(a^4a^5a^3)D(a^4a^6a^2) \\ - D(a^1a^3a^5)D(a^1a^2a^6)D(a^4a^5a^6)D(a^4a^2a^3).$$

We denote this expression more exactly by $l(a^1, a^2, a^3, a^4, a^5, a^6)$ and more briefly by l .

We find a positive lower estimate of $l_1(a^1)$ in each of the four cases characterized by the number of angles from the set $\{\varphi_1, \varphi_3, \varphi_5\}$ which are less than π separately.

Lemma 2. Let a^1, \dots, a^6 be regular nodes such that $\varphi_1 < \pi, \varphi_3 < \pi, \varphi_5 < \pi$. Then

$$l_1(a^1) \geq \frac{7}{8} \cdot P_1 P_2 P_3 P_4.$$




Fig. 1

Proof. In this case we have

$$D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) > 0 \quad \text{and}$$

$$D(a^1 a^3 a^5) D(a^1 a^2 a^6) D(a^4 a^5 a^6) D(a^4 a^2 a^3) > 0,$$

so that l is a difference of two positive values. We observe

$$(2) \quad \overline{a^4 a^5} \cap \overline{a^2 a^1} \neq \emptyset, \quad \overline{a^6 a^1} \cap \overline{a^4 a^3} \neq \emptyset, \quad \overline{a^2 a^3} \cap \overline{a^6 a^5} \neq \emptyset:$$

It follows by $\varphi_3 < \pi, \varphi_2 \leq \frac{\pi}{2}$ and $\varphi_4 \leq \frac{\pi}{2}$ that $\overline{a^4 a^5} \cap \overline{a^2 a^1} \neq \emptyset$. The rest of (2) can be justified analogically.

$$(3) \quad (a^2)' \in \overline{\overline{a^2 a^3}} \Rightarrow l(a^1, (a^2)', a^3, a^4, a^5, a^6) \leq l(a^1, a^2, a^3, a^4, a^5, a^6) \frac{|(a^2)' a^3|}{|a^2 a^3|}:$$

With respect to (2) we put $\overline{a^3 a^2} \cap \overline{a^6 a^1} = \{\tilde{a}^2\}$ and $\overline{a^2 a^3} \cap \overline{a^6 a^4} = \{\tilde{a}^3\}$; see Fig.1. Then

$$D(a^1(a^2)'a^3) = D(a^1a^2a^3)\frac{|(a^2)'a^3|}{|a^2a^3|}, \quad D(a^4a^6(a^2)') = D(a^4a^6a^2)\frac{|(a^2)'\tilde{a}^3|}{|a^2\tilde{a}^3|},$$

$$D(a^4(a^2)'a^3) = D(a^4a^2a^3)\frac{|(a^2)'a^3|}{|a^2a^3|}, \quad D(a^1(a^2)'a^6) = D(a^1a^2a^6)\frac{|(a^2)'\tilde{a}^2|}{|a^2\tilde{a}^2|}.$$

These identities and the fact that $\frac{|(a^2)'\tilde{a}^3|}{|a^2\tilde{a}^3|} \leq 1 \leq \frac{|(a^2)'\tilde{a}^2|}{|a^2\tilde{a}^2|}$ lead to

$$l(a^1, (a^2)', a^3, a^4, a^5, a^6) =$$

$$D(a^1a^5a^6)D(a^1a^2a^3)D(a^4a^5a^3)D(a^4a^6a^2)\frac{|(a^2)'a^3||a^2'\tilde{a}^3|}{|a^2a^3||a^2\tilde{a}^3|} -$$

$$D(a^1a^3a^5)D(a^1a^2a^6)D(a^4a^5a^6)D(a^4a^2a^3)\frac{|(a^2)'\tilde{a}^2||a^2'a^3|}{|a^2\tilde{a}^2||a^2a^3|} \leq$$

$$l(a^1, a^2, a^3, a^4, a^5, a^6)\frac{|(a^2)'a^3||a^2'\tilde{a}^3|}{|a^2a^3||a^2\tilde{a}^3|} \leq l(a^1, a^2, a^3, a^4, a^5, a^6)\frac{|(a^2)'a^3|}{|a^2a^3|}.$$

Hence, if we choose a point $(a^2)'$, $(a^4)'$ and $(a^6)'$ in the segment $\overline{a^2a^3}$, $\overline{a^4a^5}$ and $\overline{a^6a^1}$ in such a way that the triangle $T'_2 = \overline{a^1(a^2)'a^3}$, $T'_3 = \overline{a^3(a^4)'a^5}$ and $T'_4 = \overline{a^5(a^6)'a^1}$ has a right angle at $(a^2)'$, $(a^4)'$ and $(a^6)'$ respectively then we obtain

$$(4) \quad l(a^1, (a^2)', a^3, (a^4)', a^5, (a^6)') \leq l(a^1, a^2, a^3, a^4, a^5, a^6)\frac{P'_2P'_3P'_4}{P_2P_3P_4};$$

here P'_i denotes the area of T'_i for $i = 2, 3, 4$.

Obviously, we have

$$(5) \quad \varphi_2 = \varphi_4 = \varphi_6 = \frac{\pi}{2} \Rightarrow \varphi_1 + \varphi_3 + \varphi_5 = \frac{5}{2}\pi \quad \text{and}$$

$$(6) \quad \varphi_2 = \varphi_4 = \varphi_6 = \frac{\pi}{2} \Rightarrow$$

$$l = P_2P_3P_4[P(a^4a^6a^2) - P(a^1a^3a^5) \sin \varphi_1 \sin \varphi_3 \sin \varphi_5],$$

since in this case $l = \frac{1}{8}|a^1a^2||a^2a^3||a^3a^4||a^4a^5||a^5a^6||a^6a^1|[P(a^4a^6a^2) - P(a^1a^3a^5) \sin \varphi_1 \sin \varphi_3 \sin \varphi_5]$.

$$(7) \quad \varphi_2 = \varphi_4 = \varphi_6 = \frac{\pi}{2} \Rightarrow \sin \varphi_1 \sin \varphi_3 \sin \varphi_5 \leq \frac{1}{8} :$$

According to (5), it is sufficient to find a maximum of the expression

$$V(\varphi_1, \varphi_3) = \sin \varphi_1 \sin \varphi_3 \sin\left(\frac{5}{2}\pi - \varphi_1 - \varphi_3\right)$$

on the domain

$$\Omega = \{[\varphi_1, \varphi_3]; \frac{\pi}{2} < \varphi_1 < \pi, \frac{3\pi}{2} - \varphi_1 < \varphi_3 < \pi\}.$$

One can easily see that V attains its maximum $\frac{1}{8}$ at the point $[\frac{5\pi}{6}, \frac{5\pi}{6}]$ from Ω .

$$(8) \quad \varphi_2 = \varphi_4 = \varphi_6 = \frac{\pi}{2} \Rightarrow P(a^1 a^3 a^5) \leq P(a^4 a^6 a^2) :$$

In the proof of (8), we denote by $\widehat{a^3 a^5}$ that half of the circle with diameter $\overline{a^3 a^5}$ whose endpoints are a^3, a^5 which satisfies $a^4 \in \widehat{a^3 a^5}$. We assume that $a^3 \notin \widehat{a^3 a^5}$ and $a^5 \notin \widehat{a^3 a^5}$. In the same sense we will use the arc $\widehat{a^5 a^1}$.




Fig. 2

We first prove (8) under the following condition:

(*) There is a sharp angle between $\delta_1 + \alpha_1, \alpha_1 + \beta_1, \beta_3 + \alpha_3, \alpha_3 + \gamma_3, \gamma_5 + \alpha_5, \alpha_5 + \delta_5$.

Let us assume that $\alpha_1 + \beta_1 < \frac{\pi}{2}$ like in Fig.2. Then, necessarily, $\beta_3 + \alpha_3 > \frac{\pi}{2}$ and we put $\overline{a^2 a^3} \cap \widehat{a^3 a^5} = \{a^7\}$. It follows by $\alpha_1 + \beta_1 < \frac{\pi}{2}$ and $\overline{a^1 a^2} \parallel \overline{a^5 a^7}$ that $\angle a^7 a^5 a^1 > \frac{\pi}{2}$. Then we denote by a^8 the point of intersection of $\overline{a^7 a^5}$ and $\widehat{a^5 a^1}$ and we see that $\overline{a^1 a^2 a^7 a^8}$ is a rectangle such that

$$P(a^1 a^3 a^5) \leq \frac{1}{2} P(a^1 a^2 a^7 a^8).$$

At the same time,

$$\frac{1}{2}P(a^1a^2a^7a^8) = P(a^1a^2a^5)$$

and, as $a^4\overline{a^5} \cap a^2\overline{a^1} \neq \emptyset$ by (2), we have

$$P(a^1a^2a^5) \leq P(a^1a^2a^4).$$

Similarly, $a^6\overline{a^1} \cap a^4\overline{a^3} \neq \emptyset$ implies

$$P(a^1a^2a^4) \leq P(a^4a^6a^2)$$

and the last four relations give us (8).

Let us now assume that (*) is not true. We put $a^2\overline{a^3} \cap a^3\widehat{a^5} = \{a^7\}$, $a^6\overline{a^5} \cap a^3\widehat{a^5} = \{(a^7)'\}$ and prove (8) in the first of the two symmetric cases

$$d(a^7, \overline{a^6a^2}) \leq d((a^7)', \overline{a^6a^2}), \quad d(a^7, \overline{a^6a^2}) > d((a^7)', \overline{a^6a^2})$$

only. See Fig.3. If we denote $a^2\overline{a^1} \cap a^5\widehat{a^1} = \{a^8\}$ then $\overline{a^2a^7a^5a^8}$ is a rectangle and

$$P(a^1a^3a^5) \leq \frac{1}{2}P(a^2a^7a^5a^8) = P(a^2a^7a^5).$$




Fig. 3

As $\overline{a^2 a^3} \cap \overline{a^6 a^5} \neq \emptyset$ by (2), we have

$$P(a^2 a^7 a^5) \leq P(a^2 a^7 a^6)$$

and the condition $d(a^7, \overline{a^6 a^2}) \leq d((a^7)', \overline{a^6 a^2})$ leads to $d(a^7, \overline{a^6 a^2}) \leq d(a^4, \overline{a^6 a^2})$. Hence

$$P(a^2 a^7 a^6) \leq P(a^2 a^4 a^6)$$

and the last three relations imply (8).

Now, the statement is a consequence of (6), (7), (8) and (4). □

Lemma 3. *Let a^1, \dots, a^6 be regular nodes such that $\varphi_1 < \pi$, $\varphi_3 < \pi$ and $\varphi_5 \geq \pi$. Then*

$$l_1(a^1) > P_2(P_3)^2 P_4 \quad \text{or} \quad l_1(a^1) > P_2 P_3 (P_4)^2.$$

Proof. We have

$$D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) > 0 \quad \text{and}$$

$$D(a^1 a^3 a^5) D(a^1 a^2 a^6) D(a^4 a^5 a^6) D(a^4 a^2 a^3) \leq 0,$$

so that

$$(9) \quad l \geq D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2).$$

We can see that either $d(a^2, \overline{a^4 a^6}) > d(a^1, \overline{a^4 a^6})$ or $d(a^2, \overline{a^4 a^6}) > d(a^3, \overline{a^4 a^6})$ and, at the same time,

$$(10) \quad d(a^6, \overline{a^3 a^4}) > d(a^5, \overline{a^3 a^4}).$$




Fig. 4

Let us consider the case $d(a^2, \overline{a^4 a^6}) > d(a^3, \overline{a^4 a^6})$ from Fig.4. This relation and (10) imply $P(a^4 a^6 a^2) > P(a^4 a^5 a^3)$. This and (9) give $l > P_2(P_3)^2 P_4$. The case $d(a^2, \overline{a^4 a^6}) > d(a^1, \overline{a^4 a^6})$ leads to $l > P_2 P_3 (P_4)^2$. □

Lemma 4. Let a^1, \dots, a^6 be regular nodes such that $\varphi_1 < \pi$, $\varphi_3 \geq \pi$ and $\varphi_5 \geq \pi$. Then

$$l_1(a^1) > \frac{2}{3} \sin^2 \alpha_0 P_2(P_3)^2 P_4.$$

Proof. In this case

$$D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) > 0 \quad \text{and}$$

$$D(a^1 a^3 a^5) D(a^1 a^2 a^6) D(a^4 a^5 a^6) D(a^4 a^2 a^3) \geq 0,$$

so that l is a difference of two non-negative values. If we use the points $(a^3)'$, $(a^3)''$, $(a^5)'$, $(a^5)''$ from Fig.5 then we have




Fig. 5

$$(11) \quad \begin{aligned} &D(a^1 (a^5)' a^6) D(a^1 a^2 (a^3)') D(a^4 (a^5)' (a^3)') D(a^4 a^6 a^2) = \\ &D(a^1 (a^3)' (a^5)') D(a^1 a^2 a^6) D(a^4 (a^5)' a^6) D(a^4 a^2 (a^3)'), \end{aligned}$$

since the value of both sides of (11) is equal to

$$\frac{1}{16} d(a^1, \overline{a^6 a^2})^2 d(a^4, \overline{a^6 a^2})^2 |(a^5)' a^6| |a^2 (a^3)'| |(a^5)' (a^3)'| |a^2 a^6|.$$

$$(12) \quad l > P_2(P_3)^2 P_4 \left(1 - \frac{P(a^1 a^3 a^5) |(a^3)' (a^3)''| |(a^5)' (a^5)''|}{P(a^1 (a^3)' (a^5)') |a^3 (a^3)''| |a^5 (a^5)''|} \right) :$$

If we insert

$$D(a^1 a^5 a^6) \frac{|(a^5)'(a^5)''|}{|a^5(a^5)''|}, \quad D(a^1 a^2 a^3) \frac{|(a^3)'(a^3)''|}{|a^3(a^3)''|}, \quad D(a^4 a^5 a^3) \frac{|(a^3)'a^4|(a^5)'a^4|}{|a^3a^4||a^5a^4|},$$

$$D(a^4 a^5 a^6) \frac{|a^4(a^5)'|}{|a^4a^5|}, \quad D(a^4 a^2 a^3) \frac{|a^4(a^3)'|}{|a^4a^3|}$$

instead of $D(a^1(a^5)'a^6)$, $D(a^1a^2(a^3)')$, $D(a^4(a^5)'(a^3)')$, $D(a^4(a^5)'a^6)$, $D(a^4a^2(a^3)')$ respectively into (11) then we get

$$D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) \frac{|(a^3)'(a^3)''||a^5)'(a^5)''|}{|a^3(a^3)''||a^5(a^5)''|} =$$

$$D(a^1(a^3)'(a^5)') D(a^1 a^2 a^6) D(a^4 a^5 a^6) D(a^4 a^2 a^3).$$

Now, we multiply this equality by $P(a^1 a^3 a^5)/P(a^1(a^3)'(a^5)')$ and write the resulting left-hand side instead of $D(a^1 a^3 a^5)D(a^1 a^2 a^6)D(a^4 a^5 a^6)D(a^4 a^2 a^3)$ into l . We arrive at $l =$

$$D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) \left(1 - \frac{P(a^1 a^3 a^5)|(a^3)'(a^3)''||a^5)'(a^5)''|}{P(a^1(a^3)'(a^5)')|a^3(a^3)''||a^5(a^5)''|}\right).$$

This and the fact that $P(a^4 a^6 a^2) > P(a^4 a^5 a^3)$ give (12).

Let us denote by ω the angle between the lines $\overline{a^3 a^5}$, $\overline{(a^3)'(a^5)'}$ and put

$$V = d(a^1, \overline{a^3 a^5}), \quad v = \frac{d(a^1, \overline{(a^3)'(a^5)'})}{\cos \omega}.$$

See Fig.5. We can easily see that $|a^3 a^5| < |(a^3)'(a^5)'| \cos \omega$. By means of this fact, we verify

$$(13) \quad \frac{P(a^1 a^3 a^5)}{P(a^1(a^3)'(a^5)')} < \frac{V}{v} :$$

$$P(a^1 a^3 a^5) = \frac{1}{2} V |a^3 a^5| < \frac{1}{2} V |(a^3)'(a^5)'| \cos \omega =$$

$$\frac{1}{2} v \cos \omega |(a^3)'(a^5)'| \frac{V}{v} = P(a^1(a^3)'(a^5)') \frac{V}{v}.$$

It follows by (13) that

$$(14) \quad 1 - \frac{P(a^1 a^3 a^5)|(a^3)'(a^3)''||a^5)'(a^5)''|}{P(a^1(a^3)'(a^5)')|a^3(a^3)''||a^5(a^5)''|} > 1 - \frac{V|(a^3)'(a^3)''||a^5)'(a^5)''|}{v|a^3(a^3)''||a^5(a^5)''|}.$$

Because $\frac{V}{v} < \frac{|a^3(a^3)''|}{|(a^3)'(a^3)''|}$ and $\frac{V}{v} < \frac{|a^5(a^5)''|}{|(a^5)'(a^5)''|}$ hold obviously, we obtain

$$(15) \quad 1 - \frac{V|(a^3)'(a^3)''||a^5)'(a^5)''|}{v|a^3(a^3)''||a^5(a^5)''|} > \max \left\{ \frac{|a^3(a^3)'|}{|a^3(a^3)''|}, \frac{|a^5(a^5)'|}{|a^5(a^5)''|} \right\}.$$

We put

$$s \equiv s(a^1, a^2, a^3, a^4, a^5, a^6) \equiv \max \left\{ \frac{|a^3(a^3)'|}{|a^3(a^3)''|}, \frac{|a^5(a^5)'|}{|a^5(a^5)''|} \right\}.$$

Since the following implications

$$(16) \quad \begin{aligned} & b \in \overline{a^1 a^6}, |a^1 b| > |a^1 a^6| \Rightarrow \\ & s(a^1, a^2, a^3, a^4, a^5, b) < s(a^1, a^2, a^3, a^4, a^5, a^6), \end{aligned}$$

$$(17) \quad \overline{a^3 a^4 a^5} \subset \overline{a^3 b a^5} \Rightarrow s(a^1, a^2, a^3, b, a^5, a^6) < s(a^1, a^2, a^3, a^4, a^5, a^6)$$

are true obviously, we will find the lower bound of s under the assumptions

$$(18) \quad \beta_3 = \frac{\pi}{2} = \delta_5, \quad \gamma_4 = \alpha_0.$$




Fig. 6

Then we have

$$(19) \quad \overline{a^5 a^4} \cap \overline{a^1 a^2} = \emptyset,$$

weil $\gamma_4 = \alpha_0 \leq \beta_1 \leq \triangleleft a^2(a^3)''a^3$. By means of the symbols from Fig.6, we verify the assertion

$$(20) \quad \tilde{a}^6 \in \overset{=}{a^5}a^6 \Rightarrow s(a^1, a^2, a^3, a^4, a^5, \tilde{a}^6) \leq s(a^1, a^2, a^3, a^4, a^5, a^6) :$$

We prove that $\frac{|a^3(\tilde{a}^3)'|}{|a^3(\tilde{a}^3)''|} \leq \frac{|a^3(a^3)'|}{|a^3(a^3)''|}$ and $\frac{|a^5(\tilde{a}^5)'|}{|a^5(\tilde{a}^5)''|} \leq \frac{|a^5(a^5)'|}{|a^5(a^5)''|}$. Clearly, it is sufficient to verify the second inequality. Under the assumption $|(a^5)''(\tilde{a}^5)''| \leq |(a^5)''(a^5)'|$, we can see that

$$\begin{aligned} \frac{|a^5(\tilde{a}^5)'|}{|a^5(\tilde{a}^5)''|} &\leq \frac{|a^5(a^5)'|}{|a^5(a^5)''|} \Leftrightarrow \frac{|(a^5)'(a^5)''|}{|a^5(a^5)''|} \leq \frac{|(\tilde{a}^5)'(\tilde{a}^5)''|}{|a^5(\tilde{a}^5)''|} \Leftrightarrow \\ &\frac{|(a^5)''(\tilde{a}^5)''| + |(\tilde{a}^5)''(a^5)'|}{|(a^5)''(\tilde{a}^5)''| + |(\tilde{a}^5)''(a^5)'| + |(a^5)'a^5|} \leq \frac{|(\tilde{a}^5)'(a^5)'| + |(a^5)'(\tilde{a}^5)''|}{|a^5(a^5)'| + |(a^5)'(\tilde{a}^5)''|} \Leftrightarrow \\ &\frac{|a^5(a^5)'|}{|(\tilde{a}^5)'(a^5)'|} \leq \frac{|a^5(a^5)''|}{|(\tilde{a}^5)''(a^5)''|} \Leftrightarrow \frac{|a^5a^6||a^2\tilde{a}^6|}{|\tilde{a}^6a^6||a^2(\tilde{a}^5)'|} \leq \frac{|a^5a^6||a^1\tilde{a}^6|}{|\tilde{a}^6a^6||a^1(\tilde{a}^5)''|} \Leftrightarrow \\ &\frac{|a^2\tilde{a}^6|}{|a^2(\tilde{a}^5)'|} \leq \frac{|a^1\tilde{a}^6|}{|a^1(\tilde{a}^5)''|} \end{aligned}$$

and the last inequality is true by (19). The case $|(a^5)''(\tilde{a}^5)''| > |(a^5)''(a^5)'|$ leads to the same conclusion.

Because of (20), we can extend the assumptions (18) by

$$(21) \quad \beta_1 = \alpha_0 = \delta_1.$$

The lengths u and U from Fig.6 satisfy

$$(22) \quad s > \frac{u}{U},$$

because it holds either $\frac{u}{U} < \frac{|a^3(a^3)'|}{|a^3(a^3)''|}$ or $\frac{u}{U} < \frac{|a^5(a^5)'|}{|a^5(a^5)''|}$. Finally, under the assumptions (18), (21) we prove that

$$(23) \quad \frac{u}{U} > \frac{2}{3} \sin^2 \alpha_0 :$$

Let us put $a = |a^1a^6|$, $b = |a^1a^2|$ and choose the cartesian coordinate system in such a way that

$$a^1 = [0, 0], \quad a^6 = [0, a], \quad a^2 = [b \sin(\alpha_1 + 2\alpha_0), b \cos(\alpha_1 + 2\alpha_0)].$$




Fig. 7

See Fig.7. One can easily compute that

$$a^7 = \left[\frac{\cos \alpha_0}{\sin \alpha_1} (b \cos \alpha_0 - a \cos(\alpha_1 + \alpha_0)), a - \frac{\sin \alpha_0}{\sin \alpha_1} (b \cos \alpha_0 - a \cos(\alpha_1 + \alpha_0)) \right]$$

and, further, by means of the parametrization

$$x_1 = t \frac{\cos \alpha_0}{\sin \alpha_1} (b \cos \alpha_0 - a \cos(\alpha_1 + \alpha_0)),$$

$$x_2 = t \left[a - \frac{\sin \alpha_0}{\sin \alpha_1} (b \cos \alpha_0 - a \cos(\alpha_1 + \alpha_0)) \right],$$

$t \in < 0, 1 >$, of the segment $\overline{a^1 a^7}$, one can see that the value of the parameter t in the cross point $\overline{a^1 a^7} \cap \overline{a^6 a^2}$ is

$$t(Q) = \frac{Q \sin \alpha_1 \sin(\alpha_1 + 2\alpha_0)}{\cos \alpha_0 [2Q \cos \alpha_0 - (1 + Q^2) \cos(\alpha_1 + \alpha_0)]}.$$

Here $Q = \frac{a}{b}$ and the fact that $Q \in < \frac{a}{b_1}, \frac{a}{b_0} > = < \cos \alpha_1, \frac{1}{\cos \alpha_1} >$ is apparent from Fig.7. Because of $t(Q) = 1 - \frac{u}{U}$, an optimal lower bound of $\frac{u}{U}$ corresponds to

$$\max \{ t(Q); \cos \alpha_1 \leq Q \leq \frac{1}{\cos \alpha_1} \}.$$

We can find this maximum by a standard procedure. Then we conclude that

$$\frac{u}{U} \geq \frac{\sin \alpha_0 \sin \alpha_1 \sin(\alpha_1 + \alpha_0)}{\cos \alpha_0 [\cos \alpha_1 \sin(\alpha_1 + \alpha_0) + \sin \alpha_0]}.$$

As the expression on the right-hand side attains its minimum for $\alpha_1 = \alpha_0$, we find

$$\frac{u}{U} \geq \frac{\sin^2 \alpha_0 \sin 2\alpha_0}{\cos \alpha_0 (\cos \alpha_0 \sin 2\alpha_0 + \sin \alpha_0)} = \frac{2 \sin^2 \alpha_0}{1 + 2 \cos^2 \alpha_0} > \frac{2}{3} \sin^2 \alpha_0.$$

The statement is a consequence of (12), (14), (15), (22) and (23). □

Lemma 5. *Let a^1, \dots, a^6 be regular nodes such that $\varphi_1 \geq \pi$, $\varphi_3 \geq \pi$ and $\varphi_5 \geq \pi$. Then*

$$l_1(a^1) > P_1 P_2 P_3 P_4.$$

Proof. We have

$$D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) > 0 \quad \text{and}$$

$$D(a^1 a^3 a^5) D(a^1 a^2 a^6) D(a^4 a^5 a^6) D(a^4 a^2 a^3) \leq 0.$$

Since, at the same time, $\overline{a^4 a^6 a^2} \supset \overline{a^1 a^3 a^5}$, we conclude

$$l \geq D(a^1 a^5 a^6) D(a^1 a^2 a^3) D(a^4 a^5 a^3) D(a^4 a^6 a^2) > P_1 P_2 P_3 P_4. \quad \square$$

Main Theorem. *Let the nodes a^1, \dots, a^6 be regular. Then there exist an index $k \in \{1, 2, 3, 4\}$ and a positive constant C independent of h such that*

$$|l_i(a^i)| \geq C \cdot P_k P_2 P_3 P_4 \quad \text{for } i = 1, \dots, 6.$$

Proof. This statement is a consequence of the lemmas 1, 2, 3, 4, 5. □

REFERENCES

- [1] Dalík, J., *Quadratic interpolation polynomials in vertices of strongly regular triangulations*, in Finite Element Methods, superconvergence, post-processing and a posteriori estimates, Ed. Křížek, Neittaanmäki, Stenberg, Marcel Dekker (1996), 85–95.
- [2] Sauer, T., Xu, Y., *On multivariate Lagrange interpolation*, Math. of Comp. **64** (1995), 1147–1170.