

Jozef Doboš

Some generalizations of the notion of continuity and quasi-uniform convergence

Časopis pro pěstování matematiky, Vol. 106 (1981), No. 4, 431--434

Persistent URL: <http://dml.cz/dmlcz/108484>

Terms of use:

© Institute of Mathematics AS CR, 1981

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SOME GENERALIZATIONS OF THE NOTION OF CONTINUITY
AND QUASI-UNIFORM CONVERGENCE

JOZEF DOBOŠ, Tekovské Nemce

(Received June 30, 1980)

It is well known that the sets of quasi-continuous, somewhat continuous and cliquish functions are all closed with respect to the uniform convergence (see [3], [5], [9]). The aim of this paper is to investigate whether or not those sets are closed with respect to the quasi-uniform convergence (see [7], p. 143).

Let X, Y be two topological spaces. A function $f: X \rightarrow Y$ is said to be quasi-continuous at a point $x_0 \in X$ if for each neighbourhood $U(x_0)$ of the point x_0 (in X) and each neighbourhood $V(f(x_0))$ of the point $f(x_0)$ (in Y) there exists an open set $U \subset U(x_0)$, $U \neq \emptyset$ such that $f(U) \subset V(f(x_0))$ (see [5]).

A function $f: X \rightarrow Y$ is said to be somewhat continuous if for each set $V \subset Y$ open in Y such that $f^{-1}(V) \neq \emptyset$ there exists an open set $U \subset X$, $U \neq \emptyset$ such that $U \subset f^{-1}(V)$ (see [3]).

Let X be a topological and Y a metric space (with the metric d). A function $f: X \rightarrow Y$ is said to be cliquish at a point $x_0 \in X$ if for each neighbourhood $U(x_0)$ of the point x_0 and each $\varepsilon > 0$ there exists an open set $U \subset U(x_0)$, $U \neq \emptyset$ such that $d(f(x'), f(x'')) < \varepsilon$ holds for every two points $x', x'' \in U$ (see [9]).

A function f defined on a topological space X is said to be quasi-continuous or cliquish on X if it is quasi-continuous or cliquish, respectively, at each point $x \in X$.

The property of the quasi-continuity is equivalent to the property of the semi-continuity (see [4], [6]).

Every function $f: X \rightarrow Y$ quasi-continuous on X is also somewhat continuous on X (see [8]).

Proposition 1. *There exists a sequence of functions $f_n: R \rightarrow R$ quasi-uniformly converging to $f: R \rightarrow R$ such that f_n is quasi-continuous but f is not somewhat continuous.*

Proof. Let the sequence $\{f_n\}_{n=1}^\infty$ of functions $f_n: R \rightarrow R$ be defined by

$$f_n(x) = \chi_{\langle 0, 1/n \rangle}((-1)^n \cdot x)$$

for all $x \in R$. Obviously $f_n \rightarrow f = \chi_{\{0\}}$. Let $\varepsilon > 0$, $m \in \{0, 1, 2, \dots\}$. Denote $p = m + 2$. Then for $x \geq 0$ we have $|f_{m+p-1}(x) - f(x)| \approx 0$, and for $x < 0$ we have

$|f_{m+p}(x) - f(x)| = 0$, i.e. $\forall x \in R : \min \{|f_{m+1}(x) - f(x)|, \dots, |f_{m+p}(x) - f(x)|\} < \varepsilon$. Hence $\{f_n\}_{n=1}^{\infty}$ quasi-uniformly converges to f . We now show that f_n ($n = 1, 2, \dots$) are quasi-continuous functions. Let $n \in N$. Let U be an open neighbourhood of the point $x_0 = 0$ and V an open neighbourhood of the point $f_n(x_0)$. Then there exists $0 < \delta < 1/(2n)$ such that $(x_0 - 2\delta, x_0 + 2\delta) \subset U$. Denote $a = (-1)^n \cdot \delta$ and $U_0 = (a - \delta, a + \delta)$. Then U_0 is open, $\emptyset \neq U_0 \subset U$ and $f_n(U_0) \subset V$. Hence f_n is quasi-continuous at the point $x_0 = 0$. Let U be an open neighbourhood of the point $x_1 = (-1)^n/n$ and V an open neighbourhood of the point $f_n(x_1)$. Then there exists $0 < \delta < 1/(2n)$ such that $(x_1 - 2\delta, x_1 + 2\delta) \subset U$. Denote $b = (-1)^n \cdot (1/n - \delta)$ and $U_1 = (b - \delta, b + \delta)$. Then U_1 is open, $\emptyset \neq U_1 \subset U$ and $f_n(U_1) \subset V$. Hence f_n is quasi-continuous at the point $x_1 = (-1)^n/n$. Since f_n is continuous at each point $x \in R - \{0, (-1)^n/n\}$, conclude that f_n is quasi-continuous. Since $\text{int } f^{-1}((1/2, 2)) = \text{int } \{0\} = \emptyset$, f is not somewhat continuous.

Proposition 2. *There exists a nonempty set $M \subset R$ and a sequence of functions $f_n : M \rightarrow R$ quasi-uniformly converging to $f : M \rightarrow R$ such that f_n is cliquish but f is not cliquish.*

Proof. Let $A = \{a_1, a_2, \dots\}$, $B = \{b_1, b_2, \dots\}$ be countable subsets of R such that $A \cap B = \emptyset$, $\bar{A} = \bar{B} = R$. Denote $M = A \cup B$, $A_n = \{a_1, \dots, a_n\}$, $B_n = \{b_1, \dots, b_n\}$ for each $n \in N$. Define the sequence $\{f_n\}_{n=1}^{\infty}$ of functions $f_n : M \rightarrow R$ by

$$f_n = \begin{cases} \chi_{A_n} & \text{if } n \text{ is even,} \\ \chi_{(M-B_n)} & \text{if } n \text{ is odd.} \end{cases}$$

Obviously $f_n \rightarrow f = \chi_A$. Let $\varepsilon > 0$, $m \in \{0, 1, 2, \dots\}$. Denote $p = m + 2$. Then for $x \in A$ we have $|f_{m+p-1}(x) - f(x)| = 0$, and for $x \in B$ we have $|f_{m+p}(x) - f(x)| = 0$, i.e.

$$\forall x \in M : \min \{|f_{m+1}(x) - f(x)|, \dots, |f_{m+p}(x) - f(x)|\} < \varepsilon.$$

Hence $\{f_n\}_{n=1}^{\infty}$ quasi-uniformly converges to f . We now show that f_n ($n = 1, 2, \dots$) are cliquish functions. Let $n \in N$. Let $x_0 \in M$. Denote $\gamma = \min \{|x_0 - x| : x \in A_n \cup B_n, x \neq x_0\}$. Let U be an open set such that $x_0 \in U$. Let $\varepsilon > 0$. Denote $U_0 = (x_0, x_0 + \gamma) \cap U$. Then U_0 is open, $\emptyset \neq U_0 \subset U$ and

$$\forall x, x' \in U_0 : |f_n(x) - f_n(x')| = 0 < \varepsilon.$$

Since for each open set V we have

$$V \neq \emptyset \Rightarrow V \cap A \neq \emptyset \neq V \cap B,$$

f is not cliquish.

Definition 1. *A family \mathcal{A} of sets has the finite intersection property if the intersection of every finite subfamily of \mathcal{A} is nonempty. A centred family is a family of sets having the finite intersection property.*

Definition 2. An open almost-base for a space X is a family \mathcal{A} of open subsets of X such that every nonempty open subset of X contains some nonempty $A \in \mathcal{A}$.

Definition 3. Let $\{\mathcal{A}_n\}_{n=1}^\infty$ be a sequence of open families in a space X (an open family is a family consisting of open sets). The sequence $\{\mathcal{A}_n\}_{n=1}^\infty$ is said to be countably complete if for every centred sequence of sets $\{A_{n_k}\}_{k=1}^\infty$, where $A_{n_k} \in \mathcal{A}_{n_k}$, the set $\bigcap_{k \in \mathbb{N}} \bar{A}_{n_k}$ is nonempty.

Definition 4. A space X is said to be almost countably complete if there exists a countably complete sequence of open almost-bases for X (see [2]).

Remark. Every locally compact Hausdorff space is a regular almost countably complete space. Every complete metric space is a regular almost countably complete space (see [2]).

Theorem. Let X be a regular almost countably complete space and let (Y, d) be a metric space. Let $\{f_n\}_{n=1}^\infty$ be a sequence of cliquish functions $f_n : X \rightarrow Y$ quasi-uniformly converging to $f : X \rightarrow Y$. Then f is cliquish.

Proof. We show that f is cliquish at any point $x_0 \in X$. Let U be an open set such that $x_0 \in U$. Let $\varepsilon > 0$. We now show that there exists a sequence of nonempty open sets $U_n \subset U$ such that

$$(i) \quad \bigcap_{n \in \mathbb{N}} U_n \neq \emptyset,$$

$$(ii) \quad \forall n \in \mathbb{N} : \text{diam } f_n(U_n) < \varepsilon/6.$$

Let $\{\mathcal{B}_n\}_{n=1}^\infty$ be a countably complete sequence of open almost-bases for X . Since f_1 is cliquish at x_0 , there exists a nonempty set $U_1 \in \mathcal{B}_1$ such that $U_1 \subset U$, $\text{diam } f_1(U_1) < \varepsilon/6$. Suppose U_1, \dots, U_n have been constructed. Let $y \in U_n$. Since X is regular, there exists a closed neighbourhood W at y , such that $W \subset U_n$. Since f_{n+1} is cliquish at y , there exists a nonempty open set $U_{n+1} \in \mathcal{B}_{n+1}$ such that $U_{n+1} \subset \text{int } W$, $\text{diam } f_{n+1}(U_{n+1}) < \varepsilon/6$. Then

$$\emptyset \neq U_{n+1} \subset \bar{U}_{n+1} \subset W \subset U_n \subset U, \quad U_n \in \mathcal{B}_n.$$

Since $\{U_{n+1}\}_{n=1}^\infty$ is centred,

$$\emptyset \neq \bigcap_{n \in \mathbb{N}} \bar{U}_{n+1} \subset \bigcap_{n \in \mathbb{N}} U_n.$$

Let $y \in \bigcap_{n \in \mathbb{N}} U_n$. Since $f_n \rightarrow f$, we have

$$(1) \quad \exists m \in \mathbb{N} \quad \forall n \geq m : d(f(y), f_n(y)) < \varepsilon/6.$$

Since $\{f_n\}_{n=1}^\infty$ quasi-uniformly converges to f , we have $\exists p \in \mathbb{N} \quad \forall x \in X$:

$$\min \{d(f_{m+1}(x), f(x)), \dots, d(f_{m+p}(x), f(x))\} < \varepsilon/6.$$

Denote $U_0 = \bigcap_{n=1}^{m+p} U_n$. Let $x \in U_0$. We now show that $d(f(x), f(y)) < \varepsilon/2$. Then obviously $\forall x, x' \in U_0 : d(f(x), f(x')) < \varepsilon$.

Let $j \in \{1, \dots, p\}$ be such that $d(f_{m+j}(x), f(x)) < \varepsilon/6$. Then by (ii) we have $d(f_{m+j}(x), f_{m+j}(y)) < \varepsilon/6$, and by (1) we obtain $d(f_{m+j}(y), f(y)) < \varepsilon/6$, therefore $d(f(x), f(y)) \leq d(f_{m+j}(x), f(x)) + d(f_{m+j}(x), f_{m+j}(y)) + d(f_{m+j}(y), f(y)) < \varepsilon/2$.

Remark. For a different proof of this theorem, see [1].

Corollary. Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of cliquish functions $f_n : R \rightarrow R$ quasi-uniformly converging to $f : R \rightarrow R$. Then f is a cliquish function.

The author is very much indebted to Professor T. Šalát for many helpful remarks and suggestions offered during the preparation of this paper.

References

- [1] Doboš, J. - Šalát, T.: Some remarks on cliquish functions, Riemann integrable functions and quasi-uniform convergence. Acta F.R.N. Math. Univ. Comen. (to appear).
- [2] Frolík, Z.: Baire spaces and some generalizations of complete metric spaces. Czech. Math. J. 11 (86), 1961, 237–248.
- [3] Gentry, K. R. - Hoyle, H. B.: Somewhat continuous functions. Czech. Math. J. 21 (96), 1971, 5–12.
- [4] Levine, N.: Semi-open sets and semi-continuity in topological spaces. Amer. Math. Monthly, 70, 1963, 36–41.
- [5] Marcus, S.: Sur les fonctions quasicontinues au sens de S. Kempisty. Coll. Math. 8, 1961, 47–53.
- [6] Neubrunnová, A.: On certain generalizations of the notion of continuity. Mat. čas. 23, 1973, 4, 374–380.
- [7] Sikorski, R.: Real Functions I. (in Polish) PWN, Warszawa, 1958.
- [8] Šalát, T.: Some generalizations of the notion of continuity and Denjoy property of functions. Čas. pěst. mat. 99, 1974, 380–385.
- [9] Thielman, H.: Types of functions. Amer. Math. Monthly 60, 1953, 156–161.

Author's address: 966 54 Tekovské Nemce 261, Žiar nad Hronom.