

Ingo Bandlow

On function spaces of Corson-compact spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 35 (1994), No. 2, 347--356

Persistent URL: <http://dml.cz/dmlcz/118673>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On function spaces of Corson-compact spaces

INGO BANDLOW

Abstract. We apply elementary substructures to characterize the space $C_p(X)$ for Corson-compact spaces. As a result, we prove that a compact space X is Corson-compact, if $C_p(X)$ can be represented as a continuous image of a closed subspace of $(L_\tau)^\omega \times Z$, where Z is compact and L_τ denotes the canonical Lindelöf space of cardinality τ with one non-isolated point. This answers a question of Archangelskij [2].

Keywords: function spaces, Corson-compact spaces, elementary substructures

Classification: Primary 54C

1. Elementary substructures

We begin with a brief exposition of some definitions and facts concerning elementary substructures (see also A. Dow [5], K. Kunen [7]). A non-empty subset \mathcal{M} of a set \mathcal{H} is said to be an elementary substructure of \mathcal{H} , if for any formula $\phi(x_1, \dots, x_n)$ of the language of set theory with the only free variables x_1, \dots, x_n and for any $a_1, \dots, a_n \in \mathcal{M}$ $\phi[a_1, \dots, a_n]$ is true in \mathcal{M} if and only if it is true in \mathcal{H} .

A frequently used argument is Tarski’s Criterion:

A subset \mathcal{M} of \mathcal{H} forms an elementary substructure of \mathcal{H} if and only if for every formula $\phi(x_0, x_1, \dots, x_n)$ of the language of set theory and every $a_1, \dots, a_n \in \mathcal{M}$ such that there exists an $a \in \mathcal{H}$ such that $\phi[a, a_1, \dots, a_n]$ is true in \mathcal{H} , there is a $b \in \mathcal{M}$ such that $\phi[b, a_1, \dots, a_n]$ is true in \mathcal{H} (and therefore in \mathcal{M}).

The base of all our applications of elementary substructures is the following

Theorem 1.1 (Löwenheim-Skolem-Tarski). *For every infinite set \mathcal{H} and each subset $X \subseteq \mathcal{H}$ there exists an elementary substructure \mathcal{M} of \mathcal{H} such that $X \subseteq \mathcal{M}$ and $|\mathcal{M}| = \max\{|X|, \omega\}$.*

If θ is a cardinal, then $\mathcal{H}(\theta)$ denotes the collection of all sets hereditarily of cardinality $< \theta$. We will usually be interested in elementary substructures of $\mathcal{H}(\theta)$, where θ is a sufficiently large regular cardinal. The main reason is that for any sentence ϕ there exists a large enough regular cardinal θ such that ϕ is true (in V) if and only if it is true in $\mathcal{H}(\theta)$. In practice, one “chooses” θ without discussion how large it needs to be.

The following fact is well known and useful.

Proposition 1.2. *If θ is a regular uncountable cardinal, \mathcal{M} an elementary substructure of $\mathcal{H}(\theta)$ and $A \in \mathcal{M}$ a countable set, then $A \subseteq \mathcal{M}$.*

If \mathcal{H} is an uncountable $[\mathcal{H}]^\omega$ denotes the set of all countable subsets of \mathcal{H} . $C \subseteq [\mathcal{H}]^\omega$ is said to be unbounded if for every $X \in [\mathcal{H}]^\omega$ there is a $Y \in C$ with $X \subseteq Y$. We say C is closed if, whenever $X_n \in C$ and $X_n \subseteq X_{n+1}$ for each $n \in \omega$, then $\bigcup\{X_n : n \in \omega\} \in C$. A consequence of Theorem 1.1 is that the family of all countable elementary substructures of \mathcal{H} is closed and unbounded. Remark, that the intersection of two closed unbounded subsets of $[\mathcal{H}]^\omega$ is closed unbounded, too.

For the sake of simplicity we shall often write “Let \mathcal{M} be a suitable elementary substructure ...”. This means that all “information” we need to investigate an object, say a topological space, can be found in \mathcal{M} . For example, if X is a dyadic compact space, we suppose that there is a continuous mapping $f : D^\tau \rightarrow X$, which is an element of \mathcal{M} . Obviously there is always a closed unbounded family of “suitable” countable elementary substructures. On the other hand, “... if for any suitable countable elementary substructure \mathcal{M} the following condition is satisfied ...” means that the condition is satisfied for all countable substructures from a closed unbounded subset of $[\mathcal{H}(\theta)]^\omega$, where θ is a large enough (with respect to the object we investigate) regular cardinal.

2. The main construction

Now we are going to describe a construction for arbitrary uniform spaces. Let $\langle X, \mathcal{U} \rangle$ be a uniform space (see Engelking [6]). If \mathcal{M} is an elementary substructure with $X, \mathcal{U} \in \mathcal{M}$, the equivalence relation on X is defined by

$$x \approx y(\mathcal{M}) \text{ iff } |x - y| < V \text{ for each } V \in \mathcal{U} \cap \mathcal{M}.$$

If $x \approx y(\mathcal{M})$, we say that x and y are \mathcal{M} -equivalent. Let $X(\mathcal{M}, \mathcal{U})$ denote the set of all equivalence classes and $\phi_{\mathcal{M}}^{X, \mathcal{U}}$ the canonical mapping of X onto $X(\mathcal{M})$. (For short, we often write $X(\mathcal{M})$ and $\phi_{\mathcal{M}}^X$ and drop the \mathcal{U} .) A uniformity $\mathcal{U}(\mathcal{M})$ is given on $X(\mathcal{M})$ by all entourages $V_{\mathcal{M}}$ of the diagonal in $X(\mathcal{M})$. $\mathcal{U}(\mathcal{M})$ is defined by

$$|\phi_{\mathcal{M}}^X(x) - \phi_{\mathcal{M}}^X(y)| < V_{\mathcal{M}} \text{ iff } |x' - y'| < V \text{ for all } x', y' \in X \\ \text{such that } x' \approx x(\mathcal{M}) \text{ and } y' \approx y(\mathcal{M}),$$

where V is an arbitrary element of $\mathcal{U} \cap \mathcal{M}$.

$\phi_{\mathcal{M}}^X$ is uniformly continuous with respect to the uniformities \mathcal{U} and $\mathcal{U}(\mathcal{M})$ on X and $X(\mathcal{M})$ respectively (see Bandlow [3]). Now we shall give some easy assertions used in the sequel.

Fact 2.1. Let $f : \langle X, \mathcal{U} \rangle \rightarrow \langle Y, \mathcal{V} \rangle$ be a uniformly continuous mapping, suppose $f \in \mathcal{M}$. Then there exists a uniformly continuous mapping $f_{\mathcal{M}} : X(\mathcal{M}) \rightarrow Y(\mathcal{M})$

which makes the following diagram commutative:

$$(1) \quad \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \phi_{\mathcal{M}}^X \downarrow & & \downarrow \phi_{\mathcal{M}}^X \\ X(\mathcal{M}) & \xrightarrow{f_{\mathcal{M}}} & Y(\mathcal{M}) \end{array}$$

Fact 2.2. Let $\langle X, \mathcal{U} \rangle$ be the product of the uniform spaces $\{\langle X_t, \mathcal{U}_t \rangle : t \in T\}$; suppose $\{\langle X_t, \mathcal{U}_t \rangle : t \in T\} \in \mathcal{M}$. Then $\langle X(\mathcal{M}), \mathcal{U}(\mathcal{M}) \rangle$ is uniformly homeomorphic to the product of the uniform spaces $\{\langle X_t(\mathcal{M}), \mathcal{U}_t(\mathcal{M}) \rangle : t \in T \cap \mathcal{M}\}$ in a natural way.

Fact 2.3. If $\langle Y, \mathcal{U}_Y \rangle$ is a uniform subspace of $\langle X, \mathcal{U} \rangle$, then $Y(\mathcal{M})$ is uniformly homeomorphic to $\phi_{\mathcal{M}}^X(Y)$ in a natural way.

Fact 2.4. If X is a compact Hausdorff space, there is a unique uniformity \mathcal{U} on X which induces the original topology on X . \mathcal{U} is generated by all sets of the form

$$\mathcal{U}_{f_1, \dots, f_n}^\varepsilon = \{\langle x, y \rangle : |f_i(x) - f_i(y)| < \varepsilon, \quad i = 1, \dots, n\},$$

where f_1, \dots, f_n are arbitrary real-valued continuous functions on X , $n \in \mathbb{N}$ and $\varepsilon > 0$. Let \mathcal{M} be a suitable elementary substructure. It is easy to see that for any pair of distinct points $x, y \in X$ we have $\phi_{\mathcal{M}}^X(x) \neq \phi_{\mathcal{M}}^X(y)$ if and only if there is a function $f \in C(X) \cap \mathcal{M}$ with $f(x) \neq f(y)$. Consequently, $\phi_{\mathcal{M}}^X$ corresponds to the mapping which relates each point $x \in X$ to $(fx)_{C(X) \cap \mathcal{M}}$ from the product space $\mathbb{R}^{C(X) \cap \mathcal{M}}$. An easy consequence is that for each function $f \in C(X) \cap \mathcal{M}$ there is a continuous function $f_{\mathcal{M}} : X(\mathcal{M}) \rightarrow \mathbb{R}$ such that $f = f_{\mathcal{M}} \circ \phi_{\mathcal{M}}^X$. If $U \in \mathcal{M}$ is a functionally open subset of X , i.e. if there is a function $f \in C(X) \cap \mathcal{M}$ with $U = f^{-1}(0, 1)$, then let $U_{\mathcal{M}}$ be defined by $U_{\mathcal{M}} = f_{\mathcal{M}}^{-1}(0, 1)$. Remark that $U = (\phi_{\mathcal{M}}^X)^{-1}U_{\mathcal{M}}$. The family consisting of all open subsets of $X(\mathcal{M})$ of the form $U_{\mathcal{M}}$, where U is a functionally open subset of X and belongs to \mathcal{M} , is a base for the topology of $X(\mathcal{M})$.

We now prove that if X is a Lindelöf space, then $X(\mathcal{M})$ as a topological space does not depend on the uniformity on X .

Proposition 2.5. *Let X be a Lindelöf space and let \mathcal{U} and \mathcal{V} be uniformities on X , which induces the topology on X . If \mathcal{M} is a suitable elementary substructure, then:*

- (a) $\phi_{\mathcal{M}}^{X, \mathcal{U}}(x) = \phi_{\mathcal{M}}^{X, \mathcal{U}}(y)$ if and only if $\phi_{\mathcal{M}}^{X, \mathcal{V}}(x) = \phi_{\mathcal{M}}^{X, \mathcal{V}}(y)$ for arbitrary points $x, y \in X$ ($X(\mathcal{M}) = X(\mathcal{M}, \mathcal{U}) = X(\mathcal{M}, \mathcal{V})$).
- (b) $\mathcal{U}(\mathcal{M})$ and $\mathcal{V}(\mathcal{M})$ generate the same topology on $X(\mathcal{M})$.

PROOF: W.l.o.g. we may assume that $\mathcal{V} \subseteq \mathcal{U}$. Let x_0, y_0 be a pair of distinct points of X . Suppose that $\phi_{\mathcal{M}}^{X, \mathcal{U}}(x_0) \neq \phi_{\mathcal{M}}^{X, \mathcal{U}}(y_0)$, i.e. $|x_0 - y_0| \geq U$ for some

$U \in \mathcal{U} \cap \mathcal{M}$. We can choose a $W \in \mathcal{U} \cap \mathcal{M}$ such that $4W \subseteq U$. $\eta = \{\text{Int } B(x, W) : x \in X\}$ is an open cover of the space X and belongs to \mathcal{M} . Here $B(x, W)$ denotes the set $\{y \in X : |x - y| < V\}$. Therefore there exists a countable subcover $\eta' = \{\text{Int } B(x_n, W) : n = 1, 2, 3, \dots\}$ which also belongs to \mathcal{M} . By Proposition 1.2, we may assume that all points $x_n, n = 1, 2, 3, \dots$, are elements of \mathcal{M} . Hence we can fix $z_1, z_2 \in X \cap \mathcal{M}$ such that $x_0 \in B(z_1, W)$ and $y_0 \in B(z_2, W)$. $F = \text{cl}(B(z_1, W))$ and $G = \text{cl}(B(z_2, W))$ are elements of \mathcal{M} too, and we have $x_0 \in F, y_0 \in G$ and $F \cap G = \emptyset$. For every point $x \in F$ there exists an entourage $V_x \in \mathcal{V}$ such that $B(x, 2V_x) \cap G = \emptyset$. Similarly as above we consider now the open cover $\xi = \{\text{Int } B(x, V_x) : x \in F\}$ of F . Since F and G are elements of \mathcal{M} , ξ also belongs to \mathcal{M} . Let $\xi' = \{\text{Int } B(x_n, V_{x_n}) : n = 1, 2, 3, \dots\}$ be a countable subcover of ξ . We may assume that ξ' belongs to \mathcal{M} and, consequently, that all x_n and V_{x_n} are elements of \mathcal{M} . Hence, there exists a point $z \in F \cap \mathcal{M}$ and an entourage $V \in \mathcal{V} \cap \mathcal{M}$ such that $x_0 \in B(z, V)$ and $B(z, 2V) \cap G = \emptyset$, i.e. $|x_0 - y_0| > V$ and $\phi_{\mathcal{M}}^{X, \mathcal{V}}(x_0) \neq \phi_{\mathcal{M}}^{X, \mathcal{V}}(y_0)$. This proves the assertion (a) and we may identify $X(\mathcal{M}) = X(\mathcal{M}, \mathcal{U}) = X(\mathcal{M}, \mathcal{V})$. Let $T_{\mathcal{U}}$ and $T_{\mathcal{V}}$ denote the topologies on $X(\mathcal{M})$ generated by $\mathcal{U}(\mathcal{M})$ and $\mathcal{V}(\mathcal{M})$, respectively. We have to prove that the (identical) mapping

$$\langle X(\mathcal{M}), T_{\gamma} \rangle \longrightarrow \langle X(\mathcal{M}), T_{\mathcal{U}} \rangle$$

is continuous with respect to these topologies. Suppose $x_0 \in X$ and \mathcal{O} is a neighborhood of $\phi_{\mathcal{M}}^X(x_0)$ in $\langle X(\mathcal{M}), T_{\mathcal{U}} \rangle$. Then there is a $U \in \mathcal{U} \cap \mathcal{M}$ such that $\phi_{\mathcal{M}}^X(B(x_0, U)) \subseteq \mathcal{O}$. It is sufficient to check that there exists a $V \in \mathcal{V} \cap \mathcal{M}$ with $B(x_0, V) \subseteq B(x_0, U)$. To do this we take a $W \in \mathcal{U} \cap \mathcal{M}$ with $2W \subseteq U$ and consider the open cover $\{\text{Int } B(x, W) : x \in X\}$ of $\langle X, T_{\mathcal{U}} \rangle$. For every $x \in X$ there exists a $V_x \in \mathcal{V}$ such that $B(x, 2V_x) \subseteq \text{Int } B(x, W)$. One can assume that the open cover of $\langle X, T_{\gamma} \rangle$ $\{\text{Int } B(x, V_x) : x \in X\}$ is an element of \mathcal{M} . Using the same arguments as above we can find a point $z \in X \cap \mathcal{M}$ and an entourage $V \in \mathcal{V} \cap \mathcal{M}$ such that $x_0 \in B(z, V)$ and $B(z, 2V) \subseteq B(z, W)$. Hence, $B(x_0, V) \subseteq B(z, 2V) \subseteq B(z, W) \subseteq B(x_0, 2W) \subseteq B(x_0, U)$. The proof is now complete. □

3. Corson-compact spaces

A compact Hausdorff space X is called Corson-compact, if it is homeomorphic to a subset of

$$\sum(\mathbb{R}^T) = \left\{ x \in \mathbb{R}^T : \text{supp}(x) \text{ is countable} \right\},$$

where $\text{supp}(x) = \{t \in T : x(t) \neq 0\}$ for $x \in \mathbb{R}^T$, for some set T . Corson-compact spaces have been extensively studied by various authors (for a detailed information see Archangelskij [2], and Negrepointis [8]).

It is easy to verify that for any suitable elementary substructure \mathcal{M} , $\phi_{\mathcal{M}}^X$ maps $\text{cl}(X \cap \mathcal{M})$ homeomorphic on $X(\mathcal{M})$. The main result of the previous paper is the following characterization of Corson-compact spaces.

Theorem 3.1 (Bandlow [4]). *A compact Hausdorff space X is Corson-compact iff for any suitable countable elementary substructure \mathcal{M} , $\phi_{\mathcal{M}}^X$ maps $\text{cl}(X \cap \mathcal{M})$ homeomorphically on $X(\mathcal{M})$.*

4. The construction and $C_p(X)$

Our aim now is to consider the space $C_p(X)$ of all real-valued continuous functions on a completely regular space X in the topology of pointwise convergence. The natural uniformity is given on $C_p(X)$ by all sets of the form

$$\mathcal{U}_{x_1, \dots, x_n}^\varepsilon = \{ \langle f, g \rangle : |f(x_i) - g(x_i)| < \varepsilon, \quad i = 1, \dots, n \},$$

with $x_1, \dots, x_n \in X$, $n \in \mathbb{N}$ and $\varepsilon > 0$.

It is easy to see that two continuous functions f and g on X are \mathcal{M} -equivalent if and only if $f \upharpoonright_{X \cap \mathcal{M}} = g \upharpoonright_{X \cap \mathcal{M}}$.

In connection with $C_p(X)$, we consider on X the uniformity generated by all real-valued continuous functions. There is a natural embedding i from X into the product space $\mathbb{R}^{C(X)}$ and therefore an embedding $i_{\mathcal{M}}$ from $X(\mathcal{M})$ into $\mathbb{R}^{C(X) \cap \mathcal{M}}$.

Let $\overline{X(\mathcal{M})}$ denote the image of X by the projection mapping onto $\mathbb{R}^{\text{cl}(C(X) \cap \mathcal{M})}$. We get the following diagram:

$$(2) \quad \begin{array}{ccc} X & \xrightarrow{i} & \mathbb{R}^{C(X)} \\ \overline{\phi}_{\mathcal{M}}^X \downarrow & & \downarrow \\ \overline{X(\mathcal{M})} & \xrightarrow{\overline{i}_{\mathcal{M}}} & \mathbb{R}^{\text{cl}(C(X) \cap \mathcal{M})} \\ \psi_{\mathcal{M}}^X \downarrow & & \downarrow \\ X(\mathcal{M}) & \xrightarrow{i_{\mathcal{M}}} & \mathbb{R}^{C(X) \cap \mathcal{M}} \end{array}$$

Lemma 4.1. $\psi_{\mathcal{M}}^X$ is a bijection from $\overline{X(\mathcal{M})}$ onto $X(\mathcal{M})$.

PROOF: Let x and y be a pair of different points of X with $\overline{\phi}_{\mathcal{M}}^X(x) \neq \overline{\phi}_{\mathcal{M}}^X(y)$. Then there exists a function $g \in \text{cl}(C_p(X) \cap \mathcal{M})$ such that $g(x) \neq g(y)$. Let $\varepsilon = |g(x) - g(y)|/2$. Then there is a function $f \in C_p(X) \cap \mathcal{M}$ with $|f(x) - g(x)| < \varepsilon$ and $|f(y) - g(y)| < \varepsilon$. Hence, $g(x) \neq g(y)$ and $\phi_{\mathcal{M}}^X(x) \neq \phi_{\mathcal{M}}^X(y)$. \square

Remark 4.2. If X is compact, then $\psi_{\mathcal{M}}^X$ is a homeomorphism and we identify $\overline{X(\mathcal{M})}$ and $X(\mathcal{M})$. It is possible to show that $\psi_{\mathcal{M}}^X$ is a homeomorphism, if X is a Lindelöf p -space or pseudocompact.

If $f : X \rightarrow Y$ is a continuous mapping, then let $f^* : C(Y) \rightarrow C(X)$ denote the mapping induced by f .

Lemma 4.3. $(\overline{\phi}_{\mathcal{M}}^X)^* C_p(\overline{X(\mathcal{M})}) = \text{cl}(C_p(X) \cap \mathcal{M})$.

PROOF: Obviously, we have $C_p(X) \cap \mathcal{M} \subseteq (\phi_{\mathcal{M}}^X)^* C_p(X(\mathcal{M}))$ and $\text{cl}(C_p(X) \cap \mathcal{M}) \subseteq (\overline{\phi}_{\mathcal{M}}^X)^* C_p(\overline{X(\mathcal{M})})$ (see diagram (2)). Since $\psi_{\mathcal{M}}^X$ is a bijection mapping,

$(\psi_{\mathcal{M}}^X)^*C_p(X(\mathcal{M}))$ is a dense subspace of $C_p(\overline{X(\mathcal{M})})$ and therefore $(\phi_{\mathcal{M}}^X)^*C_p(X(\mathcal{M}))$ is a dense subspace of $(\overline{\phi_{\mathcal{M}}^X})^*C_p(\overline{X(\mathcal{M})})$. Now, it is sufficient to show that $C_p(X) \cap \mathcal{M}$ is a dense subset of $(\phi_{\mathcal{M}}^X)^*C_p(X(\mathcal{M}))$. Let g' be an arbitrary continuous function on $X(\mathcal{M})$ and $g = g' \circ \phi_{\mathcal{M}}^X$. Also, let x_1, \dots, x_n be points of X and ε a positive real number. Choose points y_1, \dots, y_k of X such that $\phi_{\mathcal{M}}^X(y_i) \neq \phi_{\mathcal{M}}^X(y_j)$ for distinct $i, j \leq k$ and $\phi_{\mathcal{M}}^X\{x_1, \dots, x_n\} = \phi_{\mathcal{M}}^X\{y_1, \dots, y_k\}$. It is enough to show that there is a function $f \in C_p(X) \cap \mathcal{M}$ with $|f(y_i) - g(y_i)| < \varepsilon$ for all $i = 1, \dots, k$. Since $\phi_{\mathcal{M}}^X(y_i) \neq \phi_{\mathcal{M}}^X(y_j)$ for distinct $i, j \leq k$, we find functionally open neighborhoods V_1, \dots, V_k of y_1, \dots, y_k , respectively, in \mathcal{M} with $\text{cl}(V_i) \cap \text{cl}(V_j) = \emptyset$ for distinct i, j . There exist functions f_1, \dots, f_k in $C_p(X) \cap \mathcal{M}$ such that $V_i = f_i^{-1}(0, 1)$ for every $i = 1, \dots, k$. Fix rational numbers q_1, \dots, q_k such that $|q_i \cdot f_i(y_i) - g(y_i)| < \varepsilon$ for all $i = 1, \dots, k$. Remark that all rational numbers are elements of \mathcal{M} . $f = q_1 \cdot f_1 + \dots + q_k \cdot f_k$ is the desired function. \square

Corollary 4.4. *Let X be a compact Hausdorff space. Then*

$$(\phi_{\mathcal{M}}^X)^*C_p(X(\mathcal{M})) = \text{cl}(C_p(X) \cap \mathcal{M}).$$

A characterization of $C_p(X)$ for Corson-compact spaces

Let X be a completely regular space. We consider X with the uniform structure induced by all real-valued continuous functions on X . The following definition plays the decisive role in what follows.

Definition 5.1. One says that the completely regular space X has the property Ω , if for sufficiently large regular cardinals θ there exists a closed unbounded family C of countable elementary substructures of $\mathcal{H}(\theta)$ such that $\phi_{\mathcal{M}}^X \text{cl}(X \cap \mathcal{M}) = X(\mathcal{M})$ for every $\mathcal{M} \in C$.

More briefly $X \in \Omega$, if $\phi_{\mathcal{M}}^X \text{cl}(X \cap \mathcal{M}) = X(\mathcal{M})$ for every suitable countable elementary substructure \mathcal{M} .

It is easy to verify that every compact Hausdorff space satisfies this condition. More generally, we have the following

Proposition 5.2. *Every Lindelöf p -space has the property Ω .*

PROOF: Since $\phi_{\mathcal{M}}^X(X \cap \mathcal{M})$ is always a dense subset of $X(\mathcal{M})$, it is enough to prove that $\phi_{\mathcal{M}}^X$ is a perfect mapping. X is a Lindelöf p -space if and only if there is a perfect mapping $g : X \rightarrow Y$, where Y has a countable base. By Proposition 2.5 we may assume that g is uniformly continuous. Since $\phi_{\mathcal{M}}^Y$ is a homeomorphism, g can be represented as a composition of $\phi_{\mathcal{M}}^X$ and $g_{\mathcal{M}}$. This implies that $\phi_{\mathcal{M}}^X$ is perfect. \square

Proposition 5.3. *Let X be a Lindelöf space, satisfying Ω . If f is a continuous mapping from X onto a completely regular space Y , then Y satisfies Ω , too.*

PROOF: Like in the proof of previous proposition, we assume that f is uniformly continuous and consider the commutative diagram

$$(3) \quad \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \phi_{\mathcal{M}}^X \downarrow & & \downarrow \phi_{\mathcal{M}}^Y \\ X(\mathcal{M}) & \xrightarrow{f_{\mathcal{M}}} & Y(\mathcal{M}) \end{array}$$

It is easy to see that $f(X \cap \mathcal{M}) = Y \cap \mathcal{M}$ and, consequently, $f(\text{cl}(X \cap \mathcal{M}) \subseteq \text{cl}(Y \cap \mathcal{M}))$. Hence, $Y(\mathcal{M}) = f_{\mathcal{M}}(\phi_{\mathcal{M}}^X \text{cl}(X \cap \mathcal{M})) = \phi_{\mathcal{M}}^Y(\text{cl}(Y \cap \mathcal{M}))$. \square

Theorem 5.4. *If $X \in \Omega$, then each compact subspace of $C_p(X)$ is a Corson-compact space.*

PROOF: Let \mathcal{M} be a suitable countable elementary substructure. Set $Z = C_p(X)$. By Theorem 3.1, it suffices to show that $\phi_{\mathcal{M}}^Z(f) \neq \phi_{\mathcal{M}}^Z(g)$ for any pair $f, g \in \text{cl}(C_p(X) \cap \mathcal{M}) = (\overline{\phi_{\mathcal{M}}^X})^* C_p(\overline{X(\mathcal{M})})$, $f \neq g$. There exist continuous functions f', g' on $\overline{X(\mathcal{M})}$ such that $f = f' \circ \overline{\phi_{\mathcal{M}}^X}$ and $g = g' \circ \overline{\phi_{\mathcal{M}}^X}$. From $X \in \Omega$ it follows that $\overline{\phi_{\mathcal{M}}^X}(X \cap \mathcal{M})$ is a dense subset of $\overline{X(\mathcal{M})}$. Hence, there exists a point $x \in X \cap \mathcal{M}$ with $f'(\overline{\phi_{\mathcal{M}}^X}(x)) \neq g'(\overline{\phi_{\mathcal{M}}^X}(x))$, i.e. $f(x) \neq g(x)$ and therefore $\phi_{\mathcal{M}}^Z(f) \neq \phi_{\mathcal{M}}^Z(g)$. \square

Remark 5.5. A compact subspace of a space $C_p(X)$, where X is an arbitrary compact Hausdorff space, is called an Eberlein space. Every Eberlein space is a Corson-compact space (Amir and Lindenstrauss [1]). Gul'ko proved Theorem 5.4 for Lindelöf Σ -spaces (see Negrepontis [8]), i.e. for continuous images of Lindelöf p -spaces.

Theorem 5.6. *Let X be a compact Hausdorff space. Then X is a Corson-compact space if and only if $C_p(X) \in \Omega$.*

PROOF: The sufficiency follows from Theorem 5.4 and the easy fact that there is a natural embedding of X into $C_p(C_p(X))$.

To prove the necessity, let X be a Corson-compact space and let \mathcal{M} be a suitable countable elementary substructure. By Theorem 3.1, $\phi_{\mathcal{M}}^X$ maps $\text{cl}(X \cap \mathcal{M})$ homeomorphic on $X(\mathcal{M})$. Hence, for every function $f \in C_p(X)$ there is a function $h \in C_p(X(\mathcal{M}))$ such that

$$f \upharpoonright_{\text{cl}(X \cap \mathcal{M})} = (h \circ \phi_{\mathcal{M}}^X) \upharpoonright_{\text{cl}(X \cap \mathcal{M})} .$$

Consequently, $\phi_{\mathcal{M}}^Z(f) = \phi_{\mathcal{M}}^Z(h \circ \phi_{\mathcal{M}}^X)$, where $Z = C_p(X)$. Since, by Corollary 4.4, $h \circ \phi_{\mathcal{M}}^X \in \text{cl}(C_p(X) \cap \mathcal{M})$, this proves that $C_p(X) \in \Omega$. \square

6. Archangelskij’s question

Let D_τ be the discrete space of cardinality τ . Let L_τ denote the space $D_\tau \cup \{\xi\}$, where $\xi \notin D_\tau$ is the only non-isolated point and every neighborhood of ξ has the form $\{\xi\} \cup D_\tau \setminus A$, where A is an arbitrary countable subset of D_τ . It is easy to see that L_τ is a Lindelöf space.

Theorem 6.1 (R. Pol [9], see Archangelskij [2]). *A compact Hausdorff space is Corson-compact if and only if $C_p(X)$ is a continuous image of a closed subspace of $(L_\tau)^\omega$ for some cardinal τ .*

It was asked by Archangelskij [2, Problem IV.3.16], whether X is Corson-compact, if $C_p(X)$ is a continuous image of a closed subspace of $(L_\tau)^\omega \times Z$ for some compact space Z .

Theorem 6.2. *Let Z and X be compact Hausdorff spaces; suppose that $C_p(X)$ can be represented as a continuous image of a closed subspace of $(L_\tau)^\omega \times Z$. Then X is Corson-compact.*

The idea to prove this theorem is very easy. By virtue of Theorem 5.6 and Proposition 5.3, it suffices to show that every closed subspace $Y \subseteq (L_\tau)^\omega \times Z$ satisfies the condition Ω . Remark that $(L_\tau)^\omega$ is always a Lindelöf space (see Step 3).

Step 1. Set $L = L_\tau$. It is easy to see that for every continuous function $f : L \rightarrow \mathbb{R}$ there is a neighborhood $O(\xi) = \{\xi\} \cup D_\tau \setminus A$, where A is a countable subset of D_τ , such that $f(t) = f(\xi)$ for every $t \in O(\xi)$. If \mathcal{M} is a suitable elementary substructure and $f \in \mathcal{M}$, we may assume that $A \in \mathcal{M}$ and, by Proposition 1.2, $A \subset \mathcal{M}$. Hence, by the definition of $\phi_{\mathcal{M}}^L$, $\phi_{\mathcal{M}}^L(\xi) = \phi_{\mathcal{M}}^L(t)$ for every $t \in D_\tau \setminus \mathcal{M}$ and $\phi_{\mathcal{M}}^L(t_1) \neq \phi_{\mathcal{M}}^L(t_2)$ for all $t_1, t_2 \in L \cap \mathcal{M}$, $t_1 \neq t_2$. Consequently, we may identify $L(\mathcal{M})$ with $\{\xi\} \cup (D_\tau \cap \mathcal{M})$. Here all points from $D_\tau \cap \mathcal{M}$ are isolated and the neighborhoods of ξ are of the form $\{\xi\} \cup ((D_\tau \cap \mathcal{M}) \setminus A)$, where A is a countable subset of D_τ and $A \in \mathcal{M}$.

Step 2. Suppose $\eta \in L^\omega \cap \mathcal{M}$. Then, by Proposition 1.2, $\eta(n) \in \mathcal{M}$ for any $n \in \omega$. Hence, $\text{cl}(L^\omega \cap \mathcal{M})$ is the following subspace of L^ω :

$$(\{\xi\} \cup (D_\tau \cap \mathcal{M}))^\omega.$$

The family of all subsets of L^ω of the form

$$A_{\eta_1, \dots, \eta_k}^{n_1, \dots, n_k} = \{\eta \in L^\omega : \eta(n_i) = \eta_i, \quad i = 1, \dots, k\},$$

where $n_1, \dots, n_k \in \omega$ and $\eta_1, \dots, \eta_k \in (\{\xi\} \cup (D_\tau \cap \mathcal{M}))$, is a network for $\text{cl}(L^\omega \cap \mathcal{M})$ in L^ω . This means that for any $\eta \in \text{cl}(L^\omega \cap \mathcal{M})$ and any open set $W \subseteq L^\omega$ with $\eta \in W$ there exists a set $A_{\eta_1, \dots, \eta_k}^{n_1, \dots, n_k}$, such that $\eta \in A_{\eta_1, \dots, \eta_k}^{n_1, \dots, n_k} \subseteq W$. Remark that $A_{\eta_1, \dots, \eta_k}^{n_1, \dots, n_k} \in \mathcal{M}$.

By Proposition 2.1, $L^\omega(\mathcal{M}) \simeq L(\mathcal{M})^\omega$ and $\phi_{\mathcal{M}}^{L^\omega}(\eta) = (\phi_{\mathcal{M}}^L(\eta(n)))_{n \in \omega}$ for every $\eta \in L^\omega$. For short, we write ϕ instead of $\phi_{\mathcal{M}}^{L^\omega}$. Remark that $\phi(\text{cl}(L^\omega \cap \mathcal{M})) = L^\omega(\mathcal{M})$.

Step 3. Now we are going to prove that L^ω is a Lindelöf space for every cardinal τ (see also Arhangelskij [2]). Let γ be an open cover of L^ω . W.l.o.g. we can assume that $\gamma \in \mathcal{M}$. We can also assume that every element of γ is a finite intersection of sets of the form

$$V(n, t) = \{\eta \in L^\omega : \eta(n) = t\},$$

where $n \in \omega$ and $t \in D_\tau$, or of the form

$$W(m, A) = \{\eta \in L^\omega : \eta(m) \in \{\xi\} \cup D_\tau \setminus A\},$$

where $m \in \omega$ and A is a countable subset of D_τ . If $V(n, t) \in \mathcal{M}$, then $t \in \mathcal{M}$ and, analogously, if $W(m, A) \in \mathcal{M}$ and, by Proposition 1.2, $A \subset \mathcal{M}$. Consequently, $(\phi)^{-1}\phi V(n, t) = V(n, t)$ and $(\phi)^{-1}\phi W(m, A) = W(m, A)$. Hence, $(\phi)^{-1}\phi U = U$ for every $U \in \gamma \cap \mathcal{M}$.

Now we claim that $\gamma \cap \mathcal{M}$ is a countable subcover of γ . By the result of Step 2, there exists a system σ of subsets of L^ω satisfying the following conditions:

- (a) $\sigma \subset \mathcal{M}$,
- (b) $\text{cl}(L^\omega \cap \mathcal{M}) \subseteq \cup \sigma$,
- (c) for every $A \in \sigma$ there exists a $U \in \gamma \cap \mathcal{M}$ with $A \subseteq U$.

Hence, $\cup(\gamma \cap \mathcal{M}) = \cup\{(\phi)^{-1}\phi U : U \in \gamma \cap \mathcal{M}\} = (\phi)^{-1}\phi(\cup(\gamma \cap \mathcal{M})) = L^\omega$.

Step 4. Let Z be a compact Hausdorff space and let Y be a closed subspace of $L^\omega \times Z$. Since $L^\omega \times Z$ is a Lindelöf space, Y is Lindelöf, too. By Proposition 2.3, we may think Y with the uniform structure induced by the uniform structure on $L^\omega \times Z$. Let \mathcal{M} be a suitable countable elementary substructure and let h denote the product of the mappings ϕ and $\phi_{\mathcal{M}}^Z$:

$$h : L^\omega \times Z \longrightarrow L(\mathcal{M})^\omega \times Z(\mathcal{M}).$$

It suffices to prove that $h(\text{cl}(Y \cap \mathcal{M})) = h(Y)$ (see Fact 2.3).

Step 5. Let $x = \langle \eta, z \rangle$ be a point of $Y \subseteq L^\omega \times Z$. We define a point $\bar{\eta} \in L^\omega$ by setting $\bar{\eta}(n) = \eta(n)$, if $\eta(n) \in \mathcal{M}$, and $\bar{\eta}(n) = \xi$, if $\eta(n) \notin \mathcal{M}$. Set $A_z = (\phi_{\mathcal{M}}^Z)^{-1}\phi_{\mathcal{M}}^Z(z)$. Remark that $h(\bar{\eta}, z') = h(\eta, z)$ for every $z' \in A_z$. Now, it is enough to prove that $\{\bar{\eta}\} \times A_z \cap \text{cl}(Y \cap \mathcal{M}) \neq \emptyset$.

(a) At first, we prove that $\{\bar{\eta}\} \times A_z \cap Y \neq \emptyset$.

Assume, on the contrary, that $\{\bar{\eta}\} \times A_z \cap Y = \emptyset$. Since $\{\bar{\eta}\} \times A_z$ is compact, there exist open sets $W \subseteq L^\omega$ and $V \subseteq Z$ such that $\bar{\eta} \in W$, $A_z \subseteq V$ and $W \times V \cap Y = \emptyset$. We may assume that $V \in \mathcal{M}$ (see Fact 2.4). Further, assume that W is a member of the canonical base for L^ω . Then there is a natural number n such that W depends only on the coordinates $i \leq n$. Let $\{i_1, \dots, i_k\} = \{i \leq n : \bar{\eta}(i) = \xi\}$, where $i_1 < i_2 < \dots < i_k \leq n$. If $i \in n \setminus \{i_1, \dots, i_k\}$, then $\bar{\eta}(i) = \eta(i) = a_i \in \mathcal{M}$. For every $j \in \{1, \dots, k\}$, we can fix a countable set $A_j \subseteq D_\tau$ such that (w.l.o.g.) W is the set of all $\xi \in L^\omega$ such that $\xi(i) = a_i$ for all $i \leq n$, $i \notin \{i_1, \dots, i_k\}$, and

$\xi(i_j) \notin A_j$ for all $j \in \{1, \dots, k\}$. Let H denote the set of all countable subsets of D_τ . For the sake of simplicity, we define for every collection $B_1, \dots, B_k \in H$

$$W(B_1, \dots, B_k) = \{\xi \in L^\omega : \xi(i) = a_i \text{ for } i \leq n, i \notin \{i_1, \dots, i_k\} \\ \text{and } \xi(i_j) \notin B_j \text{ for } j = 1, \dots, k\}.$$

The following assertion is true:

$$(\exists B_1, \dots, B_k \in H) (W(B_1, \dots, B_k) \times V \cap Y = \emptyset).$$

Since n, i_1, \dots, i_k, V and Y are elements of \mathcal{M} , we find $\overline{B}_1, \dots, \overline{B}_k \in H \cap \mathcal{M}$ satisfying this condition. If $\overline{B}_j \in \mathcal{M}$, then $\overline{B}_j \subset \mathcal{M}$. Since, by the definition of $\overline{\eta}$, $\eta(i_j) \notin D_\tau \cap \mathcal{M}$, $\eta(i_j) \notin \overline{B}_j$ for every $j \in \{1, \dots, k\}$. Now, it is easy to see that $\eta \in W(\overline{B}_1, \dots, \overline{B}_k)$. Hence, $x \in W(\overline{B}_1, \dots, \overline{B}_k) \times V$, contradicting $W(\overline{B}_1, \dots, \overline{B}_k) \times V \cap Y = \emptyset$.

(b) Now, we are going to prove that $\{\overline{\eta}\} \times A_z \cap \text{cl}(Y \cap \mathcal{M}) \neq \emptyset$. In assuming that the intersection is empty, we find open sets $W \subseteq L^\omega$ and $V \subseteq Z$, $V \in \mathcal{M}$, such that $\overline{\eta} \in W$, $A_z \subset V$ and $W \times V \cap \text{cl}(Y \cap \mathcal{M}) = \emptyset$. There exists a natural number n such that

$$C = \{\vartheta \in L^\omega : \vartheta(i) = \overline{\eta}(i) \text{ for } i = 1, \dots, n\} \subseteq W.$$

Remark that $C \in \mathcal{M}$ and $\overline{\eta} \in C$. From $C \subseteq W$ it follows that $C \times V \cap (Y \cap \mathcal{M}) = \emptyset$. Since C, V and Y are elements of \mathcal{M} , this implies that $C \times V \cap Y = \emptyset$. Hence, $\{\overline{\eta}\} \times A_z \cap Y = \emptyset$, contradicting the result of (a). This completes the proof of Theorem 6.2.

REFERENCES

- [1] Amir D., Lindenstrauss J., *The structure of weakly compact sets in Banach spaces*, Ann. Math. Ser. 2 **88:1** (1968).
- [2] Archangelskij A.V., *Topologicheskie prostranstva funkcij* (in Russian), Moscow, 1989.
- [3] Bandlow I., *A construction in set theoretic topology by means of elementary substructures*, Zeitschr. f. Math. Logik und Grundlagen d. Math. **37** (1991).
- [4] ———, *A characterization of Corson-compact spaces*, Comment. Math. Univ. Carolinae **32** (1991).
- [5] Dow A., *An introduction to applications of elementary submodels to topology*, Topology Proceedings, vol. 13, no. 1, 1988.
- [6] Engelking R., *General Topology*, Warsaw, 1977.
- [7] Kunen K., *Set Theory*, Studies in Logic 102, North Holland, 1980.
- [8] Negrepontis S., *Banach spaces and topology*, Handbook of set-theoretic topology, North Holland, 1984, 1045–1042.
- [9] Pol R., *On pointwise and weak topology in function spaces*, Preprint Nr 4/84, Warsaw, 1984.

ERNST-MORITZ-ARNDT UNIVERSITÄT GREIFSWALD, FACHBEREICH MATHEMATIK,
JAHNSTRASSE 15 A, 17489 GREIFSWALD, GERMANY

(Received January 16, 1992, revised November 2, 1993)