

Mariusz Meszka; Roman Nedela; Alexander Rosa
Circulants and the chromatic index of Steiner triple systems

Mathematica Slovaca, Vol. 56 (2006), No. 4, 371--378

Persistent URL: <http://dml.cz/dmlcz/132907>

Terms of use:

© Mathematical Institute of the Slovak Academy of Sciences, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

CIRCULANTS AND THE CHROMATIC INDEX OF STEINER TRIPLE SYSTEMS

MARIUSZ MESZKA* ROMAN NEDELA** — ALEX ROSA***

(Communicated by Martin Škoviera)

ABSTRACT. We complete the determination of the chromatic number of 6-valent circulants of the form $C(n; a, b, a + b)$ and show how this can be applied to improving the upper bound on the chromatic index of cyclic Steiner triple systems.

Introduction

A *Steiner triple system* $STS(v)$ is a pair (V, B) , where V is a set of v points and B is a collection of sets of cardinality 3, called *triples* or *blocks*, satisfying the following condition: each pair x, y of points is contained in exactly one triple. It is well known that a Steiner triple system on v points exists if and only if $v \equiv 1$ or $3 \pmod{6}$.

A *block-color class* is a system of pairwise disjoint triples. An *m -block-coloring* is a partitioning of the set B into m color classes. The *chromatic index* $\chi'(S)$ of a Steiner triple system S is the least m for which an m -block-coloring exists. The *block intersection graph* of a Steiner triple system $S = (V, B)$ is a graph with the vertex set B ; the vertices are adjacent if and only if the respective triples intersect. Since the degree of a block intersection graph equals $3(v-3)/2$, Brooks' theorem (see, e.g., [10]) gives an upper bound $\chi'(S) \leq 3(v-3)/2$ for $v > 7$. An obvious lower bound is $\chi'(S) \geq (v-1)/2$ if $v \equiv 3 \pmod{6}$, and $(v+1)/2$ for $v \equiv 1 \pmod{6}$. Hence we have $(v-1)/2 \leq \chi'(S) \leq 3(v-3)/2$ if $v \equiv 3 \pmod{6}$ and $(v+1)/2 \leq \chi'(S) \leq 3(v-3)/2$ if $v \equiv 1 \pmod{6}$. The lower

2000 Mathematics Subject Classification: Primary 05B07, 05C15.

Keywords: circulant, Steiner triple system, chromatic index.

This research was done while the first and second author were visiting Department of Mathematics and Statistics, McMaster University.

The second author was partially supported by the grant APVV51-009605.

The third author's research was supported by NSERC of Canada Grant No. OGP007268.

bound $\chi'(S) = (v - 1)/2$ is reached if and only if the Steiner triple system is resolvable.

The upper bound $\chi'(S) \leq 3(v - 3)/2$ seems to be weak in general. In fact, using probabilistic methods Pippenger and Spencer in [8] proved that $\chi'(\text{STS}(v))$ is asymptotic to $v/2$. Also no examples of STS with $v > 7$ exceeding the above lower bounds by more than two are known. For more information on the chromatic index of Steiner triple systems the reader is referred to [4: Chap. 18].

For some classes of STS the upper bound was improved. In particular, Colbourn and Colbourn [3] improved it for cyclic STS(v) by proving $\chi'(\text{STS}(v)) \leq v$. A Steiner triple system STS(v) is called *cyclic* if it is isomorphic to the one whose points are $0, 1, \dots, v-1$ and the mapping $i \mapsto i + 1 \pmod{v}$ is an automorphism. The result in [3] is based on the following idea. Let $S = \text{STS}(v)$ be a cyclic Steiner triple system. The block intersection graph has $v(v - 1)/6$ vertices and it admits an induced action of the cyclic group of order v . Then the orbits of the induced action decompose the intersection graph into $(v - 1)/6$ six-valent circulants of order v if $v \equiv 1 \pmod{6}$, and into $(v - 3)/6$ six-valent circulants of order v and one short orbit if $v \equiv 3 \pmod{6}$. By Brooks' theorem each of the circulants (with one exception) can be colored by 6 colors. Taking different sets of colors for different orbits one gets a v -block coloring of any cyclic STS. Using some known results on the chromatic number of toroidal triangulations we derive that, with two exceptions, the 6-valent circulants coming from a cyclic STS(v) can be 5-colored. Consequently, the upper bound for the chromatic index can be improved. Furthermore, we discuss colorability of 6-valent circulants in which one of the three generators is the sum of the other two. The results are applied to get better upper bounds on the chromatic index for some classes of cyclic STS(v).

Coloring 6-valent circulants with connection set $\{a, b, a+b\}$

It is not difficult to see that the induced subgraph of an orbit of length v in a Steiner triple system STS(v) in the block intersection graph is a circulant $G = C(n; a, b, c)$, where one of the generators, say c , is the sum of the others (up to replacing c by $n - c$). In what follows we assume w.l.o.g. $c = a + b$. We may think of each edge as being colored by one of the generators. A triangle of G is *3-edge-colored* if its edges are colored by the three colors a, b and c . Each edge lies in exactly two 3-edge-colored triangles. The six 3-edge-colored triangles based at a vertex j induce a cyclic rotation of the neighbours; up to replacing some of a, b, c by $-a, -b, -c$, it has a form $(j+a, j+b, j+c, j-a, j-b, j-c)$ for each vertex $j \in Z_n$. Clearly the 3-edge-colored cycles as well as the local

rotation of colors $(a, b, c, -a, -b, -c)$ are preserved by the cyclic automorphism $j \mapsto j + 1$. By Euler-Poincare theorem a component of G is a one-skeleton of a vertex-transitive triangulation of the torus. Note that there are $n/\gcd(a, b, c)$ components of G , and they are all isomorphic to $C = (n'; a', b', c')$, where $n' = n/d, a' = a/d, b' = b/d, c' = c/d$ for $d = \gcd(a, b, c)$. Many papers were devoted to the problem to determine the chromatic number of triangulations of surfaces satisfying certain conditions. Here we mention only few of those which are closely related to our problem.

Note that the complete graph K_7 can be expressed as $K_7 \cong C(7; 1, 2, 3)$, and let T_{11} be the circulant $C(11; 1, 2, 3)$. It is important to note that the chromatic number of K_7 is seven and the chromatic number of T_{11} is 6. Albertson and Hutchinson [1] (see [9] as well) proved that these two graphs are the only 6-valent graphs on the torus which cannot be 5-colored.

THEOREM 1. ([1; Theorem 4.4]) *Let G be a toroidal 6-valent graph other than K_7 and T_{11} . Then G is 5-colorable.*

COROLLARY 2. *Let $S = \text{STS}(v)$ be a cyclic Steiner triple system. Then*

- (a) $\chi'(S) \leq \frac{5v+13}{6}$ if $v \equiv 1 \pmod{6}$,
- (b) $\chi'(S) \leq \frac{5v+9}{6}$ if $v \equiv 3 \pmod{6}$.

Proof. As already mentioned, the vertices of the block intersection graph decompose into $(v-1)/6$ orbits of length v if $v \equiv 1 \pmod{6}$, and into $(v-3)/6$ orbits of length v and one short orbit if $v \equiv 3 \pmod{6}$. The subgraph induced on a full orbit is a six-valent circulant $C(v; a, b, c)$ with $c = a + b$ for some a, b . A connectivity component of such a circulant triangulates the torus. By Theorem 1 each such circulant can be 5-colored except when it is a disjoint union of K_7 's, or a disjoint union of T_{11} 's. This can happen only if $7|v$ or $11|v$, respectively. Moreover, at most one orbit can be a union of K_7 's and at most one orbit can be a union of T_{11} 's. We color the vertices of the block intersection graph by using distinct sets of five colors for each full orbit which is neither a union of K_7 's, nor a union of T_{11} 's. For the latter two orbits, if they appear, we use 7 and 6 colors, respectively. Finally, in case $v \equiv 3 \pmod{6}$ we use one color to color the short orbit. By adding the number of colors used we get the upper bounds. □

The bound in Corollary 2(b) cannot be improved in general by the method used in the proof of Corollary 2. To see this, let $v = 3p, p$ a prime, such that

- (*) $p \equiv 3 \pmod{4}$, x is such that the order of $x \pmod{p}$ is $\frac{p-1}{2}$, and the order of $x \pmod{3p}$ is $p-1$; such x always exists.

Now, for each prime p and x satisfying (*) and $v = 3p$. there exist cyclic Steiner triple systems of order v such that for each full orbit of triples the

corresponding 6-valent circulant is 5-chromatic, so that the bound of Corollary 2 becomes $\chi'(S) \leq \frac{5v-9}{6}$. Indeed, the set $\{x^{2i}, 2 \cdot x^{2i}, 3 \cdot x^{2i} : i = 0, 1, \dots, \frac{v-3}{6}\}$ is a solution to the 2nd Heffter's difference problem (cf. [4]); note the crucial fact that $\frac{v-1}{2}$ is odd. Clearly, all circulants corresponding to the full orbits of any cyclic STS(v) whose difference triples are as above are multiplier-isomorphic to $C(v; 1, 2, 3)$.

The first few values of v satisfying condition (*) with $x = 2$ are $v = 21, 69, 141, 213, 237, 309, 501, \dots$. There are infinitely many primes p and corresponding x satisfying (*). Let us note that there exist cyclic STSs also of orders $\equiv 1 \pmod{6}$ (e.g. $v = 37$) for which the circulants corresponding to any full orbit are 5-chromatic.

TABLE

Order	Spectrum
7	7
9	-
13	10
15	9
19	12, 15
21	13, 14, 16
25	18, 20
27	18
31	20, 21, 22, 23
33	21, 22, 23, 25, 26
37	24, 25, 26, 27, 28, 30
39	23, 24, 25, 26, 27, 28, 29
43	28, 29, 30, 31, 32, 33, 34
45	26, 27, 28, 29, 30, 31

On the other hand, for many cyclic STSs our method gives a much better upper bound. For example, for the projective STS of order 31 (whose blocks are the lines of $\text{PG}(4, 2)$), the bound of Corollary 2 yields the value of 20. We found an 18-block-colouring of $\text{PG}(4, 2)$, which implies that the chromatic index of

$PG(4, 2)$ equals 18 ($\chi' \geq 18$ follows from Wilson's argument, see [4; p. 345]). The spectrum of values for bounds obtained by adding the chromatic numbers of circulants corresponding to the individual orbits of STSs of small orders is given in the table above.

Let $\min_v = \frac{v-1}{2}$ if $v \equiv 3 \pmod{6}$ and $\min_v = \frac{v+1}{2}$ if $v \equiv 1 \pmod{6}$. We checked on a computer that for all cyclic Steiner triple systems of order $v \leq 43$ (note that the number of cyclic Steiner triple systems of order 43 is known to be 9508, not a small number [4]), we have for the chromatic index the bounds $\min_i \leq \chi' \leq \min_v + 2$. This suggests that the following could be true:

CONJECTURE. *The chromatic index of every cyclic STS(v), $v > 7$, takes on one of three values: \min_v , $\min_v + 1$, $\min_v + 2$.*

It would be tempting to conjecture that the same holds for any Steiner triple system but admittedly the evidence for such a claim is scarce.

Let us note that the problem of determining the chromatic number of general circulants is NP-hard ([2]). However, we are able to say more in the special case that we consider, namely for the 6-valent circulants $C(n; a, b, a+b)$.

Some recent results proved by Collins and Hutchinson [6] and in more general setting by Hutchinson, Richter and Seymour [7] suggest that most of the considered circulants are even 4-colorable. Indeed, Yeh and Zhu [11; Theorem 5] proved the following statement conjectured by Collins, Fisher and Hutchinson in [5]:

THEOREM 3. (Yeh, Zhu [11]) *Let $G = C(n; 1, r, r+1)$, $n \geq 2(r+1) \geq 6$, be a circulant (possibly with multiple edges for $n = 2(r+1)$). Then G is 4-colorable except when one of the following cases occurs:*

- (1) $n = 2r + 2$, $n = 2r + 3$, $n = 3r + 1$ or $3r + 2$,
and n is not divisible by 4,
- (2) $r = 2$, and 4 does not divide n ,
- (3) (r, n) is one of $(3, 13)$, $(3, 17)$, $(3, 18)$, $(3, 25)$, $(4, 17)$, $(6, 17)$, $(6, 25)$, $(6, 33)$, $(7, 19)$, $(7, 25)$, $(7, 26)$, $(9, 25)$, $(10, 25)$, $(10, 26)$, $(10, 37)$, $(14, 33)$.

Using the above theorem one can classify k -chromatic 6-valent circulants for all admissible integers k , $3 \leq k \leq 7$, as follows.

THEOREM 4. *Let $G = C(n; a, b, c)$ be a connected 6-valent circulant, where $n \geq 7$, $c = a + b$ or $n - c = a + b$ are pairwise distinct positive integers different from $n/2$. Let $\chi(G)$ be the chromatic number of G . Then*

- (1) $\chi(G) = 7$ if and only if $G \cong K_7 \cong C(7; 1, 2, 3)$,
- (2) $\chi(G) = 6$ if and only if $G \cong T_{11} \cong C(11; 1, 2, 3)$,

- (3) $\chi(G) = 5$ if and only if $G \cong C(n; 1, 2, 3)$ and $n \neq 7, 11$ is not divisible by 4, or G is isomorphic to one of the following circulants: $C(13; 1, 3, 4)$, $C(17; 1, 3, 4)$, $C(18; 1, 3, 4)$, $C(19; 1, 7, 8)$, $C(25; 1, 3, 4)$, $C(26; 1, 7, 8)$, $C(33; 1, 6, 7)$, $C(37; 1, 10, 11)$.
- (4) $\chi(G) = 3$ if and only if $3|n$ and none of a, b, c is divisible by 3.
- (5) $\chi(G) = 4$ in all the remaining cases.

Proof.

(1) follows from Brooks' theorem.

(2) comes from Albertson and Hutchinson [1]. It is easy to see that none of the circulants listed in (3) is 4-colorable. Assume G is not 4-colorable. If $\gcd(n, a) = 1$, $\gcd(n, b) = 1$ or $\gcd(n, c) = 1$, then G is (multiplier-)isomorphic to $C(n; 1, r, r+1)$. Applying Theorem 3 and reducing the number of exceptions by taking representatives up to isomorphism we get the list of 5-chromatic six-valent circulants. Assume the greatest common divisors $\gcd(n, a)$, $\gcd(n, b)$ and $\gcd(n, c)$ to be all > 1 . We show that in this case G can be 4-colored. Let, say, $\gcd(n, a) = d > 1$. Since each of the pairs (a, b) , (b, c) , (a, c) generates Z_n , it follows that at most one of $\gcd(a, n)$, $\gcd(b, n)$ and $\gcd(c, n)$ is even. Hence we may assume that both $\gcd(a, n) = d$ and $\gcd(b, n) = k$ are odd. Clearly $db \in \langle a \rangle$, hence $db = xa$ for some x , $0 \leq x \leq m - 1$, where $m = n/d$. If $\gcd(x, m) = 1$, then $k = \gcd(n, b) = 1$, which is a contradiction. Moreover, $\gcd(m, x) \neq 2$ because $\gcd(m, x) = 2$ would imply $|b| = n/2 = n/\gcd(b, n)$, forcing $\gcd(b, n) = 2$ contradicting the assumptions. Hence, $\gcd(m, x) > 2$. Let $x = 0$. Then the circulant G forms a so-called right diagonal $m \times d$ grid, $m, d \geq 3$, which by [6; Theorem 3.3] is 4-colorable. If $x \neq 0$, then consider the factor of G by $\langle xa \rangle$. It is a 6-valent circulant $\bar{G} = C(\bar{n}; \bar{a}, \bar{b}, \bar{c})$ such that one of the generators is the sum of the other two, $\langle \bar{a} \rangle$ is a subgroup of index d and of cardinality $\bar{m} = \bar{n}/d \geq 3$. Moreover, $d\bar{b} = 0$, and consequently, the order of \bar{b} is d . As above, the 4-colorability of \bar{G} follows from [6; Theorem 3.3]. The 4-coloring of the original circulant G can be constructed by lifting the 4-coloring of \bar{G} along the covering $G \rightarrow \bar{G}$.

If $3|n$ and none of the generators is divisible by 3, then the assignment $j \mapsto \bar{j}$, where \bar{j} is the residue class of $j \pmod 3$, defines a 3-coloring of the circulant G .

Assume there is a 3-coloring of G . If one of the $\gcd(n, a)$, $\gcd(n, b)$ or $\gcd(n, c)$ equals one, then the circulant is isomorphic to $C(n; 1, r, r+1)$ for some $1 < r < r + 1 < n/2$. The vertices $0, r, r + 1$ are colored by three different colors, say A, B and C , respectively. It follows that the vertex 1 is colored by B , vertex $r + 2$ by A, \dots . It is easy to see that this coloring extends to a coloring of G if and only if $3|n$ and $r \equiv 1 \pmod 3$. Hence, all the generators are not $\equiv 0 \pmod 3$, and this property remains valid when multiplying by integers coprime to n as well as when replacing generators by their inverses. Assume now that none of $\gcd(n, a)$, $\gcd(n, b)$, $\gcd(n, c)$ equals one. If, say, $\gcd(n, a) = d > 1$,

then considering 3-colorability of the subgraph H induced by $\langle a \rangle \cup (b + \langle a \rangle)$ we see that H can be 3-colored only if $3 \mid n/d$. Hence $3 \mid n$. Moreover, the 3-coloring is unique up to a permutation of colors. Since a is a generator of a cyclic subgroup of order divisible by 3, it is coprime to 3. The same argument applies to b and c , proving that all a , b and c are coprime to 3. \square

Let us call circulants of type (1), (2) or (3) from Theorem 4 *exceptional*.

COROLLARY 5. *Let $S = \text{STS}(v)$ be a cyclic Steiner triple system of order v , and assume that no circulant induced on an orbit of the block intersection graph contains as a component an exceptional circulant. Then $\chi'(S) \leq \frac{2}{3}(v-1)$ if $v \equiv 1 \pmod{6}$, and $\chi'(S) \leq \frac{2}{3}v - 1$ if $v \equiv 3 \pmod{6}$.*

Acknowledgement

Thanks to Martin Mačaj for valuable suggestions concerning the impossibility of an improvement of Corollary 2(b). We would like to thank the Department of Mathematics and Statistics, McMaster University for its hospitality.

REFERENCES

- [1] ALBERTSON, M. HUTCHINSON, J.: *On six-chromatic toroidal graphs*, Proc. London Math. Soc. (3) **41** (1980), 533–556.
- [2] CODENOTTI, B. GERACE, I.—VIGNA, S.: *Hardness results and spectral techniques for combinatorial problems on circulant graphs*, Linear Algebra Appl. **285** (1998), 123–142.
- [3] COLBOURN, C.—COLBOURN, M.: *The chromatic index of cyclic Steiner 2-designs*, Int. J. Math. Math. Sci. **5** (1982), 823–825.
- [4] COLBOURN, C. ROSA, A.: *Triple Systems*, Clarendon Press, Oxford, 1999.
- [5] COLLINS, K. FISHER, D. HUTCHINSON, J.: *On 3- and 4-coloring some circulant graphs*. Presented at DREI'98. Graph Theory and Combinatorial Optimization, August 1998.
- [6] COLLINS, K. HUTCHINSON, J.: *Four-coloring six-regular graphs on the torus*. In: CRM Proc. Lecture Notes 23, Amer. Math. Soc., Providence, RI, 1999, pp. 21–33.
- [7] HUTCHINSON, J. RICHTER, R. B. SEYMOUR, P.: *Coloring Eulerian triangulations*, J. Combin. Theory Ser. B **84** (2002), 225–239.
- [8] PIPPENGER, N. SPENCER, J.: *Asymptotic behavior of the chromatic index for hypergraphs*, J. Combin. Theory Ser. A **51** (1989), 24–42.
- [9] THOMASSEN, C.: *Five-coloring maps on surfaces*, J. Combin. Theory Ser. B **59** (1993), 89–105.
- [10] WEST, D.: *Introduction to Graph Theory* (2nd ed.), Prentice Hall of India, New Delhi, 2005.

- [11] YEH, H.-G.—ZHU, X.: *4-colorable 6-regular toroidal graphs*, *Discrete Math.* **273** (2003), 261–274.

Received October 29, 2004

* *Faculty of Applied Mathematics*
AGH University of Science and Technology
PL 30059 Kraków
POLAND

** *Mathematical Institute*
Slovak Academy of Sciences
SK-974 00 Banská Bystrica
SLOVAKIA
E-mail: nedela@savbb.sk

*** *Department of Mathematics and Statistics*
McMaster University
Hamilton
Ontario
CANADA