

A. Mokhtari; Toufik Moussaoui; D. O'Regan

Existence and multiplicity of solutions for a $p(x)$ -Kirchhoff type problem via variational techniques

Archivum Mathematicum, Vol. 51 (2015), No. 3, 163–173

Persistent URL: <http://dml.cz/dmlcz/144427>

Terms of use:

© Masaryk University, 2015

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

EXISTENCE AND MULTIPLICITY OF SOLUTIONS
FOR A $p(x)$ -KIRCHHOFF TYPE PROBLEM
VIA VARIATIONAL TECHNIQUES

A. MOKHTARI, T. MOUSSAOUI, AND D. O'REGAN

ABSTRACT. This paper discusses the existence and multiplicity of solutions for a class of $p(x)$ -Kirchhoff type problems with Dirichlet boundary data of the following form

$$\begin{cases} -\left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = f(x, u), & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a smooth open subset of \mathbb{R}^N and $p \in C(\overline{\Omega})$ with $N < p^- = \inf_{x \in \Omega} p(x) \leq p^+ = \sup_{x \in \Omega} p(x) < +\infty$, a, b are positive constants and $f: \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function. The proof is based on critical point theory and variable exponent Sobolev space theory.

1. INTRODUCTION

In this paper we study

$$(1) \quad \begin{cases} -\left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = f(x, u), & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded domain of \mathbb{R}^N with smooth boundary $\partial\Omega$ and $N \geq 1$, $p \in C(\overline{\Omega})$ with $N < p^- = \inf_{x \in \Omega} p(x) \leq p^+ = \sup_{x \in \Omega} p(x) < +\infty$, a, b are positive constants and $f: \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function.

Problem (1) is related for example to vibrations and deformations of plates or tended cords. The operator $-\operatorname{div}(|\nabla u|^{p(x)-2} \nabla u)$ is said to be the $p(x)$ -Laplacian, and becomes the p -Laplacian when $p(x) = p$. The variable exponent Sobolev space $W_0^{1,p(x)}(\Omega)$ is a natural generalization of the classical Sobolev space $W_0^{1,p}(\Omega)$ and related preliminaries concerning $W_0^{1,p(x)}(\Omega)$ and $p(x)$ -Laplacian equations will be given in Section 2.

2010 *Mathematics Subject Classification*: primary 34B27; secondary 35J60, 35B05.

Key words and phrases: existence results, genus theory, nonlocal problems Kirchhoff equation, critical point theory.

Received March 18, 2015. Editor V. Müller.

DOI: 10.5817/AM2015-3-163

In this paper we examine existence and multiplicity of solutions of the $p(x)$ -Kirchhoff equation associated to problem (1) by applying a minimization principle and the genus theory introduced by Krasnoselskii (see [4], [13]). Problem (1) is related to the stationary problem of a model introduced by Kirchhoff [12]. More precisely, Kirchhoff proposed a model given by the equation

$$\rho \frac{\partial^2 u}{\partial t^2} - \left(\frac{\rho_0}{h} + \frac{E}{2L} \int_0^L \left| \frac{\partial u}{\partial x} \right|^2 dx \right) \rho \frac{\partial^2 u}{\partial x^2} = 0,$$

G. Dai and J. Wei established the existence of infinitely many nonnegative solutions for problem (1) by applying a general variational principle due to B. Ricceri (see Theorem 3.1 and Theorem 3.2 in [8]). In this paper, we first prove the existence of a solution to problem (1), by using a minimization principle with the hypothesis $\limsup_{|u| \rightarrow +\infty} \frac{F(x,t)}{|u|^{\theta(x)}} \leq a(x)$, where $F(x, u) = \int_0^u f(x, t) dt$ and a is a measurable function which belongs to $L^\infty(\Omega)$ and $\theta \in C(\bar{\Omega})$ satisfies $\theta^+ = \sup_{\bar{\Omega}} \theta(x) < 2p^-$. Our second result gives the existence and multiplicity of solutions using Clarke's theorem under the following assumptions:

- there exists positive constants $C_1, C_2 > 0$ and a function q measurable on Ω such that $C_1 t^{q(x)-1} \leq f(x, t) \leq C_2 t^{q(x)-1}$ for all $t \geq 0$ and for all $x \in \Omega$, where $q \in L^\infty(\Omega)$ and $1 < q^- = \text{essinf}_\Omega q(x) \leq q(x) \leq q^+ = \text{esssup}_\Omega q(x) < p^- = \text{essinf}_\Omega p(x)$,
- $f(x, -t) = -f(x, t)$ for all $t \in \mathbb{R}$ and for all $x \in \bar{\Omega}$.

These hypotheses are a generalization of the hypotheses introduced in [7].

This paper is organized as follows. In Section 2, we present some necessary preliminaries on variable exponent Lebesgue and Sobolev spaces and we recall some definitions and basic properties of the Krasnoselskii genus. In Section 3, using critical point theory, we establish existence and multiplicity results for problem (1).

2. PRELIMINARIES

Suppose that Ω is a smooth bounded open domain of \mathbb{R}^N with a smooth boundary $\partial\Omega$ and $p \in C(\bar{\Omega})$ satisfies

$$1 < p^- \doteq \inf_{x \in \Omega} p(x) \leq p^+ \doteq \sup_{x \in \Omega} p(x) < +\infty.$$

The variable exponent Lebesgue space $L^{p(x)}(\Omega)$ is defined by

$$L^{p(x)}(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ is measurable, } \int_\Omega |u(x)|^{p(x)} dx < +\infty \right\}$$

endowed with the norm

$$|u|_{p(x)} = \inf \left\{ \lambda > 0 : \int_\Omega \left| \frac{u(x)}{\lambda} \right|^{p(x)} dx \leq 1 \right\}.$$

Define the variable exponent Sobolev space $W^{1,p(x)}(\Omega)$ by

$$W^{1,p(x)}(\Omega) = \{ u \in L^{p(x)}(\Omega) : |\nabla u| \in L^{p(x)}(\Omega) \}$$

endowed with the norm $\|u\|_{1,p(x)} = |u|_{p(x)} + |\nabla u|_{p(x)}$.

Denote by $C(\overline{\Omega})$ the space of continuous functions on $\overline{\Omega}$ endowed with the norm $|u|_\infty = \sup_{x \in \overline{\Omega}} |u(x)|$. $W_0^{1,p(x)}(\Omega)$ denotes the closure of $C_0^\infty(\Omega)$ in $W^{1,p(x)}(\Omega)$.

Proposition 2.1 (See [10]). *$L^{p(x)}(\Omega)$, $W_0^{1,p(x)}(\Omega)$ and $W^{1,p(x)}(\Omega)$ are separable, reflexive and uniformly convex Banach spaces.*

Proposition 2.2 (See [10]). *Let $\rho(u) = \int_\Omega |u(x)|^{p(x)} dx$. For any $u, u_k \in L^{p(x)}(\Omega)$, $k = 1, 2, \dots$, we have*

- (1) *For $u \neq 0$, $|u|_{p(x)} = \lambda \Leftrightarrow \rho(\frac{u}{\lambda}) = 1$.*
- (2) *$|u|_{p(x)} < 1 (= 1; > 1) \Leftrightarrow \rho(u) < 1 (= 1; > 1)$.*
- (3) *If $|u|_{p(x)} > 1$, then $|u|_{p(x)}^{p^-} \leq \rho(u) \leq |u|_{p(x)}^{p^+}$.*
- (4) *If $|u|_{p(x)} < 1$, then $|u|_{p(x)}^{p^+} \leq \rho(u) \leq |u|_{p(x)}^{p^-}$.*
- (5) *$\lim_{k \rightarrow +\infty} |u_k|_{p(x)} = 0 \Leftrightarrow \lim_{k \rightarrow +\infty} \rho(u_k) = 0$.*
- (6) *$\lim_{k \rightarrow +\infty} |u_k|_{p(x)} = +\infty \Leftrightarrow \lim_{k \rightarrow +\infty} \rho(u_k) = +\infty$.*

Proposition 2.3 (See [10]). *If $u, u_k \in L^{p(x)}(\Omega)$, $k = 1, 2, \dots$, then the following statements are equivalent to each other:*

- (1) *$\lim_{k \rightarrow +\infty} |u_k - u|_{p(x)} = 0$ (i.e. $u_k \rightarrow u$ in $L^{p(x)}(\Omega)$).*
- (2) *$\lim_{k \rightarrow +\infty} \rho(u_k - u) = 0$.*
- (3) *$u_k \rightarrow u$ in measure in Ω and $\lim_{k \rightarrow +\infty} \rho(u_k) = \rho(u)$.*

Proposition 2.4 (See [9]). *The Poincaré-type inequality holds, that is, there exists a positive constant c_Ω such that*

$$|u|_{p(x)} \leq c_\Omega |\nabla u|_{p(x)}, \quad \forall u \in W_0^{1,p(x)}(\Omega).$$

Thus $|\nabla u|_{p(x)}$ is an equivalent norm in $W_0^{1,p(x)}(\Omega)$. We will use this equivalent norm in the following discussion and write $\|u\| = |\nabla u|_{p(x)}$ for simplicity.

We now recall the Krasnoselskii genus and more information on this subject may be found in ([11], [1], [4], [13]). Let E be a real Banach space. Let us denote by Σ the class of all closed subsets $A \subset E - \{0\}$ that are symmetric with respect to the origin, that is, $u \in A$ implies $-u \in A$.

Definition 2.1. Let $A \in \Sigma$. The Krasnoselskii genus $\gamma(A)$ is defined as being the least positive integer n such that there is an odd mapping $\varphi \in C(A, \mathbb{R}^n - \{0\})$. If such an n does not exist we set $\gamma(A) = +\infty$. Furthermore, by definition, $\gamma(\emptyset) = 0$.

Theorem 2.1 (See [11]). *Let $E = \mathbb{R}^N$ and $\partial\Omega$ be the boundary of an open, symmetric and bounded subset $\Omega \subset \mathbb{R}^N$ with $0 \in \Omega$. Then $\gamma(\partial\Omega) = N$.*

Note $\gamma(S^{N-1}) = N$. If E is of infinite dimension and separable and S is the unit sphere in E , then $\gamma(S) = +\infty$.

Proposition 2.5 (See [11]). *Let $A, B \in \Sigma$. Then:*

- if there exists an odd map $f \in C(A, B)$, then $\gamma(A) \leq \gamma(B)$. Consequently, if there exists an odd homeomorphism $f: A \rightarrow B$, then $\gamma(A) = \gamma(B)$.
- if $A \subset B$, then $\gamma(A) \leq \gamma(B)$.
- $\gamma(A \cup B) \leq \gamma(A) + \gamma(B)$.

Theorem 2.2 (Minimization principle). *Let X be a real reflexive Banach space. If the functional $J: X \rightarrow \mathbb{R}$ is weakly lower semi-continuous and coercive (i.e. $\lim_{\|u\| \rightarrow +\infty} J(u) = +\infty$), then there exists $u_0 \in X$ such that $J(u_0) = \inf_{u \in X} J(u)$. Moreover, if J is also Gateaux differentiable on X , then $J'(u_0) = 0$.*

Definition 2.2. Let $J \in C^1(X, \mathbb{R})$. If any sequences $(u_n) \subset X$ for which $(J(u_n))$ is bounded and $J'(u_n) \rightarrow 0$ when $n \rightarrow +\infty$ in X' possesses a convergent subsequence, then we say that J satisfies the Palais-Smale condition (denoted by the (P-S) condition).

We now state a theorem due to Clarke.

Theorem 2.3 (See [5], [15]). *Let $J \in C^1(E, \mathbb{R})$ be a functional satisfying the Palais-Smale condition. Also suppose that:*

- J is bounded from below and even;
- there is a compact set $K \in \Sigma$ such that $\gamma(K) = k$ and $\sup_{x \in K} J(x) < J(0)$.

Then J possesses at least k pairs of distinct critical points and their corresponding critical values are less than $J(0)$.

3. MAIN RESULTS

In this section we will discuss the existence of weak solutions of (1).

Definition 3.1. We say that $u \in W_0^{1,p(x)}(\Omega)$ is a weak solution of problem (1) if and only if

$$\left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx \right) \int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla v dx = \int_{\Omega} f(x, u) v dx,$$

for all $v \in W_0^{1,p(x)}(\Omega)$.

The energy functional corresponding to problem (1) is defined as follows,

$$J(u) = a \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \frac{b}{2} \left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx \right)^2 - \int_{\Omega} F(x, u) dx$$

where $F(x, t) = \int_0^t f(x, s) ds$. It is easy to see that $J \in C^1(W_0^{1,p(x)}(\Omega), \mathbb{R})$ and for all $u, v \in W_0^{1,p(x)}(\Omega)$

$$J'(u) \cdot v = \left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx \right) \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)-2} \nabla u \nabla v dx - \int_{\Omega} f(x, u) v dx.$$

Thus the critical points of J are the weak solutions of (1).

Theorem 3.1. *Assume that*

$$(H1) \quad \limsup_{|u| \rightarrow +\infty} \frac{F(x, u)}{|u|^{\theta(x)}} \leq a(x),$$

where $\theta \in C(\overline{\Omega})$ with $\theta^- = \inf_{x \in \overline{\Omega}} \theta(x) > 1$ and $a \in L^\infty(\Omega)$. If $\theta^+ = \sup_{x \in \overline{\Omega}} \theta(x) < 2p^-$, then (1) has a weak solution.

Proof. From the continuity of F and assumption (H1) we deduce that there exists a positive constant C such that

$$F(x, u) \leq a(x)|u|^{\theta(x)} + C, \quad \forall u \in \mathbb{R}, \forall x \in \Omega.$$

We have for $\|u\| > 1$ that

$$\begin{aligned} J(u) &= a \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \frac{b}{2} \left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx \right)^2 - \int_{\Omega} F(x, u) dx \\ &\geq \frac{a}{p^+} \rho_p(\nabla u) + \frac{b}{2(p^+)^2} (\rho_p(\nabla u))^2 - \int_{\Omega} (a(x)|u|^{\theta(x)} + C) dx, \\ &\geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - C \operatorname{meas}(\Omega) - |a|_{L^\infty} \int_{\Omega} |u|^{\theta(x)} dx. \end{aligned}$$

Since $p^- > N$, the embedding $W_0^{1,p(x)} \hookrightarrow C(\overline{\Omega})$ is continuous and we see that

$$\int_{\Omega} |u|^{\theta(x)} dx \leq \int_{\Omega} |u|_{\infty}^{\theta(x)} dx \leq C_0 \int_{\Omega} \|u\|^{\theta(x)} \leq C_0 \|u\|^{\theta^+},$$

where $C_0 = \max_{x \in \overline{\Omega}} C^{\theta(x)}$ and C is the constant of the embedding, and this implies that

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - C \operatorname{meas}(\Omega) - |a|_{L^\infty} C_0 \|u\|^{\theta^+}.$$

Since $\theta^+ < 2p^-$ then J is coercive. Now J is also weakly lower semicontinuous (see Theorem 3.2.9 in [9]), so we see that J has a global minimum point $u \in W_0^{1,p(x)}(\Omega)$, which is a weak solution to problem (1). \square

Example 3.1. We consider $\Omega = (0, 1)$ and $f(x, t) = a(x)\theta(x)|t|^{\theta(x)-2}t$, for all $t \in \mathbb{R}$, and we put for $x \in \Omega$, $a(x) = \sin x$, $\theta(x) = x + \frac{3}{2}$ and $p(x) = x^2 + 2$. Our problem becomes

$$\begin{cases} -\left(a + b \int_0^1 \frac{1}{x^2+2} |u'|^{x^2+2} dx\right) (|u'|^{x^2} u')' = \frac{\sin x}{x^2+2} |u|^{-\frac{2x^2+3}{x^2+2}} u, & \text{in } (0, 1) \\ u(0) = u(1) = 0, \end{cases}$$

and it admits at least one weak solution.

Remark 3.1. To obtain a nontrivial solution, we may assume that $f(x_0, 0) \neq 0$ for some $x_0 \in [0, 1]$.

Theorem 3.2. *Assume that:*

(H2) *there exist positive constants $C_1, C_2 > 0$ and a function q measurable on Ω such that $C_1 t^{q(x)-1} \leq f(x, t) \leq C_2 t^{q(x)-1}$ for all $t \geq 0$ and for all $x \in \Omega$, where $q \in L^\infty(\Omega)$ and $1 < q^- = \operatorname{ess\,inf}_{x \in \Omega} q(x) \leq q(x) \leq q^+ = \operatorname{ess\,sup}_{x \in \Omega} q(x) < p^- = \inf_{x \in \Omega} p(x)$,*

(H3) *$f(x, -t) = -f(x, t)$ for all $t \in \mathbb{R}$ and for all $x \in \bar{\Omega}$.*

Then (1) has infinitely many weak solutions.

For the proof of Theorem 3.2 we will need the following steps.

Step 1: J is bounded from below.

Indeed, for any $u \in W_0^{1,p(x)}(\Omega)$, we have

$$J(u) \geq \frac{a}{p^+} \int_{\Omega} |\nabla u|^{p(x)} \, dx + \frac{b}{2(p^+)^2} \left(\int_{\Omega} |\nabla u|^{p(x)} \, dx \right)^2 - \frac{C_2}{q^-} \int_{\Omega} |\nabla u|^{q(x)} \, dx.$$

Let $\rho_p(u) = \int_{\Omega} |u|^{p(x)} \, dx$ and $\rho_q(u) = \int_{\Omega} |u|^{q(x)} \, dx$. We have the following four cases:

(i) If $\rho_p(u) > 1$ and $\rho_q(u) < 1$, then by Proposition 2.2, we obtain

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - \frac{C_2}{q^-} |u|_{q(x)}^{q^-}.$$

Since $q(x) \leq p(x)$ for all $x \in \Omega$,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - \frac{C_2 C_3}{q^-} |u|_{p(x)}^{q^-},$$

where C_3 is the constant of the continuous embedding of $L^{p(x)}$ in $L^{q(x)}$.

Using the Poincaré inequality, we have

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - \frac{C_2 C_3 c_{\Omega}^{q^-}}{q^-} \|u\|_{p(x)}^{q^-}.$$

We note that $2p^- > q^-$, so J is bounded from below.

(ii) If $\rho_p(u) > 1$ and $\rho_q(u) > 1$ then,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - \frac{C_2 C_3 c_{\Omega}^{q^-}}{q^-} \|u\|_{p(x)}^{q^+}.$$

We note that $2p^- > q^+$, so J is bounded from below.

(iii) If $\rho_p(u) < 1$ and $\rho_q(u) < 1$ then,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^+} + \frac{b}{2(p^+)^2} \|u\|^{2p^+} - \frac{C_2 C_3 c_{\Omega}^{q^-}}{q^-} \|u\|_{p(x)}^{q^-}.$$

(iv) If $\rho_p(u) < 1$ and $\rho_q(u) > 1$ then,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^+} + \frac{b}{2(p^+)^2} \|u\|^{2p^+} - \frac{C_2 C_3 c_{\Omega}^{q^-}}{q^-} \|u\|_{p(x)}^{q^+}.$$

Since $2p^+ > q^-$ and $2p^+ > q^+$ in (iii) and (iv) successively then J is bounded from below.

Step 2: J satisfies the (P-S) condition.

Indeed, let (u_n) be a Palais-Smale sequence for J . Thus there exists a positive constant C such that $J(u_n) \leq C$. Arguing as above, we obtain for all $u \in W_0^{1,p(x)}(\Omega)$, the following cases:

(1) If $\rho_p(u) < 1$ and $\rho_q(u) < 1$ then,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^+} + \frac{b}{2(p^+)^2} \|u\|^{2p^+} - \frac{C_2 C_3 c_\Omega^{q^-}}{q^-} \|u\|_{p(x)}^{q^-}.$$

(2) If $\rho_p(u) < 1$ and $\rho_q(u) > 1$ then,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^+} + \frac{b}{2(p^+)^2} \|u\|^{2p^+} - \frac{C_2 C_3 c_\Omega^{q^-}}{q^-} \|u\|_{p(x)}^{q^+}.$$

(3) If $\rho_p(u) > 1$ and $\rho_q(u) < 1$ then,

$$J(u) \geq \frac{a}{p^+} \|u\|^{p^-} + \frac{b}{2(p^+)^2} \|u\|^{2p^-} - \frac{C_2 C_3 c_\Omega^{q^-}}{q^-} \|u\|_{p(x)}^{q^-}.$$

(4) If $\rho_p(u) > 1$ and $\rho_q(u) > 1$ then,

$$J(u) \geq \frac{a}{p^-} \|u\|^{p^-} + \frac{b}{2(p^-)^2} \|u\|^{2p^-} - \frac{C_2 C_3 c_\Omega^{q^-}}{q^-} \|u\|_{p(x)}^{q^+}.$$

Since $q^+ < p^-$, in all cases we now deduce that the sequence (u_n) is bounded in $W_0^{1,p(x)}(\Omega)$. Thus, passing to a subsequence if necessary, there exists $u \in W_0^{1,p(x)}(\Omega)$ such that $u_n \rightharpoonup u$ weakly in $W_0^{1,p(x)}(\Omega)$. Since $a, b > 0$ we get

$$a + b \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx > a > 0.$$

Consider the sequence

$$K_n = J'(u_n)u_n + \int_\Omega f(x, u_n)u_n dx - J'(u_n)u - \int_\Omega f(x, u_n)u dx.$$

From the Lebesgue dominated convergence theorem and the Sobolev embedding, we have that

$$\int_\Omega f(x, u_n)u_n dx \rightarrow \int_\Omega f(x, u)u dx, \quad \int_\Omega f(x, u_n)u dx \rightarrow \int_\Omega f(x, u)u dx,$$

so we have that $K_n \rightarrow 0$ and it can be seen that

$$\begin{aligned} K_n &= \left(a + b \int_\Omega \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx \right) \int_\Omega |\nabla u_n|^{p(x)} \\ &\quad - \left(a + b \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx \right) \int_\Omega |\nabla u_n|^{p(x)-2} \nabla u_n \nabla u dx. \end{aligned}$$

Let

$$L_n = - \left(a + b \int_\Omega \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx \right) \int_\Omega |\nabla u|^{p(x)-2} \nabla u_n \nabla u dx$$

$$\begin{aligned}
 &+ \left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx \right) \int_{\Omega} |\nabla u|^{p(x)} dx \\
 &= - \left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx \right) \left[\int_{\Omega} |\nabla u|^{p(x)-2} \nabla u_n \nabla u dx - \int_{\Omega} |\nabla u|^{p(x)} dx \right].
 \end{aligned}$$

From the weak convergence of (u_n) , we have that $L_n \rightarrow 0$. Hence,

$$\begin{aligned}
 K_n + L_n &= \left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx \right) \left[\int_{\Omega} |\nabla u_n|^{p(x)} - \int_{\Omega} |\nabla u_n|^{p(x)-2} \nabla u_n \nabla u dx \right. \\
 &\quad \left. - \int_{\Omega} |\nabla u|^{p(x)-2} \nabla u_n \nabla u dx + \int_{\Omega} |\nabla u|^{p(x)} dx \right] \\
 &= \left(a + b \int_{\Omega} \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx \right) \\
 &\quad \times \int_{\Omega} (|\nabla u_n|^{p(x)-2} \nabla u_n - |\nabla u|^{p(x)-2} \nabla u) (\nabla u_n - \nabla u) dx.
 \end{aligned}$$

We can generalize the elementary inequalities from [14] to the variable exponent case and we obtain

$$(|x|^{p(\cdot)-2} x - |y|^{p(\cdot)-2} y)(x - y) \geq C_{p(\cdot)} |x - y|^{p(\cdot)} \quad \text{if } p(\cdot) \geq 2,$$

where $C_{p(\cdot)} \geq \min \left\{ 1, \frac{1}{2^{p^+-2}} \right\}$, and

$$(|x|^{p(\cdot)-2} x - |y|^{p(\cdot)-2} y)(x - y) \geq \frac{C_{p(\cdot)} |x - y|^2}{(|x| + |y|)^{2-p(\cdot)}} \quad \text{if } 1 < p(\cdot) < 2.$$

We obtain that,

$$K_n + L_n \geq a C_{p(\cdot)} \rho_{p(\cdot)}(\nabla(u_n - u)),$$

and as $C_{p(\cdot)}$ is dominated by a constant we deduce that $\rho_p(\nabla(u_n - u))$ converges to 0 as $n \rightarrow +\infty$. By Proposition 2.3, we conclude that $\|u_n - u\| \rightarrow 0$ in $W_0^{1,p(x)}(\Omega)$.

Proof of Theorem 3.2. We notice that $W_0^{1,p^+}(\Omega) \subset W_0^{1,p(x)}(\Omega)$. Consider $\{e_1, e_2, \dots\}$, a Schauder basis of the space $W_0^{1,p^+}(\Omega)$ (see [17]), and for each $k \in \mathbb{N}$, consider X_k , the subspace of $W_0^{1,p^+}(\Omega)$ generated by k vectors $\{e_1, e_2, \dots, e_k\}$. Clearly X_k is subspace of $W_0^{1,p(x)}(\Omega)$. So we notice that $X_k \subset L^{q(x)}(\Omega)$ because $X_k \subset W_0^{1,p^+}(\Omega) \subset L^{q(x)}$. Thus, the norms $\|\cdot\|$ and $|\cdot|_{q(x)}$ are equivalent on X_k because X_k is a finite dimension space. Consequently, there exists a positive constant C_k such that

$$-|u|_{q(x)} \leq -C_k \|u\|, \quad \text{for all } u \in X_k.$$

Thus we have

$$J(u) \leq \frac{a}{p^-} \rho_p(\nabla u) + \frac{b}{2(p^-)^2} (\rho_p(\nabla u))^2 - \frac{C_1}{q^+} \rho_q(u).$$

- If $\rho_p(\nabla u) < 1$ and $\rho_q(u) < 1$, then

$$\begin{aligned} J(u) &\leq \frac{a}{p^-} \|u\|^{p^-} + \frac{b}{2(p^-)^2} \|u\|^{2p^-} - \frac{C_1}{q^+} |u|_{q(x)}^{q^+} \\ &\leq \frac{a}{p^-} \|u\|^{p^-} + \frac{b}{2(p^-)^2} \|u\|^{2p^-} - \frac{C_1}{q^+} C_k \|u\|^{q^+} \\ &= \|u\|^{q^+} \left[\frac{a}{p^-} \|u\|^{p^- - q^+} + \frac{b}{2(p^-)^2} \|u\|^{2p^- - q^+} - \frac{C_1}{q^+} C_k \right]. \end{aligned}$$

We choose $R > 0$ small enough such that

$$\frac{a}{p^-} R^{p^- - q^+} + \frac{b}{2(p^-)^2} R^{2p^- - q^+} < \frac{C_1}{q^+} C_k.$$

Thus, for $0 < r < R$, we consider the set $K = \{u \in X_k : \|u\| = r\}$. For all $u \in K$, we have

$$\begin{aligned} J(u) &\leq r^{q^+} \left[\frac{a}{p^-} r^{p^- - q^+} + \frac{b}{2(p^-)^2} r^{2p^- - q^+} - \frac{C_1}{q^+} C_k \right] \\ &< R^{q^+} \left[\frac{a}{p^-} R^{p^- - q^+} + \frac{b}{2(p^-)^2} R^{2p^- - q^+} - \frac{C_1}{q^+} C_k \right] \\ &< 0 = J(0). \end{aligned}$$

We can apply similar reasoning to the other cases since:

- If $\rho_p(\nabla u) < 1$ and $\rho_q(u) > 1$, then

$$J(u) \leq \|u\|^{q^-} \left[\frac{a}{p^-} \|u\|^{p^- - q^-} + \frac{b}{2(p^-)^2} \|u\|^{2p^- - q^-} - \frac{C_1}{q^+} C_k \right].$$

- If $\rho_p(\nabla u) > 1$ and $\rho_q(u) < 1$, then

$$J(u) \leq \|u\|^{q^+} \left[\frac{a}{p^-} \|u\|^{p^+ - q^+} + \frac{b}{2(p^-)^2} \|u\|^{2p^+ - q^+} - \frac{C_1}{q^+} C_k \right].$$

- If $\rho_p(\nabla u) > 1$ and $\rho_q(u) > 1$, then

$$J(u) \leq \|u\|^{q^-} \left[\frac{a}{p^-} \|u\|^{p^+ - q^-} + \frac{b}{2(p^-)^2} \|u\|^{2p^+ - q^-} - \frac{C_1}{q^+} C_k \right].$$

We can be considered the odd homeomorphism $h : K \rightarrow S^{k-1}$ defined by $h(u) = (\alpha_1, \alpha_2, \dots, \alpha_k)$, where S^{k-1} is the sphere in \mathbb{R}^k . From Theorem 2.1 and proposition 2.5 we conclude that $\gamma(K) = k$. thanks to theorem 2.3, J has at least k pairs of different critical points. Since k is arbitrary, we obtain infinitely many critical points of J . \square

Example 3.2. We consider $\Omega = (0, 1)$ and let $f(x, t) = C_1 t^{q(x)-1}$ if $t \geq 0$ and $f(x, t) = -C_1 (-t)^{q(x)-1}$ if $t < 0$, where $p(x) = x^2 + 2$ and $q(x) = \frac{1}{x^2+2} + 1$, for all $x \in \Omega$. Then, the problem

$$\begin{cases} -\left(a + b \int_{\Omega} \frac{1}{x^2+2} |u'|^{x^2+2} dx\right) (|u'|^{x^2} u')' = \text{sgn}(u) C_1 |u|^{\frac{1}{x^2+2}}, & \text{in } (0, 1) \\ u(0) = u(1) = 0, \end{cases}$$

has infinitely many solutions.

REFERENCES

- [1] Ambrosetti, A., Malchiodi, A., *Nonlinear analysis and semilinear elliptic problems*, Cambridge Stud. Adv. Math., vol. 14, Cambridge Univ. Press, 2007.
- [2] Antontsev, S.N., Rodrigues, J.F., *A model porous medium equation with variable exponent of nonlinearity: existence, uniqueness and localization properties of solutions*, Nonlinear Anal. (2005), 515–545.
- [3] Antontsev, S.N., Rodrigues, J.F., *On stationary thermo-rheological viscous flows*, Ann. Univ. Ferrara Sez. VII (N.S.) **52** (2006), 19–36.
- [4] Castro, A., *Metodos variacionales y analisis funcional no linear*, X Colóquio Colombiano de Matematicas, 1980.
- [5] Clarke, D.C., *A variant of the Lusternik-Schnirelman theory*, Indiana Univ. Math. J. **22** (1972), 65–74.
- [6] Corrêa, F.J.S.A., Figueiredo, G.M., *On an elliptic equation of p -Kirchhoff-type via variational methods*, Bull. Austral. Math. Soc. **74** (2006), 263–277.
- [7] Corrêa, F.J.S.A., Figueiredo, G.M., *On a p -Kirchhoff equation via Krasnoselskii's genus*, Appl. Math. Lett. **22** ('2009), 819–822.
- [8] Dai, G., Wei, J., *Infinitely many non-negative solutions for a $p(x)$ -Kirchhoff-type problem with Dirichlet boundary condition*, Nonlinear Anal. **73** (2010), 3420–3430.
- [9] Diening, L., Harjulehto, P., Hästö, P., Ružička, M., *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Math., vol. 2017, Springer, New York, 2011.
- [10] Fan, X.L., Zhao, D., *On the spaces $L^{p(x)}$ and $W^{m,p(x)}$* , J. Math. Anal. Appl. **263** (2001), 424–446.
- [11] Kavian, O., *Introduction 'a la théorie des points critiques et applications aux problèmes elliptiques*, Springer-Verlag, 1993.
- [12] Kirchhoff, G., *Mechanik*, Teubner, Leipzig, Germany, 1883.
- [13] Krasnoselskii, M.A., *Topological methods in the theory of nonlinear integral equations*, MacMillan, New York, 1964.
- [14] Peral, I., *Multiplicity of solutions for the p -Laplacian*, Second School of Nonlinear Functional Analysis and Applications to Differential Equations, ICTP, Trieste, 1997.
- [15] Rabinowitz, P. H., *Minimax methods in critical point theory with applications to differential equations*, Conference Board of the Mathematical Sciences, by the American Mathematical Society, Providence, Rhode Island, 1984.
- [16] Ružička, M., *Electro-rheological Fluids: Modeling and Mathematical Theory*, Springer-Verlag, Berlin, 2000.
- [17] Triebel, H., *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland, Amsterdam, 1978.
- [18] Zhikov, V.V., *Averaging of functionals of the calculus of variations and elasticity theory*, Math. USSR Izv. **9** (1987), 33–66.

A. MOKHTARI, T. MOUSSAOUI,
LABORATORY OF FIXED POINT THEORY AND APPLICATIONS,
DEPARTMENT OF MATHEMATICS,
E.N.S. KOUBA, ALGIERS, ALGERIA
E-mail: mokhtarimaths@yahoo.fr moussaoui@ens-kouba.dz

NATIONAL UNIVERSITY OF IRELAND,
SCHOOL OF MATHEMATICS, STATISTICS AND APPLIED MATHEMATICS,
GALWAY, IRELAND

KING ABDULAZIZ UNIVERSITY,
NAAM RESEARCH GROUP,
JEDDAH, SAUDI ARABIA
E-mail: donal.oregan@nuigalway.ie