

Josef Voldřich

The ranges of nonlinear operators of the polynomial type

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 23 (1982), No. 4, 671--684

Persistent URL: <http://dml.cz/dmlcz/106186>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1982

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

THE RANGES OF NONLINEAR OPERATORS  
OF THE POLYNOMIAL TYPE  
Josef VOLDŘICH

**Abstract:** In this paper we prove the existence results for the equation  $Au + Su = f$ , where  $A$  is a polynomial operator on a reflexive Banach space,  $S$  is a strongly continuous nonlinearity.

**Key words:** Polynomial operators, perturbations, strong subasymptote.

Classification: 47H15

---

1. Introduction. J. Frehse investigated a class of nonlinear functional equations and nonlinear operators of polynomial type (see e.g. [1]). The ranges of these operators are closed linear subspaces with a finite codimension and the equation

$$(1.1) \quad Au = f$$

has at least one solution if  $f$  satisfies the Fredholm condition. Further, J. Frehse deals with the solvability of the equation

$$(1.2) \quad Au + Su = f,$$

where  $S$  is the Landesman-Lazer type nonlinearity (see e.g. [2]).

This paper continues, in some sense, the works [1],[2] and deals with the solvability of the equation (1.2) in section 2, where  $S$  is "subpolynomial-type" nonlinearity. In section 3 the abstract theorems are applied to the examples of polynomial

operators, for example, to the problem

$$\left\{ \begin{array}{l} (\Delta - \lambda) [(\Delta u - \lambda u)^5 + (\Delta u - \lambda u)^3] + \\ \quad + |u|^{\sigma} \operatorname{sign} u = f \text{ in } \Omega, \\ u = 0 \text{ on } \partial\Omega. \end{array} \right.$$

There are also presented results concerning the solvability of (1.2) in section 4, where the operator  $S$  has a vanishing strong subasymptote. For example, there is considered the problem

$$\left\{ \begin{array}{l} (\Delta - \lambda) [(\Delta u - \lambda u)^5 + (\Delta u - \lambda u)^3] + \frac{u}{1 + u^2} = f \text{ in } \Omega, \\ u = 0 \text{ on } \partial\Omega. \end{array} \right.$$

The proof which is published in [5], is analogous to that contained in the papers [3],[4] where equations with linear non-invertible operators in the main part are considered.

**2. Abstract theorems.** We shall investigate continuous maps  $A: B \rightarrow B^*$  where  $B$  is a real reflexive Banach space with a norm  $\|\cdot\|$ ,  $B^*$  is its dual space. We consider following conditions:

(2.1) There exists  $a \geq 0$  such that it holds

(i) if  $\limsup_{t \rightarrow +\infty} t^{-a} |\langle A(u+tv), v \rangle| < +\infty$

then  $\langle A(u+tv), v \rangle = \langle Au, v \rangle$  whenever  $t \in \mathbb{R}$ ,  $u, v \in B$ ,

(ii) if  $\limsup_{t \rightarrow +\infty} t^{-a} |\langle A(tw), v \rangle| < +\infty$

then  $\langle A(tw), v \rangle = \langle A(0), v \rangle$  whenever  $t \in \mathbb{R}$ ,  $v, w \in B$ .

(2.2) If  $u, v \in B$ ,  $\varphi(t) = \langle A(u+tv), u+tv \rangle$  and

(i)  $\liminf_{t \rightarrow +\infty} t^{-1} \varphi(t) \geq 0$ ,

(ii)  $\limsup_{t \rightarrow +\infty} t^{-1} \varphi(t) < +\infty$ ,

then  $\lim_{t \rightarrow +\infty} t^{-1} \varphi(t) = 0$ .

Any continuous operator  $A$  satisfying conditions (2.1) and (2.2) will be said  $a$ -polynomial.

An operator  $A$  satisfying

$$(2.3) \liminf_{\|u\| \rightarrow \infty} \|u-v\|^{-1} \langle Au-Av, u-v \rangle \geq 0 \text{ for each } v \in B$$

will be called the asymptotically monotone operator.

(2.4) There exist constants  $K, c > 0$ ,  $p > 1$  and a finite dimensional subspace  $V \subset B$  with a bounded linear projection  $Q: B \rightarrow V$  such that

$$\langle Au, u \rangle \geq c \|u\|^p - K \|Qu\|^p - K \text{ whenever } u \in B.$$

2.5. Definition. A continuous operator  $A: B \rightarrow B^*$  is said regular if the variational inequality

$$\langle Au-f, u-v \rangle \leq 0, \quad v \in K,$$

has a solution  $u \in K$  for any bounded closed convex set  $K \subset B$  and for every  $f \in B^*$ .

The main result of Frehse's work [1] is as follows.

2.6. Theorem. Let  $A: B \rightarrow B^*$  be a regular operator satisfying conditions (2.1)(i) with  $a = 0$ , (2.2)-(2.4) and let  $A(0) = 0$ . Then the equation  $Au = f$  has at least one solution if and only if  $f \perp (R(A))^\perp$ .

Moreover,  $\dim R(A)^\perp \leq \dim V < +\infty$ .

We shall use the next lemma in proofs of the following theorems.

2.7. Lemma. Let  $A: B \rightarrow B^*$  be an asymptotically monotone  $a$ -polynomial operator,  $A(0) = 0$ . Suppose that for some  $v \in B$  there exist constants  $\sigma, C, K \geq 0$  such that the inequality

$$(2.8) \quad \langle Aw, v \rangle \leq C + K \|w\|^\sigma$$

holds for every  $w \in B$ . If  $a \geq \sigma$  then  $v \perp R(A)$ .

Proof. The inequality (2.8) implies  $\langle A(w+tv), v \rangle \leq C + K \|w+tv\|^\sigma$  and from the asymptotical monotonicity of the

operator  $A$  (i.e.  $\liminf_{|t| \rightarrow \infty} |t|^{-1} \langle A(w+tv) - Aw, tv \rangle \geq 0$ ) we obtain  $\langle A(w+tv), v \rangle \geq \langle Aw, v \rangle - \varepsilon$  for every  $t \geq t_0$  with some  $t_0 > 0$ ,  $\varepsilon > 0$ . Together with the supposition (2.8) and the condition (2.1) we have

$$(2.9) \quad \langle A(w+tv), v \rangle = \langle Aw, v \rangle \text{ for every } t \in \mathbb{R}.$$

Using the inequality  $\liminf_{|t| \rightarrow +\infty} |t|^{-1} \langle A(w+tv) - A(2w), -w+tv \rangle \geq 0$  we get that  $\limsup_{|t| \rightarrow +\infty} |t|^{-1} \langle A(w+tv), w \rangle \leq K(w)$  with some constant  $K(w)$ . It yields together with (2.9)  $\limsup_{|t| \rightarrow +\infty} |t|^{-1} \varphi(t) < +\infty$ , where  $\varphi(t) = \langle A(w+tv), w+tv \rangle$ . From conditions (2.2), (2.3),  $A(0) = 0$  it follows that

$$(2.10) \quad \lim_{|t| \rightarrow +\infty} t^{-1} \varphi(t) = 0.$$

Let  $s \in \mathbb{R}$  be fixed. It is obvious that

$$\liminf_{|t| \rightarrow +\infty} |t|^{-1} \langle A(w+tv) - A(sw), (1-s)w+tv \rangle \geq 0$$

and this together with (2.9) yields

$$\liminf_{|t| \rightarrow +\infty} |t|^{-1} [(1-s)\varphi(t) + s \langle Aw, tv \rangle - \langle A(sw), (1-s)w+tv \rangle] \geq 0.$$

According to this fact and with respect to the condition (2.10) we have  $s \langle Aw, v \rangle - \langle A(sw), v \rangle \geq 0$ ,  $-s \langle Aw, v \rangle + \langle A(sw), v \rangle \geq 0$  and

$$(2.11) \quad s \langle Aw, v \rangle = \langle A(sw), v \rangle, \quad s \in \mathbb{R}.$$

If  $a < 1$  then  $0 \leq \sigma < 1$  and as  $s \langle Aw, v \rangle \leq C + K|s|^\sigma \|w\|^\sigma$  we get  $\langle Aw, v \rangle = 0$ , taking the limits  $s \rightarrow \pm\infty$ . This completes the proof for  $a < 1$ .

Let  $a \geq 1$ . There exists  $\gamma > 0$  such that  $\|Au\| \leq 1 + \|A(0)\| = 1$  for every  $u \in B$ ,  $\|u\| \leq \gamma$ . The inequality

$$\langle Aw, v \rangle = \frac{\|w\|}{\gamma} \langle A(\frac{w}{\|w\|} \gamma), v \rangle \geq -\frac{\|w\|}{\gamma} \|v\|, \quad w \neq 0,$$

is an immediate consequence of (2.11). Therefore, there exists the constant  $L > 0$  such that  $\langle Aw, v \rangle \geq -L\|w\|$ ,  $w \in B$ . Using the inequality (2.8) and the fact that  $a \geq 1$  we obtain

$$\limsup_{t \rightarrow +\infty} t^{-a} |\langle A(tw), v \rangle| < +\infty.$$

From (2.1) we get  $\langle Aw, v \rangle = \langle A(0), v \rangle = 0$ . It means that  $v \perp R(A)$  and the proof of the lemma is complete.

Let  $S: B \rightarrow B^*$  be an operator satisfying conditions

$$(2.12) \quad \|Su\|_{B^*} \leq \alpha + \beta \|u\|^{\sigma}, \quad \alpha, \beta, \sigma \geq 0,$$

(2.13) there exist constants  $G, H > 0$  such that the inequality

$$\liminf_{\|u_j\| \rightarrow +\infty} \|u_j\|^{-1} \langle Su_j - Sw, u_j - w \rangle \geq -G - H \|w\|^{\sigma}$$

is fulfilled for every  $w \in B$ .

**2.14. Definition.** Let  $V$  be a closed linear subspace of  $B$ ,  $V_r = \{u \in V, \|u\| \leq r\}$ . A mapping  $\Psi: V_1 \rightarrow R$  will be said a strong subasymptote of the operator  $S$  with respect to  $V$  if

$$(2.15) \quad \Psi(z) \leq \liminf_{j \rightarrow +\infty} \langle Su_j, \|u_j\|^{-1}(u_j - w) \rangle, \quad w \in B,$$

holds for any sequence  $\{u_j\}_{j=1}^{+\infty}$  such that  $\|u_j\| \rightarrow +\infty$  and  $\|u_j\|^{-1}u_j \rightarrow z$  (i.e. weakly) for  $j \rightarrow +\infty$ , where  $z \neq 0, z \in V$ .

**2.16. Theorem.** Let  $A, S: B \rightarrow B^*$  be continuous operators with the following properties

- (i)  $A$  is an asymptotically monotone  $a$ -polynomial operator,  $A(0) = 0$  and  $A$  satisfies (2.4),
- (ii)  $S$  satisfies (2.12), (2.13) and  $p > 1 + \sigma$ ,  $a \geq \sigma$ ,
- (iii)  $A + S$  is a regular operator.

If  $\Psi: (R(A)^\perp)_1 \rightarrow R$  is a strong subasymptote of the operator  $S$  with respect to  $(R(A)^\perp)_1$  and if

$$(2.17) \quad \langle f, z \rangle < \Psi(z) \text{ for every } z \in (R(A)^\perp)_1, z \neq 0,$$

then the equation (1.2) has at least one solution.

**Proof.** Let us suppose that the equation is not solvable and let  $u_r$  be the solution of the variational inequality

$$(2.18) \quad \langle Au + Su - f, u - w \rangle \leq 0, \quad w \in B_r.$$

Observe that  $u_r \in \partial B_r$  and therefore  $\|u_r\| = r$ . Choose a sequence  $\{r_i\}_{i=1}^{+\infty}$  so that  $\|u_{r_i}\|^{-1} u_{r_i} \rightharpoonup z$  weakly in  $B$ . According to (2.18) with  $w = 0$  and in view of the growth of  $S$  (see (2.12)) we get the inequality  $\langle Au_{r_i}, u_{r_i} \rangle \leq L \|u_{r_i}\|^{1+\sigma}$  for  $i \geq i_0$  with some positive constant  $L$ . Since  $p > 1 + \sigma$  we obtain from (2.4) that  $\liminf_{i \rightarrow +\infty} \|Qu_{r_i}\|^p \|u_{r_i}\|^{-p} \geq \frac{c}{K} > 0$ . The fact that  $\dim R(Q) < +\infty$  implies  $Q(u_{r_i} \|u_{r_i}\|^{-1}) \rightarrow Qz$  in  $B$  for  $i \rightarrow +\infty$  and  $\|Qz\| > 0$ , therefore  $z \neq 0$ .

We claim  $z \perp R(A)$ . Observe that

$$\liminf_{i \rightarrow +\infty} \|u_{r_i}\|^{-1} \langle Au_{r_i} - Aw, u_{r_i} - w \rangle \geq 0,$$

$$\liminf_{i \rightarrow +\infty} \|u_{r_i}\|^{-1} \langle f - Au_{r_i} - Su_{r_i}, u_{r_i} - w \rangle \geq 0$$

and therefore

$$(2.19) \quad \liminf_{i \rightarrow +\infty} \|u_{r_i}\|^{-1} \langle f - Su_{r_i} - Aw, u_{r_i} - w \rangle \geq 0.$$

From (2.13) we have

$$\liminf_{i \rightarrow +\infty} \|u_{r_i}\|^{-1} \langle f - Aw - Sw, u_{r_i} - w \rangle \geq -G - H \|w\|^{\sigma}$$

and this gives the estimate

$$\liminf_{i \rightarrow +\infty} \langle -Aw, u_{r_i} \|u_{r_i}\|^{-1} \rangle \geq -G - H \|w\|^{\sigma} - (\alpha + \beta \|w\|^{\sigma}) - |\langle f, z \rangle|.$$

Consequently,  $\langle Aw, z \rangle \leq G + |\langle f, z \rangle| + \alpha + (\beta + H) \|w\|^{\sigma}$  and the Lemma 2.7 implies  $z \perp R(A)$ .

Observe that the inequality (2.19) yields

$$\langle f, z \rangle - \langle Aw, z \rangle - \liminf_{i \rightarrow +\infty} \langle Su_{r_i}, \|u_{r_i}\|^{-1} (u_{r_i} - w) \rangle \geq 0.$$

As  $\Psi$  is the strong subasymptote of the operator  $S$  we get  $\langle f, z \rangle - \Psi(z) \geq 0$ , which is the contradiction with (2.17) and the proof is complete.

2.20. Proposition. The condition (2.17) is necessary for the solvability of (1.2), if  $\langle Su, z \rangle < \Psi(z)$  for every  $u \in B$ ,  $z \neq 0$ ,  $z \in (R(A)^\perp)_1$ .

Proof. If  $Au + Su = f$  then  $\langle f, z \rangle = \langle Su, z \rangle < \Psi(z)$  for  $z \in (R(A)^\perp)_1$ .

In the case  $\delta < 1$ , the strong subasymptote of the operator  $S$  can be replaced by more verifiable conditions:

$$(2.21) \quad \liminf_{\|u_i\| \rightarrow +\infty} \|u_i\|^{-1} \langle Su_i - Sw_i, u_i - w_i \rangle \geq -G$$

for every bounded sequence  $\{w_i\}_{i=1}^{+\infty}$ .

(2.22) For every  $z \in (R(A)^\perp)$ ,  $z \neq 0$ , there exist  $t_z \in R$ ,  $v_z \in B$  such that  $\langle S(t_z z + v_z), z \rangle > G$ , where  $G$  is the constant from (2.21).

$$(2.23) \quad \liminf_{t \rightarrow +\infty} \langle S(tz_i + v), -z_i \rangle \leq \langle S(tz + v), -z \rangle$$

holds for any  $t \in R$ ,  $v \in B$  and any sequence  $\{z_i\}_{i=1}^{+\infty} \subset B$ ,  $z_i \rightharpoonup z$  weakly for  $i \rightarrow +\infty$ ,  $z \in (R(A)^\perp)$ ,  $z \neq 0$ .

A strongly continuous operator  $S$  satisfies the condition (2.23).

2.24. Theorem. Let  $A, S: B \rightarrow B^*$  be continuous operators with the following properties

(i)  $A$  is an asymptotically monotone  $a$ -polynomial operator satisfying (2.4),  $A(0) = 0$ ,

(ii)  $S$  satisfies (2.12), (2.21)-(2.23) and  $p > 1 + \delta$ ,  $a \geq \delta$ ,  $\delta < 1$ ,

(iii)  $A + S$  is a regular operator.

Then the equation  $Au + Su = 0$  has at least one solution.

Proof. The condition (2.21) implies (2.13). Let us suppose that the equation  $Au + Su = 0$  is not solvable. Analogously as in the proof of Theorem 2.16 there exists a sequence



$\{u_{r_i}\}_{i=1}^{+\infty}$ ,  $\|u_{r_i}\| \rightarrow +\infty$ ,  $\|u_{r_i}\|^{-1} u_{r_i} \rightarrow z$  weakly in  $B$  for  $i \rightarrow +\infty$ ,  $z \in R(A)^{\perp}$ ,  $z \neq 0$ , and  $\langle Au_{r_i} + Su_{r_i}, u_{r_i} - w \rangle \leq 0$  for every  $w \in B_{r_i}$ . As the operator  $S$  satisfies (2.21) and (2.22) we have

$$\begin{aligned}
 -G &\leq \liminf_{i \rightarrow +\infty} \|u_{r_i}\|^{-1} \langle S(t_z u_{r_i} \|u_{r_i}\|^{-1} + v_z) - Su_{r_i}, \\
 &\quad t_z u_{r_i} \|u_{r_i}\|^{-1} + v_z - u_{r_i} \rangle = \\
 &= \liminf_{i \rightarrow +\infty} \langle S(t_z u_{r_i} \|u_{r_i}\|^{-1} + v_z) - Su_{r_i}, -u_{r_i} \|u_{r_i}\|^{-1} \rangle
 \end{aligned}$$

because  $\sigma < 1$ . The operator  $A + S$  is regular and therefore we get  $\langle Au_{r_i} + Su_{r_i}, -u_{r_i} \rangle \geq 0$  and

$$\liminf_{i \rightarrow +\infty} \langle S(t_z u_{r_i} \|u_{r_i}\|^{-1} + v_z) + Au_{r_i}, -u_{r_i} \|u_{r_i}\|^{-1} \rangle \geq -G.$$

Further,  $A$  is asymptotically monotone, e.g.

$$\liminf_{i \rightarrow +\infty} \langle -Au_{r_i}, -u_{r_i} \|u_{r_i}\|^{-1} \rangle \geq 0$$

and

$$\liminf_{i \rightarrow +\infty} \langle S(t_z u_{r_i} \|u_{r_i}\|^{-1} + v_z), -u_{r_i} \|u_{r_i}\|^{-1} \rangle \geq -G.$$

From (2.23) we obtain  $\langle S(t_z z + v_z), z \rangle \leq G$ , which is the contradiction with (2.22).

**3. Examples.** Let  $P_j: R^s \rightarrow R$ ,  $j = 1, 2, \dots, s$ , be polynomials satisfying the following conditions (with  $C, K, c > 0$ )

$$(3.1) \quad |P_j(\xi)| \leq C(1 + |\xi|^{p-1}) \text{ for every } \xi \in R^s,$$

$$(3.2) \quad \sum_{j=1}^s P_j(\xi) \xi_j \geq c |\xi|^p - K \text{ for every } \xi \in R^s,$$

$$(3.3) \quad \sum_{j=1}^s (P_j(\xi) - P_j(\eta)) (\xi_j - \eta_j) \geq 0 \text{ for all } \xi, \eta \in R^s.$$

Let  $\Omega \subset R^N$  be a bounded domain with a smooth boundary and let  $V = W^{2m, p}(\Omega) \cap W_0^{m, p}(\Omega)$ ,  $p > 1$ . We define

$L_j u = \sum_{|\kappa|, |q| \leq m} (-1)^r D^r (a_{r,q}^{(j)}(x) D^q u)$ ,  $j = 1, \dots, s$ ,  
 for every  $u \in V$  where  $a_{r,q}^{(j)} \in C^\infty(\bar{\Omega})$  ( $|\kappa|, |q| \leq m$ ,  $j = 1, \dots, s$ ). Let

$$\sum_{|\kappa|, |q| \leq m} (-1)^m a_{r,q}^{(j)}(x) \zeta^{r+q} \geq \alpha |\zeta|^{2m}, \quad j = 1, \dots, s,$$

hold with some  $\alpha > 0$  for every  $\zeta \in \mathbb{R}^N$ . Let us define the operator  $A: V \rightarrow V^*$  by

$$\langle Au, v \rangle = \sum_{j=1}^s \int_{\Omega} P_j(L_1 u, \dots, L_s u) L_j v, \quad v \in V.$$

Using the Theorem 2.6, we see that the equation  $Au = f$  is solvable if  $(f - A(0)) \perp (R(A) - A(0))^\perp$ . Let us remark that for  $s = 1$  it is possible to show: if we consider the operator  $A: V/\text{Ker}[L_1] \rightarrow (V/\text{Ker}[L_1])^*$  then this result follows from the theory of monotone operators and  $(R(A) - A(0))^\perp = \text{Ker}[L_1]$ .

Let the function  $\varphi$  be continuous, odd, increasing,  $\lim_{t \rightarrow +\infty} \varphi(t) = +\infty$  and  $|\varphi(t)| \leq \bar{\alpha} + \bar{\beta}|t|^\sigma$ ,  $t \in \mathbb{R}$ , with some  $\bar{\alpha}, \bar{\beta}, \sigma > 0$ . Let  $2mp > N$ . We define the operator  $S: V \rightarrow V^*$  by

$$\langle Su, v \rangle = \int_{\Omega} \varphi(u) v, \quad v \in V.$$

We note that the inequality (2.12) holds with some constants  $\alpha, \beta$ . Let us assume the conditions

$$(3.4) \quad \limsup_{t \rightarrow +\infty} \varphi(\omega t) [\varphi(t)]^{-1} = \chi(\omega) < +\infty$$

for every  $\omega \geq 1$ , where  $\chi$  is a continuous function with

$$\lim_{\omega \rightarrow 1^+} \chi(\omega) = 1,$$

$$(3.5) \quad \text{meas } \Omega > 2 \text{ meas } \{x \in \Omega; z(x) = 0\}$$

for every  $z \in (R(A) - A(0))^\perp$ ,  $z \neq 0$ .

**3.6. Proposition.** The mapping  $\Psi: ((R(A) - A(0))^\perp)_1 \rightarrow \{K\}$ , where  $K$  is a real number, is a strong subasymptote of the

operator  $S$  defined above with respect to  $(R(A) - A(0))^\perp$ .

Proof. We assume that  $A(0) = 0$  and that for a sequence  $\{u_n\}_{n=1}^{+\infty} \subset V$  it is  $\|u_n\|^{-1} \rightarrow +\infty$ ,  $(u_n - w) \|u_n\|^{-1} \rightarrow z$  weakly for  $n \rightarrow +\infty$ ,  $z \in R(A)^\perp$ ,  $z \neq 0$ ,  $w \in V$ . It suffices to show that

$$\liminf_{n \rightarrow +\infty} \int_{\Omega} \varphi(u_n) \frac{u_n - w}{\|u_n\|} - K \geq 0.$$

As  $W^{2m,p}(\Omega)$  is compactly imbedded into  $C(\bar{\Omega})$  we have  $u_n \|u_n\|^{-1} \rightarrow z$  and  $(u_n - w) \|u_n\|^{-1} \rightarrow z$  in  $L_\infty(\Omega)$ . If we denote  $\Omega_\varepsilon^+ = \{x \in \Omega; z(x) \geq \varepsilon\}$ ,  $\Omega_\varepsilon^- = \{x \in \Omega; z(x) \leq -\varepsilon\}$ ,  $\Omega_\varepsilon = \Omega_\varepsilon^- \cup \Omega_\varepsilon^+$  then according to (3.4) and (3.5) there exist  $\varepsilon > 0$ , an integer  $k_0 > 1$  such that the inequality

$$(3.6) \quad \text{meas } \Omega_\varepsilon - \frac{k+1}{k-1} \chi \left( \frac{k+1}{k-1} \right) \text{meas}(\Omega \setminus \Omega_\varepsilon) > 0$$

holds for every  $k \geq k_0$ . There exists a natural number  $n_0$  such that

$$z(x) - \frac{\varepsilon}{k_0} \leq \frac{u_n(x) - w(x)}{\|u_n\|} \leq \frac{\varepsilon}{k_0} + z(x) \quad \text{a.e. in } \Omega,$$

$$z(x) - \frac{\varepsilon}{k_0} \leq \frac{u_n(x)}{\|u_n\|} \leq \frac{\varepsilon}{k_0} + z(x) \quad \text{a.e. in } \Omega$$

for every  $n \geq n_0$ . So we get

$$\begin{aligned} \int_{\Omega} \varphi(u_n) \frac{u_n - w}{\|u_n\|} &\geq \int_{\Omega_\varepsilon^+} \varphi(u_n) \frac{u_n - w}{\|u_n\|} + \int_{\Omega_\varepsilon^-} \varphi(-u_n) \frac{-u_n + w}{\|u_n\|} - \\ &- \int_{\Omega \setminus \Omega_\varepsilon} \varepsilon \frac{k_0 + 1}{k_0} \varphi\left(\varepsilon \frac{k_0 + 1}{k_0} \|u_n\|\right) \geq \\ &\geq \int_{\Omega_\varepsilon} \varepsilon \frac{k_0 - 1}{k_0} \varphi\left(\varepsilon \frac{k_0 - 1}{k_0} \|u_n\|\right) - \int_{\Omega \setminus \Omega_\varepsilon} \varepsilon \frac{k_0 + 1}{k_0} \varphi\left(\varepsilon \frac{k_0 + 1}{k_0} \|u_n\|\right) \geq \\ &\geq \varepsilon \frac{k_0 - 1}{k_0} \varphi\left(\varepsilon \frac{k_0 - 1}{k_0} \|u_n\|\right) \text{meas } \Omega_\varepsilon - \end{aligned}$$

$$- \frac{k_0 + 1}{k_0} \varepsilon \left[ \chi \left( \frac{k_0 + 1}{k_0 - 1} \right) + \vartheta_n \right] \text{meas}(\Omega \setminus \Omega_\varepsilon) \varphi \left( \varepsilon \frac{k_0 - 1}{k_0} \|u_n\| \right),$$

where  $\vartheta_n \rightarrow 0$  for  $n \rightarrow +\infty$ . Observe that

$$\int_{\Omega} \varphi(u_n) \frac{u_n - w}{\|u_n\|} \geq \varepsilon \frac{k_0 - 1}{k_0} \varphi \left( \varepsilon \frac{k_0 - 1}{k_0} \|u_n\| \right) [\text{meas } \Omega_\varepsilon -$$

$$- \frac{k_0 + 1}{k_0 - 1} \left( \chi \left( \frac{k_0 + 1}{k_0 - 1} \right) + \vartheta_n \right) \text{meas}(\Omega \setminus \Omega_\varepsilon)].$$

Denote the expression in the square brackets by  $c_n$ . It follows from (3.6) that  $\lim_{n \rightarrow +\infty} c_n > 0$  and therefore

$$\lim_{n \rightarrow +\infty} \varepsilon \frac{k_0 - 1}{k_0} \varphi \left( \varepsilon \frac{k_0 - 1}{k_0} \|u_n\| \right) c_n = +\infty.$$

The proof is finished.

If the operator  $A$  satisfies the condition (3.5) then the Theorem 2.16 can be applied. If  $\sigma < 1$  then the operator  $S - f$  satisfies the conditions (2.21)-(2.23) and the Theorem 2.24 can be used. In these cases, if  $p > 1 + \sigma$ , a  $\delta > 0$  then the equation  $Au + Su = f$  has at least one solution.

For example, the problem

$$(\Delta - \lambda)[(\Delta u - \lambda u)^5 + (\Delta u - \lambda u)^3] + |u|^\sigma \text{sign } u = f \text{ in } \Omega,$$

$$u = 0 \text{ on } \partial\Omega$$

has at least one weak solution  $u \in W_0^{1,6}(\Omega) \cap W^{2,6}(\Omega)$  for  $0 < \sigma < 3$ .

4. Problems with a bounded nonlinearity. Let  $B$  be a linear closed subspace of  $W^{k,p}(\Omega)$ ,  $k, p > N$ ,  $p > 1$ ,  $A(0) = 0$ ,

$$(4.1) \quad \langle Su, v \rangle = \int_{\Omega} \varphi(u) v, \text{ for } u, v \in B,$$

where the function  $\varphi$  is continuous, odd,  $\lim_{|t| \rightarrow +\infty} \varphi(t) = 0$ . Then

$\|Su\|_{B^*} \leq \varrho$  for every  $u \in B$  with some constant  $\varrho$ . Further,

we shall assume the following conditions be satisfied

(4.2) for all  $w \in R(A)^\perp$ ,  $t \in R$ ,  $v \in B$  it is  $A(v + tw) = Av$ ,

(4.3) there exists a bounded linear projection  $Q: B \rightarrow R(A)^\perp$  and  $\langle Au, u \rangle \geq C \|u\|^p - K \|Qu\|^p - L$  for every  $u \in B$ , where  $p > 1$ ,  $C, K, L > 0$ .

**4.4. Proposition.** Let the function  $t \mapsto \langle A(u + tv), w \rangle$  be a polynomial for any fixed  $u, v, w \in B$ . If  $A$  is regular and satisfies (2.3), (2.4),  $A(0) = 0$ , then the condition (4.2) is fulfilled.

The proof can be found in Frehse's papers or in [5].

Let  $\Psi: (0, +\infty) \rightarrow (0, +\infty)$  be the increasing function satisfying

$$\sup_{\substack{w \in R(A)^\perp \\ \|w\|_{C(\bar{\Omega})} = 1}} \int_{\Omega_\varepsilon(w)} |w| \leq \Psi(\varepsilon),$$

where  $\Omega_\varepsilon(w) = \{x \in \Omega; 0 < |w(x)| < \varepsilon\}$  and such that

$$\limsup_{\varepsilon \rightarrow 0_+} [\Psi(\varepsilon)]^{-1} \Psi(\omega\varepsilon) < +\infty \text{ for every } \omega \in (0, +\infty).$$

**4.5. Theorem.** Let a regular asymptotically monotone 0-polynomial operator  $A$  satisfy the conditions (4.2), (4.3),  $A(0) = 0$  and let  $S$  be given by (4.1). If

$$(4.6) \quad \lim_{\xi \rightarrow +\infty} [\Psi(\frac{1}{\xi})]^{-1} \min_{\tau \in \langle a, \xi \rangle} \varphi(\tau) = +\infty$$

for some  $a > 0$  then the equation  $Au + Su = f$  has at least one solution for an arbitrary  $f \in R(A)^\perp$ .

Sketch of the proof. Let us consider the function

$$\tilde{\varphi}: \xi \mapsto \begin{cases} \varphi(\xi) & \text{for } |\xi| \leq b, \\ \varphi(b) & \text{for } \xi > b, \\ \varphi(-b) & \text{for } \xi < -b, \end{cases}$$

and the corresponding equation  $Au + \tilde{S}u = f$ . From the Theorem 2.16 this equation has a solution  $u$  because

$$0 = \sup_{\substack{w \in R(A)^\perp \\ \|w\|_{C(\bar{\Omega})} = 1}} |\langle f, w \rangle| < |\tilde{\mathcal{Q}}(b)| \inf_{\substack{w \in R(A)^\perp \\ \|w\|_{C(\bar{\Omega})} = 1}} \int_{\Omega} |w|.$$

Using the condition (4.2) we can obtain a priori estimate

$$\|Q^c u\|_{C(\bar{\Omega})} \leq c_1 = c_1(\|f\|_{B^*}).$$

Further, methods from [3],[4] give a priori estimate

$$\|Qu\|_{C(\bar{\Omega})} \leq c_3 = c_3(a, \tilde{\mathcal{Q}}, f),$$

where  $a > 0$ ,

$$c_3 = \frac{a + c_1}{\Psi^{-1}(c_2(\inf_{\xi \geq a} \tilde{\mathcal{Q}}(\xi) + \sup_{\xi \in \mathbb{R}} |\tilde{\mathcal{Q}}(\xi)|)^{-1})},$$

$$c_2 = c_2(a, \tilde{\mathcal{Q}}, f) = \inf_{\substack{w \in R(A)^\perp \\ \|w\|_{C(\bar{\Omega})} = 1}} (\inf_{\xi \geq a} \tilde{\mathcal{Q}}(\xi) \int_{\Omega} |w|).$$

If there exist numbers  $a, b \in \mathbb{R}$ ,  $0 < a < b$ , such that  $b > c_1(\tilde{\mathcal{Q}}, f) + c_3(a, \tilde{\mathcal{Q}}, f)$  then the solution  $u$  of the equation  $Au + \tilde{S}u = f$  is also the solution of the equation  $Au + Su = f$  because  $\tilde{S}u = Su$ . The condition (4.6) guarantees the existence of such numbers  $a, b$ .

For example, the problem

$$\begin{cases} (\Delta - \lambda) [(\Delta u - \lambda u)^5 + (\Delta u - \lambda u)^3] + \frac{u}{1 + u^2} = f \text{ in } \Omega, \\ u = 0 \text{ on } \partial\Omega, \end{cases}$$

has at least one weak solution  $u \in W_0^{1,6}(\Omega) \cap W^{2,6}(\Omega)$  if  $f \perp \text{Ker}[\Delta - \lambda \text{id}]$ .

It is also possible to apply the abstract results to the existence of solution of the Neuman problem

$$\sum_{i=1}^N \frac{\partial}{\partial x_i} [(\alpha + |\nabla u|^2)^{\frac{p}{2}} - 1] \frac{\partial u}{\partial x_i} + \frac{u}{1 + |u|^k} = f \text{ in } \Omega$$

$$\frac{\partial u}{\partial n} = 0 \text{ on } \partial\Omega,$$

where  $c > 0$ ,  $p > 1$ ,  $k \geq 2$ . If  $f \in L_1(\Omega)$ ,  $\int_{\Omega} f(x) dx = 0$ , this problem has at least one weak solution  $u \in W^{1,p}(\Omega)$ .

#### R e f e r e n c e s

- [1] J. FREHSE: Solvability and alternative theorems for a class of nonlinear functional equations in Banach spaces, Ark. Math. 17(1979), no. 1, 93-105.
- [2] J. FREHSE: Landesman-Lazer alternative theorems for a class of nonlinear functional equations, Math. Ann. 238(1978), no.1, 59-65.
- [3] S. FUČÍK: Solvability of nonlinear equations and boundary value problems, Society of Czechoslovak mathematicians and physicists, Prague, 1980.
- [4] S. FUČÍK, M. KRBEČ: Boundary value problems with bounded nonlinearity and general null-space of the linear part, Math. Z. 155(1977), 129-138.
- [5] J. VOLDŘICH: Nonlinear noncoercive operator equations (in Czech), Graduate theses, Charles University, Prague, 1980.

Katedra matematiky VŠSE, Nejedlého sady 14, 30614 Plzeň, Czechoslovakia

(Oblatum 9. 4. 1982)