

Lech Górniewicz

Periodic problems for ODEs via multivalued Poincaré operators

Archivum Mathematicum, Vol. 34 (1998), No. 1, 93--104

Persistent URL: <http://dml.cz/dmlcz/107636>

Terms of use:

© Masaryk University, 1998

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Periodic Problems for ODE's via Multivalued Poincaré Operators

Lech Górniewicz

Faculty of Mathematics and Informatics
Nicholas Copernicus University
Chopina 12/18, 87-100 Toruń, Poland
Email: gorn@mat.uni.torun.pl

Abstract. We shall consider periodic problems for ordinary differential equations of the form

$$\begin{cases} x'(t) = f(t, x(t)), \\ x(0) = x(a), \end{cases} \quad (\text{I})$$

where $f : [0, a] \times R^n \rightarrow R^n$ satisfies suitable assumptions.

To study the above problem we shall follow an approach based on the topological degree theory. Roughly speaking, if on some ball of R^n , the topological degree of, associated to (I), multivalued Poincaré operator P turns out to be different from zero, then problem (I) has solutions.

Next by using the multivalued version of the classical Liapunov-Krasnoselskiĭ guiding potential method we calculate the topological degree of the Poincaré operator P . To do it we associate with f a guiding potential V which is assumed to be locally Lipschitzean (instead of C^1) and hence, by using Clarke generalized gradient calculus we are able to prove existence results for (I), of the classical type, obtained earlier under the assumption that V is C^1 .

Note that using a technique of the same type (adopting to the random case) we are able to obtain all of above mentioned results for the following random periodic problem:

$$\begin{cases} x'(\xi, t) = f(\xi, t, x(\xi, t)), \\ x(\xi, 0) = x(\xi, a), \end{cases} \quad (\text{II})$$

where $f : \Omega \times [0, a] \times R^n \rightarrow R^n$ is a random operator satisfying suitable assumptions.

This paper stands a simplification of earlier works of F. S. De Blasi, G. Pianigiani and L. Górniewicz (see: [7], [8]), where the case of differential inclusions is considered.

AMS Subject Classification. 34C25, 55M20, 47H10

Keywords. Periodic processes, topological degree, Poincaré translation operator

1 Background in geometric topology

Throughout this note \mathbb{R}^n , $n \geq 1$, is an n -dimensional real Euclidean space, with inner product $\langle \cdot, \cdot \rangle$ and induced norm $\| \cdot \|$. A closed (resp. open) ball in \mathbb{R}^n with center x and radius $r \geq 0$ is denoted by $B^n(x, r)$ (resp. $B_0^n(x, r)$). Furthermore we put:

$$\begin{aligned} B^n(r) &= B^n(0, r), & B_0^n(r) &= B_0^n(0, r), \\ S^{n-1}(r) &= B^n(r) \setminus B_0^n(r), & \mathbb{P}^n &= \mathbb{R}^n \setminus \{0\}. \end{aligned}$$

\mathbb{Z} stands for the set of all integers.

For $A \subset \mathbb{R}^n$ we denote by \overline{A} , the closure of A . If $A \subset \mathbb{R}^n$ is nonempty, we put

$$|A|^- = \inf\{\|a\| \mid a \in A\}.$$

As usual, $\varphi : X \rightarrow Y$ (resp. $\varphi : X \multimap Y$) denotes a single valued (resp. multivalued) map φ from X to Y .

In the sequel, any topological space X we consider is always supposed to be metric. When the clarity is not affected, by “space X ” we mean “topological metric space X ”.

A space X is called *contractible* if there is a continuous homotopy $h : X \times [0, 1] \rightarrow X$ and a point $x_0 \in X$ such that:

$$\begin{aligned} h(x, 0) &= x && \text{for every } x \in X, \\ h(x, 1) &= x_0 && \text{for every } x \in X. \end{aligned}$$

A nonempty compact space X is called an R_δ -set if there is a decreasing sequence $\{X_k\}$ of compact contractible spaces X_k such that

$$X = \bigcap_{k=1}^{+\infty} X_k.$$

A space X is called an *absolute neighbourhood retract* ($X \in ANR$) if, for every space Y and any closed set $C \subset Y$ and any continuous map $f : C \rightarrow X$, there is an open neighbourhood U of C in Y and a continuous map $g : U \rightarrow X$ such that:

$$g(x) = f(x) \quad \text{for every } x \in C. \quad (3)$$

A space X is called an *absolute retract* ($X \in AR$) if, for any space Y and any closed $C \subset Y$ and any continuous map $f : C \rightarrow X$, there is a continuous map $g : Y \rightarrow X$ satisfying (3).

Clearly $X \in AR$ implies $X \in ANR$ (the converse is not true). Moreover, if $X, Y \in ANR$ then $X \times Y \in ANR$. It is also easy to verify that any $X \in AR$ is contractible.

A multivalued map $\varphi : X \multimap Y$ with nonempty values is called *upper semicontinuous* (u.s.c.), if $\{x \in X \mid \varphi(x) \subset U\}$ is open in X for each open $U \subset Y$.

As usual $C([a, b], \mathbb{R}^n)$ stands for the Banach space of all continuous maps $x : [a, b] \rightarrow \mathbb{R}^n$, endowed with the norm of uniform convergence. Clearly it holds $C([a, b], \mathbb{R}^n) \in AR$.

Now, following [11], (see also [6,7,10]), we recall some definitions of the topological degree for multivalued maps. Applications to periodic problems (comp. (9)) will be given later, in Sections 3 and 4.

For any $X \in ANR$ we set

$$J(B^n(r), X) = \{F : B^n(r) \multimap X \mid F \text{ is u.s.c. with } R_\delta\text{-values}\}.$$

For any continuous $f : X \rightarrow \mathbb{R}^n$, when $X \in ANR$, we put

$$J_f(B^n(r), \mathbb{R}^n) = \{\varphi : B^n(r) \multimap \mathbb{R}^n \mid \varphi = f \circ F \text{ for some } F \in J(B^n(r), X), \\ \text{and } \varphi(S^{n-1}(r)) \subset \mathbb{P}^n\}.$$

Finally, we define

$$CJ(B^n(r), \mathbb{R}^n) = \bigcup \{J_f(B^n(r), \mathbb{R}^n) \mid f : X \rightarrow \mathbb{R}^n \text{ is continuous,} \\ \text{with } X \in ANR\}.$$

It is well known (see: [11,6,7,10]) that the notion of topological degree for multivalued maps in the class $CJ(B^n(r), \mathbb{R}^n)$ is available. To define it we need an appropriate notion of homotopy in $CJ(B^n(r), \mathbb{R}^n)$.

Definition 1. Let $\varphi_1, \varphi_2 \in CJ(B^n(r), \mathbb{R}^n)$ be two maps of the form

$$\begin{aligned} \varphi_1 &= f_1 \circ F_1 & B^n(r) &\xrightarrow{F_1} X \xrightarrow{f_1} \mathbb{R}^n \\ \varphi_2 &= f_2 \circ F_2 & B^n(r) &\xrightarrow{F_2} X \xrightarrow{f_2} \mathbb{R}^n. \end{aligned}$$

We say that φ_1 and φ_2 are *homotopic* in $CJ(B^n(r), \mathbb{R}^n)$ if there exists an u.s.c. R_δ -valued homotopy $\chi : B^n(r) \times [0, 1] \multimap X$ and a continuous homotopy $h : X \times [0, 1] \rightarrow \mathbb{R}^n$ satisfying:

- (i) $\chi(u, 0) = F_1(u), \quad \chi(u, 1) = F_2(u) \quad \text{for every } u \in B^n(r),$
- (ii) $h(x, 0) = f_1(x), \quad h(x, 1) = f_2(x) \quad \text{for every } x \in X,$
- (iii) for every $(u, \lambda) \in S^{n-1}(r) \times [0, 1]$ and $x \in \chi(u, \lambda)$ we have $h(x, \lambda) \neq 0$.

The map $H : B^n(r) \times [0, 1] \multimap \mathbb{R}^n$ given by

$$H(u, \lambda) = h(\chi(u, \lambda), \lambda)$$

is called a *homotopy* in $CJ(B^n(r), \mathbb{R}^n)$ between φ_1 and φ_2 .

By using approximation results for multivalued maps (see: [11,6,7,10]) one can prove the following theorem concerning the construction of a degree for maps $\varphi \in CJ(B^n(r), \mathbb{R}^n)$.

Theorem 2. *There exists a map*


$$\text{Deg} : CJ(B^n(r), \mathbb{R}) \rightarrow \mathbb{Z},$$

called topological degree function, satisfying the following properties:

- (i) *If $\varphi \in CJ(B^n(r), \mathbb{R}^n)$ is of the form $\varphi = f \circ F$ with F single valued and continuous, then $\text{Deg}(\varphi) = \text{deg}(\varphi)$, when $\text{deg}(\varphi)$ stands for the ordinary Brouwer degree of the single valued continuous map $\varphi : B^n(r) \rightarrow \mathbb{R}^n$.*
- (ii) *If $\text{Deg}(\varphi) \neq 0$, where $\varphi \in CJ(B^n(r), \mathbb{R}^n)$, then there exists $u \in B_0^n(r)$ such that $0 \in \varphi(u)$.*
- (iii) *If $\varphi \in CJ(B^n(r), \mathbb{R}^n)$ and $\{u \in B^n(r) \mid 0 \in \varphi(u)\} \subset B_0^n(\tilde{r})$ for some $0 < \tilde{r} < r$, then the restriction $\tilde{\varphi}$ of φ to $B^n(\tilde{r})$ is in $CJ(B^n(\tilde{r}), \mathbb{R}^n)$ and $\text{Deg}(\tilde{\varphi}) = \text{Deg}(\varphi)$.*
- (iv) *Let $\varphi_1, \varphi_2 \in CJ(B^n(r), \mathbb{R}^n)$ be two maps of the form*

$$\begin{aligned} \varphi_1 &= f_1 \circ F_1, & B^n(r) &\xrightarrow{F_1} X \xrightarrow{f_1} \mathbb{R}^n \\ \varphi_2 &= f_2 \circ F_2, & B^n(r) &\xrightarrow{F_2} Y \xrightarrow{f_2} \mathbb{R}^n, \end{aligned}$$

where $X, Y \in \text{ANR}$. If there exists a continuous map $h : X \rightarrow Y$ such that the diagram



is commutative, that is $F_2 = h \circ F_1$ and $f_1 = f_2 \circ h$, then $\text{Deg}(\varphi_1) = \text{Deg}(\varphi_2)$.

- (v) *If φ_1, φ_2 are homotopic in $CJ(B^n(r), \mathbb{R}^n)$, then $\text{Deg}(\varphi_1) = \text{Deg}(\varphi_2)$.*

2 Construction of a random topological degree

For details concerning this section we recommend [8] where the present state of a random topological degree and a random periodic problem for differential inclusions is presented.

Let (Ω, Σ) be a measurable space and $\varphi : \Omega \multimap X$ be a multivalued mapping with nonempty values; φ is called *measurable* if $\{\omega \in \Omega \mid \varphi(\omega) \subset A\} \in \Sigma$ for every closed A in a metric space X .

If X is a metric space we shall use the symbol $\mathcal{B}(X)$ to denote the *Borel σ -algebra* on X . Then $\Sigma \otimes \mathcal{B}(X)$ denotes the smallest σ -algebra on $\Omega \times X$ which contains all the sets $A \times B$, where $A \in \Sigma$ and $B \in \mathcal{B}(X)$. We say that a single valued map $v : X \rightarrow Y$ is a selection of $\varphi : X \multimap Y$ (written $v \subset \varphi$) provided $v(x) \in \varphi(x)$ for every $x \in X$.

The following lemma is crucial in what follows:

Lemma 3 ([8]). *Let X be a separable metric space, A be a closed subset of X and $\varphi : \Omega \times A \multimap X$ be a measurable mapping (with respect to the σ -algebra $\Sigma \otimes \mathcal{B}(A)$) with compact nonempty values. Assume further that for every $\omega \in \Omega$ the set $\text{Fix } \varphi_\omega = \{x \in X \mid x \in \varphi(\omega, x)\}$ is compact and nonempty. Then the map $F : \Omega \multimap X$ defined by:*

$$F(\omega) = \text{Fix } \varphi_\omega \quad \text{for every } \omega \in \Omega,$$

has a measurable selection, where $\varphi_\omega(x) = \varphi(\omega, x)$ for every $x \in A$.

Sketch of proof. First, let us define the function $f : \Omega \times A \rightarrow [0, +\infty)$ as follows:

$$f(\omega, x) = \text{dist}(x, \varphi(\omega, x)) = \inf\{d(x, y) \mid y \in \varphi(\omega, x)\}$$

for every $\omega \in \Omega$ and $x \in A$. Since φ is measurable one can get that f is measurable.

Now, it is obvious that the graph

$$\Gamma_F = \{(\omega, x) \in \Omega \times X \mid x \in F(\omega)\}$$

of F is equal to

$$f^{-1}(0) = \{(\omega, x) \in \Omega \times A \mid f(\omega, x) = 0\}.$$

It implies that Γ_F is a measurable subset of $\Omega \times X$ and consequently by virtue of Aumann's selection theorem there exists a measurable selection of F . \square

Definition 4. A multivalued map $\varphi : \Omega \times Y \multimap X$ with compact nonempty values is called a *random operator* provided φ is measurable and satisfies the following condition:

$$\varphi(\omega, \cdot) : Y \multimap X \text{ is u.s.c. for every } \omega \in \Omega. \tag{4}$$

Now, assume that $Y \subset X$ and $\varphi : \Omega \times Y \multimap X$ is a random operator. We say that φ has a *random fixed point* provided there exists a single valued measurable map $\xi : \Omega \rightarrow Y$, called the random fixed point of φ , such that:

$$\xi(\omega) \in \varphi(\omega, \xi(\omega)) \quad \text{for every } \omega \in \Omega.$$

We let

$$\text{Fix}^{ra}(\varphi) = \{\xi : \Omega \rightarrow Y \mid \xi \text{ is a random fixed point of } \varphi\}.$$

In view of Lemma 3 it is easy to see that in many cases existence of deterministic fixed points implies existence of random fixed points. Namely, we have:

Proposition 5. *Let X be a separable AR-space and $\varphi : \Omega \times X \multimap X$ be a random operator. Assume further that φ has R_δ -values and $\overline{\varphi_\omega(X)}$ is compact for every $\omega \in \Omega$. Then φ has a random fixed point.*

Sketch of proof. In fact, in view of Schauder Fixed Point Theorem (see [10]) we get that $\text{Fix } \varphi_\omega$ is compact and nonempty. Then the map $F : \Omega \multimap X$, $F(\omega) = \text{Fix } \varphi_\omega$ has a measurable selection $\xi \in \mathcal{F}$ (comp. Lemma 3). Of course, $\xi \in \text{Fix}^{ra}(\varphi)$. \square

Note (comp. [4,10,11]) that Proposition 5 can be formulated in many other cases.

Below, we would like to show that the topological degree theory considered in Section 1 can be taken up onto the random case (see: [7,8]).

According to Section 1 we shall use the following notations.

For any $X \in ANR$ we let:

$$J^{ra}(\Omega \times B^n(r), X) = \{F : \Omega \times B^n(r) \multimap X \mid F \text{ is random operator with } R^\delta\text{-values}\};$$

for any continuous $f : X \rightarrow \mathbb{R}^n$ we let

$$J_f^{ra}(\Omega \times B^n(r), \mathbb{R}^n) = \{\varphi : \Omega \times B^n(r) \multimap \mathbb{R}^n \mid \varphi = f \circ F \text{ for some } F \in J^{ra}(\Omega \times B^n(r), X) \text{ and } \varphi(\Omega \times S^{n-1}(r)) \subset \mathbb{P}^n\};$$

finally, we define

$$CJ^{ra}(\Omega \times B^n(r), \mathbb{R}^n) = \bigcup \{J_f^{ra}(\Omega \times B^n(r), \mathbb{R}^n) \mid f : X \rightarrow \mathbb{R}^n \text{ is continuous and } X \in ANR\}.$$

In the set $CJ^{ra}(\Omega \times B^n(r), \mathbb{R}^n)$ we can introduce the appropriate notion of homotopy (comp. Section 1 for deterministic case or [8]).

Now, observe that if $\varphi \in CJ^{ra}(\Omega \times B^n(r), \mathbb{R}^n)$, then $\varphi_\omega \in CJ(\Omega \times B^n(r), \mathbb{R}^n)$ for every $\omega \in \Omega$ and, consequently, topological degree $\text{Deg}(\varphi_\omega)$ of φ_ω is well defined (see Section 1 or [7]). Therefore we are allowed to define:

Definition 6. We define a multivalued map $\mathcal{D} : CJ^{ra}(\Omega \times B^n(r), \mathbb{R}^n) \multimap \mathbb{Z}$ by putting

$$\mathcal{D}(\varphi) = \{\text{Deg}(\varphi_\omega) \mid \omega \in \Omega\}.$$

Then the map \mathcal{D} is called the *random topological degree* on $CJ^{ra}(\Omega \times B^n(r), \mathbb{R}^n)$; we say that the random topological degree $\mathcal{D}(\varphi)$ of φ is different from zero (written $\mathcal{D}(\varphi) \neq 0$) if and only if $\text{Deg}(\varphi_\omega) \neq 0$ for every $\omega \in \Omega$.

Finally, let us remark that Theorem 2 holds true for random operators. We recommend also [6] and [10] for further possible consequences of the random topological degree.

3 The Poincaré operator

In this section we define the Poincaré translation map along trajectories of ordinary differential equations ([1,2,3,4,5,7,8,9,10,11,12]): A map $f : [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called *Carathéodory* if it satisfies:

$$t \rightarrow f(t, x) \text{ is measurable for every } x \in \mathbb{R}^n, \tag{5}$$

$$x \rightarrow f(t, x) \text{ is continuous for almost all } t \in [0, a], \tag{6}$$

$$\|f(t, x)\| \leq \mu(t)(1 + \|x\|) \text{ for every } (t, x) \in [0, a] \times \mathbb{R}^n, \tag{7}$$

where $\mu : [0, a] \rightarrow [0, +\infty)$ is an integrable function.

For a Carathéodory map $f : [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ we shall consider the following two problems:

Cauchy problem

$$\begin{cases} x'(t) = f(t, x(t)), \\ x(0) = x_0, \end{cases} \tag{8}$$

and

Periodic problem

$$\begin{cases} x'(t) = f(t, x(t)), \\ x(0) = x(a), \end{cases} \tag{9}$$

where a solution $x : [0, 1] \rightarrow \mathbb{R}^n$ is an absolutely continuous map such that:

$$x'(t) = f(t, x(t)) \text{ for almost all } t \in [0, a].$$

For each $x_0 \in \mathbb{R}^n$ we denote by

$$S^f(x_0) = \{x : [0, 1] \rightarrow \mathbb{R}^n \mid x \text{ is a solution of (8)}\}$$

and by

$$P^f = \{x : [0, a] \rightarrow \mathbb{R}^n \mid x \text{ is a solution of (9)}\}.$$

We recall the well known result so called Aronszajn Theorem (comp. [7,10,12]).

Theorem 7 (Aronszejn). *If $f : [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Carathéodory map, then the map*

$$S_f : \mathbb{R}^n \rightarrow C([0, a], \mathbb{R}^n)$$

defined by $S_f(x) = S^f(x)$, for every $x \in \mathbb{R}^n$, is an u.s.c. map with R_δ -values, where $C([0, a], \mathbb{R}^n)$ is a Banach space of continuous mappings with the usual max-norm.

Now, we wish to study problem (9). To do it we shall consider the diagram:

$$\mathbb{R}^n \xrightarrow{S_f} C([0, a], \mathbb{R}^n) \xrightarrow{e_t} \mathbb{R}^n,$$

where for every $t \in [0, a]$ the map e_t is defined by $e_t(x) = x(0) - x(t)$.

For any $t \in [0, a]$ the map $P_f^t = e_t \circ S_f$ is called the *Poincaré translation map* associated to the problem (8).

The following proposition is self-evident:

Proposition 8. *Problem (9) has a solution if and only if there is $x \in \mathbb{R}^n$ such that $0 \in P_f^a(x)$.*

In what follows we can assume, without loss of generality, that $0 \notin P_f^a(x)$ for every $x \in \mathbb{R}^n$ such that $\|x\| = r$ for some $r > 0$. Then we have:

Theorem 9. *If $\text{Deg}(P_f^a) \neq 0$, then $P^f \neq \emptyset$, where we consider P_f^a as a mapping in $CJ(B^n(r), \mathbb{R}^n)$ for r given above.*

Proof. Since S_f is u.s.c. with R_δ -values and e_a is continuous, we have that

$$P_f^a \in CJ(B^n(r), \mathbb{R}^n), \text{ for } X = C([0, a], \mathbb{R}^n)$$

being an AR -space. □

Therefore by our assumption $\text{Deg}(P_f^a)$ on $B^n(r)$ is well defined. Consequently, our result follows from Theorem 2, (ii) and Proposition 8.

In order to show that $\text{Deg}(P_f^a) \neq 0$ we shall use the *guiding potential* introduced by Liapunov (comp. [16,17,18]) and subsequently developed by Krasnoselski (comp. [13]), Mawhin [14] and others (see: [7,8,10,15]).

4 Guiding potentials

The guiding potential method has been recently employed in [10,15] to study periodic problems (9). Unlike these papers, where the guiding potential V is supposed to be C^1 , here we assume that V is only locally Lipschitzean (see: [7]). We recall some notions of Clarke generalized gradient calculus [9] we shall need later.

Let $V: \mathbb{R}^n \rightarrow \mathbb{R}$ be a locally Lipschitzean function. For $x_0 \in \mathbb{R}^n$ and $v \in \mathbb{R}^n$, let $V^0(x_0, v)$ be the generalized directional derivative of V at x_0 in the direction v , that is

$$V^0(x_0, v) = \limsup_{\substack{x \rightarrow x_0 \\ t \rightarrow 0^+}} \frac{V(x + tv) - V(x_0)}{t}.$$

Then the *generalized gradient* $\partial V(x_0)$ of V at x_0 is defined by

$$\partial V(x_0) = \{y \in \mathbb{R}^n \mid \langle y, v \rangle \leq V^0(x_0, v) \text{ for every } v \in \mathbb{R}^n\}.$$

The map $\partial V: \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by the above equality is u.s.c. with nonempty compact convex values ([9, pp. 27, 29]). If, in addition, V is also convex, then $\partial V(x_0)$ coincides with the subdifferential of V at x_0 in the sense of convex analysis ([10, p. 36]), that is

$$\partial V(x_0) = \{y \in \mathbb{R}^n \mid \langle y, x - x_0 \rangle \leq V(x) - V(x_0) \text{ for every } x \in \mathbb{R}^n\}.$$

If V is C^1 , then $\partial V(x_0)$ reduces to the singleton set $\{\text{grad } V(x_0)\}$ ([10, p. 33]).

Definition 10. Let $V : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be locally Lipschitzean. If, for some $r_0 > 0$, V satisfies

$$\langle \partial V(x), \partial V(x) \rangle^- > 0 \quad \text{for every } \|x\| \geq r_0, \tag{10}$$

then V is called a *direct potential*, where $\langle \partial V(x), \partial V(x) \rangle^- = \inf\{\langle u, v \rangle \mid u, v \in \partial V(x)\}$. If, in addition, V is convex and instead of (4.1) satisfies

$$0 \notin \partial V(x) \quad \text{for every } \|x\| \geq r_0, \tag{11}$$

then V is called a *direct convex potential*.

Observe that Definition 10 implies (11), the converse is not true in general. Moreover, if V is C^1 , then either (10) or (11) is equivalent to saying that $\text{grad } V(x) \neq 0$ for every $\|x\| \geq r_0$. In view of that, the above definition can be interpreted as a generalization of the definition of a direct potential V in the sense of Krasnoselskiĭ [13] (see also [14,15,18]), where V is assumed to be a C^1 function.

Let $V : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a direct potential. Observe that $\partial V \in CJ(B^n(r), \mathbb{R}^n)$ if $r \geq r_0$, and thus, by Theorem 2, the topological degree $\text{Deg}(\partial V)$ is well defined and independent of r . Hence, it makes sense to define the index $\text{Ind}(V)$ of the direct potential V , by putting

$$\text{Ind}(V) = \text{Deg}(\partial V),$$

when ∂V stands for the restriction of ∂V to $B^n(r), r \geq r_0$. Of course the definition of $\text{Ind}(V)$ is analogous if V is direct convex potential.

Proposition 11 ([6]). *If $V : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a direct potential (or a direct convex potential) satisfying $\lim_{\|x\| \rightarrow +\infty} V(x) = +\infty$, then $\text{Ind}(V) = 1$.*

A connection between the notion of direct potential and ordinary differential equations is given by the following:

Definition 12. Let $f : [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a Carathéodory map. A direct potential $V : \mathbb{R}^n \rightarrow \mathbb{R}$ is called a *guiding potential* for f if there exists $r_0 > 0$ such that:

$$\langle f(t, x), \partial V(x) \rangle^- \geq 0 \quad \text{for every } t \text{ and } \|x\| \geq r_0. \tag{12}$$

Our main result of this section is the following:

Theorem 13. *Assume that $f : [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Carathéodory map and $V : \mathbb{R}^n \rightarrow \mathbb{R}$ is a guiding potential for f such that $\text{Ind}(V) \neq 0$. Then $P^f \neq \emptyset$.*

In the proof of Theorem 13 we proceed analogously as in the proof of (4.4) in [12] (comp. also [6]). By the homotopy property of the topological degree finally we obtain $\text{Deg}(P_f^a) = |\text{Ind}(V)|$ and therefore our result follows from Theorem 9.

All technical details are omitted here and we left them to the reader.

5 The random case

In this section we would like point out that all results of section 3 and 4 can be formulated in the random case. We shall restrict our considerations to showing formulations and some ideas of the proofs (we recommend for more details [8]).

Definition 14. Let $f : \Omega \times [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a single valued map such that

$$f(\cdot, \cdot, x) : \Omega \times [0, a] \rightarrow \mathbb{R}^n \text{ is measurable,} \quad (13)$$

$$f(\omega, t, \cdot) : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ is continuous,} \quad (14)$$

$$\begin{aligned} \|f(\omega, t, x)\| &\leq \mu(\omega, t)(1 + \|x\|) \text{ for every } t, \omega \text{ and } x, \text{ where} \\ \mu : \Omega \times [0, a] &\rightarrow [0, +\infty) \text{ is a map such that } \mu(\cdot, t) \text{ is mea-} \\ &\text{surable and } \mu(\omega, \cdot) \text{ is Lebesgue integrable.} \end{aligned} \quad (15)$$

Then f is called a random Carathéodory operator.

Now, for given random Carathéodory operator and a measurable map $\xi_0 : \Omega \rightarrow \mathbb{R}^n$ we consider the following Cauchy problem:

$$\begin{cases} x'(\omega, t) = f(\omega, t, x(\omega, t)) \\ x(\omega, 0) = \xi_0(\omega), \end{cases} \quad (16)$$

where the solution $x : \Omega \times [0, a] \rightarrow \mathbb{R}^n$ is a map such that $x(\cdot, t)$ is measurable, $x(\omega, \cdot)$ is absolutely continuous and the derivative $x'(\omega, t)$ is considered with respect to t .

Moreover we consider the following *random periodic problem*:

$$\begin{cases} x'(\omega, t) = f(\omega, t, x(\omega, t)), \\ x(\omega, t) = x(\omega, a). \end{cases} \quad (17)$$

To solve it we need the random Poincaré translation operator. Observe that for every $\omega \in \Omega$ and $y \in \mathbb{R}^n$ we can consider the following Cauchy problem:

$$\begin{cases} x'(t) = f_\omega(t, x(t)) = f(\omega, t, x(t)), \\ x(0) = y. \end{cases} \quad (18)$$

Now, we define the following map:

$$\begin{aligned} P : \Omega \times \mathbb{R}^n &\rightarrow C([0, a], \mathbb{R}^n), \\ P(\omega, y) &= S^{f_\omega}(y). \end{aligned}$$

We have:

Proposition 15 ([8]). *Under the above assumptions the map P is a random operator.*

Note that for the proof of Proposition 15 we need the Fubini and Kuratowski Ryll-Nardzewski Selection Theorem (see [8]).

Consequently, the map

$$P_a = e_a \circ P \quad (19)$$

is called the random Poincaré operator associated to (17).

Now we are able to formulate Proposition 8, Theorem 9 and results of Section 4 for the random periodic problem. We refer to do it to the reader (comp. [8]).

References

1. Andres, J., *Periodic derivatives of solutions to nonlinear differential equations*, Czech. Math. J. **40**, 3 (1990), 353–360
2. ———, *Derivo-periodic boundary value problems for nonautonomous ordinary differential equations*, Riv. Mat. Pura Appl., **13** (1993), 62–90
3. ———, *On the multivalued Poincaré operator*, Topological Methods in Nonlinear Analysis (to appear)
4. Andres, J., Górniewicz, L. and Lewicka, M., *Partially dissipative periodic processes*, Topology in Nonlinear Analysis (K. Geba and L. Górniewicz, eds.) **35** (1996), Banach Center Publications, Warszawa, 109–118
5. Andres, J., Gabor, G. and Górniewicz, L., *Boundary value problems on infinite intervals*, Trans. Amer. Math. Soc. (to appear)
6. Bader, R. and Kryszewski, W., *Fixed point index for compositions of set-valued maps with proximally ∞ -connected values on arbitrary ANR's*, Set Valued Analysis **2**, 3 (1994), 459–480
7. De Blasi, F. S., Górniewicz, L. and Pianigiani, G., *Topological degree and periodic solutions of differential inclusions*, Nonlinear Analysis TMA (to appear)
8. ———, *Random topological degree and differential inclusions*, reprint
9. Clarke, F., *Optimization and nonsmooth analysis*, John Wiley, New York, 1983
10. Górniewicz, L., *Topological approach to differential inclusions*, Topological Methods in Differential Equations and Inclusions (M. Frigon and A. Granas, eds.), NATO ASI Series C 472, Kluwer Academic Publ., 1995, 129–190
11. Górniewicz, L., Granas, A. and Kryszewski, W., *On the homotopy method in the fixed point index theory of multivalued mappings of compact ANR's*, J. Math. Anal. Appl. **161** (1991), 457–473
12. Górniewicz, L. and Plaskacz, S., *Periodic solutions of differential inclusions in R^n* , Boll. Un. Mat. Ital. **7** (7-A 1993), 409–420
13. Krasnoselskiĭ M. A. and Zabreiko, P., *Geometric methods of nonlinear analysis*, Springer-Verlag, Berlin, 1975
14. Mawhin, J., *Topological degree methods in nonlinear boundary value problems*, Regional Conference in Mathematics, vol. 40, American Mathematical Society, Providence, R. I., 1979
15. Plaskacz, S., *Periodic solutions of differential inclusions on compact subsets of R^n* , J. Math. Anal. Appl. **148** (1990), 202–212
16. Yorke, J. A., *Extending Liapunov's second method to non Lipschitz Liapunov functions*, Bull. Amer. Math. Soc. **2** (1968), 322–325

17. ———, *Differential inequalities and non Lipschitz scalar functions*, Math. Systems Theory **4** (1970), 140–151
18. Yoshizawa, T., *Stability theory by Liapunov's second method*, Math. Soc. Japan, Tokyo, 1996