

Michael E. Filippakis

Periodic solutions for systems with nonsmooth and partially coercive potential

*Archivum Mathematicum*, Vol. 42 (2006), No. 3, 225--232

Persistent URL: <http://dml.cz/dmlcz/108000>

## Terms of use:

© Masaryk University, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**PERIODIC SOLUTIONS FOR SYSTEMS WITH NONSMOOTH  
AND PARTIALLY COERCIVE POTENTIAL**

MICHAEL E. FILIPPAKIS

ABSTRACT. In this paper we consider nonlinear periodic systems driven by the one-dimensional  $p$ -Laplacian and having a nonsmooth locally Lipschitz potential. Using a variational approach based on the nonsmooth Critical Point Theory, we establish the existence of a solution. We also prove a multiplicity result based on a nonsmooth extension of the result of Brezis-Nirenberg [3] due to Kandilakis-Kourogenis-Papageorgiou [13].

## 1. INTRODUCTION

The purpose of this paper is to prove an existence and a multiplicity result for nonlinear periodic systems driven by the one-dimensional  $p$ -Laplacian with nonsmooth Laplacian.

Recently there has been an increasing interest for problems involving the one-dimensional  $p$ -Laplacian and various solvability techniques were used. We mention the works of Dang-Oppenheimer [6], Del Pino-Manasevich-Murua [7], Fabry-Fayyad [8], Gasinski-Papageorgiou [9], Guo [10], Manasevich-Mawhin [16] and the references therein. From the above works Gasinski-Papageorgiou use a variational approach, while the others use degree theory combined with techniques from nonlinear analysis and the right hand side nonlinearity is continuous (i.e. the corresponding potential function is  $C^1$ ). Also we should mention that in Dang-Oppenheimer, Guo and Manasevich-Mawhin the right hand side nonlinearity also depends on  $x'$  and consequently their hypotheses are stronger. Here the potential function  $j(t, x)$  is only measurable in  $t \in T$  and locally Lipschitz in  $x \in \mathbb{R}^N$  (not necessarily  $C^1$ ). We assume that  $j(t, \cdot)$  is only partially coercive, i.e.  $j(t, x) \rightarrow +\infty$  as  $\|x\| \rightarrow \infty$  uniformly for almost all  $t \in E \subseteq T$ , with  $|E| > 0$  (here by  $|\cdot|$  we denote the Lebesgue measure on  $\mathbb{R}$ ). This way we extend the very recent work of Tang-Wu [18] where  $p = 2$  (semilinear problem) and the potential function  $j(t, \cdot)$  is

---

2000 *Mathematics Subject Classification*: 34A60.

*Key words and phrases*: locally linking Lipschitz function, generalized subdifferential, nonsmooth critical point theory, nonsmooth Palais-Smale condition,  $p$ -Laplacian, periodic system.

The author was supported by a grant of the National Scholarship Foundation of Greece (I.K.Y.).

Received January 14, 2005.

$C^1$  (smooth problem). Initially semilinear problems with fully coercive potential, were studied by Berger-Schechter [2] and Mawhin-Willem [17].

Our approach is variational and it is based on the nonsmooth Critical Point Theory as this was formulated by Chang [4] and extended recently by Kourogenis-Papageorgiou [14]. The multiplicity result that we prove is based on a recent nonsmooth extension of the result of Brezis-Nirenberg [3] due to Kandilakis-Kourogenis-Papageorgiou [13].

### 2. MATHEMATICAL BACKGROUND

Let  $X$  be a Banach space,  $X^*$  its topological dual. By  $\langle \cdot, \cdot \rangle$  we denote the duality brackets for the pair  $(X, X^*)$ . Given a locally Lipschitz function  $\varphi : X \rightarrow \mathbb{R}$ , the *generalized directional derivative* of  $\varphi$  at  $x \in X$  in the direction  $h \in X$ , is defined by

$$\varphi^0(x; h) \stackrel{df}{=} \limsup_{\substack{x' \rightarrow x \\ \lambda \downarrow 0}} \frac{\varphi(x' + \lambda h) - \varphi(x')}{\lambda}.$$

The function  $h \rightarrow \varphi^0(x; h)$  is sublinear, continuous and so it is the support function of a nonempty,  $w^*$ -compact, convex set  $\partial\varphi(x) \subseteq X^*$  defined by

$$\partial\varphi(x) \stackrel{df}{=} \{x^* \in X^* : \langle x^*, h \rangle \leq \varphi^0(x; h) \text{ for all } h \in X\}.$$

The multifunction  $x \rightarrow \partial\varphi(x)$  is known as the *generalized* (or *Clarke*) *subdifferential* of  $\varphi$ . If  $\varphi$  is continuous convex (hence locally Lipschitz), then the generalized subdifferential and the subdifferential in the sense of convex analysis coincide. Also if  $\varphi \in C^1(X)$  (hence it is locally Lipschitz), then  $\partial\varphi = \{\varphi'(x)\}$ .

A point  $x \in X$  is a critical point of the locally Lipschitz function  $\varphi : X \rightarrow \mathbb{R}$ , if  $0 \in \partial\varphi(x)$ . A local extremum of  $\varphi$  is a critical point. The well-known Palais-Smale condition (PS-condition for short), in the present nonsmooth setting takes the following form:

“A locally Lipschitz function  $\varphi : X \rightarrow \mathbb{R}$  satisfies the nonsmooth PS-condition, if every sequence  $\{x_n\}_{n \geq 1} \subseteq X$  such that  $|\varphi(x_n)| \leq M_1$  for some  $M_1 > 0$ , all  $n \geq 1$  and  $m(x_n) = \inf \{\|x^*\| : x^* \in \partial\varphi(x_n)\} \rightarrow 0$  as  $n \rightarrow \infty$ , has a strongly convergent subsequence.”

### 3. EXISTENCE THEOREM

The nonlinear, nonsmooth periodic system under consideration is the following:

$$(3.1) \quad \begin{cases} (\|x'(t)\|^{p-2}x'(t))' \in \partial j(x(t)) & \text{a.e. on } T = [0, b] \\ x(0) = x(b), \quad x'(0) = x'(b), & 2 \leq p < \infty. \end{cases}$$

Here by  $\partial j(t, x)$  we denote the Clarke subdifferential of the locally Lipschitz potential function  $j(t, \cdot)$ . Our hypotheses on  $j(t, x)$  are the following:

- $H(j)_1$ :  $j : T \times \mathbb{R}^N \rightarrow \mathbb{R}$  is a function such that  $j = j_1 + j_2$  and for  $i = 1, 2$ ;
  - (i) for all  $x \in \mathbb{R}^N$ ,  $t \rightarrow j_i(t, x)$  is measurable;
  - (ii) for almost all  $t \in T$ ,  $x \rightarrow j_i(t, x)$  is locally Lipschitz;

- (iii) for every  $M > 0$ , there exists  $\alpha_M \in L^1(T)$  such that
 
$$\sup [ |j(t, x)|, \|u\| : \|x\| \leq M, u \in \partial j(t, x) ] \leq \alpha_M(t) \quad \text{a.e. on } T;$$
- (iv)  $j_1(t, x) \rightarrow +\infty$  as  $\|x\| \rightarrow \infty$  uniformly for almost all  $t \in E$ ,  $|E| > 0$  and there exists  $\xi \in L^1(T)$  such that for almost all  $t \in T$  and all  $x \in \mathbb{R}^N$   $\xi(t) \leq j_1(t, x)$ ;
- (v) there exists  $\theta \in L^1(T)$  such that for almost all  $t \in T$ , all  $x \in \mathbb{R}^N$  and all  $u \in \partial j_2(t, x)$ ,  $\|u\| \leq \theta(t)$  and  $\int_0^b j_2(t, x) dt \geq -c_0$  for all  $x \in \mathbb{R}^N$  with  $c_0 > 0$ .

In the proof of our existence theorem we shall need the following auxiliary result due to Tang-Wu [18] (see Lemma 3) relating uniform coercivity and subadditivity.

**Lemma 3.1.** *If  $j : T \times \mathbb{R}^N \rightarrow \mathbb{R}$  is a function such that for all  $x \in \mathbb{R}^N$ ,  $t \rightarrow j(t, x)$  is measurable, for almost all  $t \in T$   $x \rightarrow j(t, x)$  is continuous, for every  $M > 0$  there exists  $\alpha_M \in L^1(T)$  such that for almost all  $t \in T$  and all  $\|x\| \leq M$ ,  $|j(t, x)| \leq \alpha_M(t)$  and  $j(t, x) \rightarrow +\infty$  as  $\|x\| \rightarrow \infty$  uniformly for almost all  $t \in E$ ,  $|E| > 0$ , then there exist  $g \in C(\mathbb{R}^N)_+$  subadditive function such that  $g(x) \rightarrow +\infty$  as  $\|x\| \rightarrow \infty$  and  $g(x) \leq \|x\| + 4$  and  $\eta \in L^1(T)$  for which we have for almost all  $t \in E$  and all  $x \in \mathbb{R}^N$   $j(t, x) \geq g(x) + \eta(t)$ .*

**Remark 3.2.** Here by  $|E|$  we denote the Lebesgue measure of  $E$ .

**Theorem 3.3.** *If hypotheses  $H(j)_1$  hold, then problem (3.1) has a solution  $x \in C^1(T, \mathbb{R}^N)$ .*

**Proof.** Let  $\varphi : W_{\text{per}}^{1,p}(T, \mathbb{R}^N) \rightarrow \mathbb{R}$  be the energy functional defined by

$$\varphi(x) = \frac{1}{p} \|x'\|_p^p + \int_0^b j(t, x(t)) dt = \frac{1}{p} \|x'\|_p^p + \int_0^b j_1(t, x(t)) dt + \int_0^b j_2(t, x(t)) dt.$$

We know (see for example Chang [4] or Hu-Papageorgiou [12]) that  $\varphi$  is locally Lipschitz. By virtue of Lemma 3.1, we can find  $E \subseteq T$ , with  $|E| > 0$  such that for almost all  $t \in E$  and all  $x \in \mathbb{R}^N$  we have

$$j_1(t, x) \geq g(x) + \eta(t)$$

with  $g \in C(\mathbb{R}^N)_+$  subadditive, coercive and  $\eta \in L^1(T)$ . We have

$$\begin{aligned} \int_0^b j_1(t, x(t)) dt &= \int_E j_1(t, x(t)) dt + \int_{T \setminus E} j_1(t, x(t)) dt \\ &\geq \int_E g(x(t)) dt + \int_E \eta(t) dt + \int_{T \setminus E} \xi(t) dt. \end{aligned}$$

Consider the following direct sum decomposition

$$W_{\text{per}}^{1,p}(T, \mathbb{R}^N) = \mathbb{R}^N \oplus V$$

with  $V = \left\{ v \in W_{\text{per}}^{1,p}(T, \mathbb{R}^N) : \int_0^b v(t) dt = 0 \right\}$ . So if  $x \in W_{\text{per}}^{1,p}(T, \mathbb{R}^N)$ , we can write in a unique way  $x = \bar{x} + \hat{x}$ , with  $\bar{x} \in \mathbb{R}^N$  and  $\hat{x} \in V$ . Exploiting the subadditivity

of  $g$ , we have

$$\begin{aligned} g(\bar{x}) &= g(x(t) - \hat{x}(t)) \leq g(x(t)) + g(-\hat{x}(t)) \quad \text{for all } t \in T, \\ &\Rightarrow g(\bar{x}) - g(-\hat{x}(t)) \leq g(x(t)) \quad \text{for all } t \in T. \end{aligned}$$

Moreover, because of Lemma 3.1 we have

$$g(-\hat{x}(t)) \leq \|\hat{x}(t)\| + 4 \leq \|\hat{x}\|_\infty + 4.$$

We have

$$\begin{aligned} \int_E g(x(t)) dt &\geq \int_E g(\bar{x}) dt - \int_E g(-\hat{x}(t)) dt \\ &= g(\bar{x})|E| - (\|\hat{x}\|_\infty + 4)|E|. \end{aligned}$$

But from the Poincare-Wirtinger inequality (see Mawhin-Willem [17], p.8) we know that

$$\|\hat{x}\|_\infty \leq b^{\frac{1}{q}} \|\hat{x}'\|_p = b^{\frac{1}{q}} \|x'\|_p.$$

So we obtain

$$\int_E g(x(t)) dt \geq g(\bar{x})|E| - \left(b^{\frac{1}{q}} \|x'\|_p + 4\right)|E|.$$

Let  $\Gamma(t) = \{(v, \lambda) \in \mathbb{R}^N \times (0, 1) : v \in \partial j_2(t, \bar{x} + \lambda \hat{x}(t)), j_2(t, \bar{x} + \hat{x}(t)) - j_2(t, \bar{x}) = (v, \hat{x}(t))_{\mathbb{R}^N}\}$ . From the Mean Value Theorem (see for example Clarke [5], p.41), we know that for almost all  $t \in T$ ,  $\Gamma(t) \neq \emptyset$ . By redefining  $\Gamma(\cdot)$  on the exceptional Lebesgue-null set, we may assume without any loss of generality that  $\Gamma(t) \neq \emptyset$  for all  $t \in [0 \cdot b]$ . We claim that for every direction  $h \in \mathbb{R}^N$  the function  $(t, \lambda) \rightarrow j_2^0(t, \bar{x} + \lambda \hat{x}(t); h)$  is measurable. Indeed from the definition of the generalized derivative, we have

$$\begin{aligned} j_2^0(t, \bar{x} + \lambda \hat{x}(t)) &= \\ &= \inf_{m \geq 1} \sup_{r, s \in Q \cap (-\frac{1}{m}, \frac{1}{m})} \frac{j_2(t, \bar{x} + \lambda \hat{x}(t) + r + sh) - j_2(t, \bar{x} + \lambda \hat{x}(t) + r)}{s}. \end{aligned}$$

Since  $j_2$  is jointly measurable (see Hu-Papageorgiou [11], p.142), it follows that  $(t, \lambda) \rightarrow j_2^0(t, \bar{x} + \lambda \hat{x}(t); h)$  is measurable. Set  $S(t, \lambda) = \partial j_2(t, \bar{x} + \lambda \hat{x}(t))$  and let  $\{h_m\}_{m \geq 1} \subseteq \mathbb{R}^N$  be a countable dense set. Because  $j_2^0(t, \bar{x} + \lambda \hat{x}(t); \cdot)$  is continuous, we have

$$\begin{aligned} GrS &= \{(t, \lambda, u) \in T \times (0, 1) \times \mathbb{R}^N : u \in S(t, \lambda)\} \\ &= \bigcap_{m \geq 1} \{(t, \lambda, u) \in T \times (0, 1) \times \mathbb{R}^N : (u, h_m)_{\mathbb{R}^N} \leq j_2^0(t, \bar{x} + \lambda \hat{x}(t); h_m)\} \\ &\Rightarrow GrS \in \mathcal{L}(T) \times B((0, 1)) \times B(\mathbb{R}^N), \end{aligned}$$

with  $\mathcal{L}(T)$  being the Lebesgue  $\sigma$ -field of  $T$  and  $B((0, 1))$  (resp.  $B(\mathbb{R}^N)$ ) the Borel  $\sigma$ -field of  $(0, 1)$  (resp. of  $\mathbb{R}^N$ ). So we can apply the Yankon-von Neumann-Aumann selection theorem (see Hu-Papageorgiou [11], p.158) to obtain measurable functions  $v : T \rightarrow \mathbb{R}^N$  and  $\lambda : T \rightarrow (0, 1)$  such that  $(v(t), \lambda(t)) \in \Gamma(t)$  for all  $t \in T$

and  $j_2(t, \bar{x} + \hat{x}(t)) - j_2(t, \bar{x}) = (v(t), \hat{x}(t))_{\mathbb{R}^N}$ ,  $v(t) \in \partial j_2(t, \bar{x} + \lambda(t)\hat{x}(t))$  a.e. on  $T$ . Using hypothesis  $H(j)_1(v)$  and the Poincare-Wirtinger inequality, we obtain

$$\begin{aligned} \int_0^b j_2(t, x(t)) dt &= \int_0^b j_2(t, \bar{x} + \hat{x}(t)) \\ &\geq \int_0^b j_2(t, \bar{x}) dt - b^{\frac{1}{p}} \|x'\|_p \|\theta\|_1. \end{aligned}$$

Thus finally we have

$$\varphi(x) \geq \frac{1}{p} \|x'\|_p^p + g(\bar{x})|E| - (b^{\frac{1}{q}} \|x'\|_p + 4)|E| - \|\xi\|_1 - c_0 - b^{\frac{1}{q}} \|x'\|_p \|\theta\|_1.$$

From this inequality and the coercivity of  $g$ , it follows that  $\varphi$  is coercive. Exploiting the compact embedding of  $W_{\text{per}}^{1,p}(T, \mathbb{R}^N)$  into  $C(T, \mathbb{R}^N)$ , we can easily check that  $\varphi$  is weakly lower semicontinuous. So by the Weierstrass theorem we can find  $x \in W_{\text{per}}^{1,p}(T, \mathbb{R}^N)$  such that  $\varphi(x) = \inf \varphi$ . Then we have  $0 \in \partial\varphi(x)$ . Let  $A : W_{\text{per}}^{1,p}(T, \mathbb{R}^N) \rightarrow W_{\text{per}}^{1,p}(T, \mathbb{R}^N)^*$  be the nonlinear operator defined by

$$\langle A(x), y \rangle = \int_0^b -\|x'(t)\|^{p-2} (x'(t), y'(t))_{\mathbb{R}^N} dt.$$

We have  $A(x) = u$  with  $u \in S^q_{\partial j(\cdot, x(\cdot))}$ . For every  $\psi \in C_0^\infty((0, b), \mathbb{R}^N)$  we have

$$\int_0^b -\|x'(t)\|^{p-2} (x'(t), \psi'(t))_{\mathbb{R}^N} dt = \int_0^b (u(t), \psi(t))_{\mathbb{R}^N} dt$$

Recalling that  $(\|x'(\cdot)\|^{p-2} x'(\cdot)) \in W^{-1,q}(T, \mathbb{R}^N) = W_0^{1,p}(T, \mathbb{R}^N)^*$  (see Adams [1], p.50), we have that

$$\langle (\|x'\|^{p-2} x')', \psi \rangle_0 = \int_0^b (u(t), \psi(t))_{\mathbb{R}^N} dt = \langle u, \psi \rangle_0,$$

where  $\langle \cdot, \cdot \rangle_0$  denotes the duality brackets for the pair  $(W_{\text{per}}^{1,p}(T, \mathbb{R}^N), W^{-1,q}(T, \mathbb{R}^N))$ . Since  $C_0^\infty((0, b), \mathbb{R}^N)$  is dense in  $W_{\text{per}}^{1,p}(T, \mathbb{R}^N)$  it follows that

$$(3.2) \quad (\|x'(t)\|^{p-2} x'(t))' = u(t) \in \partial j(t, x(t)) \quad \text{a.e. on } T.$$

Also for every  $y \in W_{\text{per}}^{1,p}(T, \mathbb{R}^N)$ , using Green's identity (integration by parts), we obtain

$$\begin{aligned} \langle A(x), y \rangle &= (\|x'(b)\|^{p-2} x'(b), y(b))_{\mathbb{R}^N} - (\|x'(0)\|^{p-2} x'(0), y(0))_{\mathbb{R}^N} \\ &\quad - \int_0^b ((\|x'(t)\|^{p-2} x'(t))', y(t))_{\mathbb{R}^N} dt \quad \text{for all } y \in W_{\text{per}}^{1,p}(T, \mathbb{R}^N)_{\mathbb{R}^N} dt. \end{aligned}$$

Because  $A(x) = u$ , and using (3.2), we obtain

$$\begin{aligned} (\|x'(b)\|^{p-2} x'(b), y(b))_{\mathbb{R}^N} &= (\|x'(0)\|^{p-2} x'(0), y(0))_{\mathbb{R}^N} \quad \text{for all } y \in W_{\text{per}}^{1,p}(T, \mathbb{R}^N), \\ \Rightarrow \|x'(b)\|^{p-2} x'(b) &= \|x'(0)\|^{p-2} x'(0), \\ \Rightarrow x'(0) &= x'(b). \end{aligned}$$

Note that since  $x \in W_{\text{per}}^{1,p}(T, \mathbb{R}^{\mathbb{N}})$ , we have  $x(0) = x(b)$ . Finally since  $\|x'\|^{p-2}x' \in W_{\text{per}}^{1,q}(T, \mathbb{R}^{\mathbb{N}}) \Rightarrow \|x'(\cdot)\|^{p-2}x'(\cdot) \in C_{\text{per}}^1(T, \mathbb{R}^{\mathbb{N}})$ . Because the map  $y \rightarrow \|y\|^{p-2}y$  is a homeomorphism of  $\mathbb{R}^{\mathbb{N}}$ , we infer that  $x' \in C_{\text{per}}(T, \mathbb{R}^{\mathbb{N}})$ , hence  $x \in C_{\text{per}}^1(T, \mathbb{R}^{\mathbb{N}})$  and it solves (3.1).  $\square$

#### 4. MULTIPLICITY RESULT

Next by strengthening our hypotheses on  $j(t, \cdot)$  with a condition about its behavior near zero, we obtain a multiplicity result for problem (3.1). For this we will need the following nonsmooth version of the Local Linking theorem due to Brezis-Nirenberg [3]. This theorem was proved recently by Kandilakis-Kourogenis-Papageorgiou [13].

**Theorem 4.1.** *If  $X$  is a reflexive Banach space such that  $X = Y \oplus V$  with  $\dim Y < +\infty$ ,  $\varphi : x \rightarrow \mathbb{R}$  is a locally Lipschitz functional which satisfies the nonsmooth PS-condition,  $\varphi(0) = 0$  and*

(a) *there exists  $r > 0$  such that*

$$\varphi(y) \leq 0 \quad \text{for } y \in Y, \|y\| \leq r \quad \text{and} \quad \varphi(v) \geq 0 \quad \text{for } v \in V, \|v\| \leq r,$$

(ii)  *$\varphi$  is bounded below and  $\inf \varphi < 0$ ,*

*then  $\varphi$  has at least two nontrivial critical points.*

Our hypotheses on the nonsmooth potential  $j(t, x)$  are the following:

$H(j)_2$ :  $j : T \times \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}$  is a function which satisfies hypotheses  $H(j)_1$  and

(vi)  $\lim_{x \rightarrow 0} \frac{pj(t, x)}{\|x\|^p} = 0$  uniformly for almost all  $t \in T$  and there exists  $r_0 > 0$  such that for almost all  $t \in T$  and all  $\|x\| \leq r_0$  we have  $j(t, x) \leq 0$ .

**Theorem 4.2.** *If hypotheses  $H(j)_2$  hold, then problem (3.1) has at least two nontrivial solutions in  $C^1(T, \mathbb{R}^{\mathbb{N}})$ .*

**Proof.** Let  $\varphi : W_{\text{per}}^{1,p}(T, \mathbb{R}^{\mathbb{N}}) \rightarrow \mathbb{R}$  be the locally Lipschitz Weyl functional defined by

$$\varphi(x) = \frac{1}{p}\|x'\|_p^p + \int_0^b j(t, x(t)) dt.$$

From the proof of Theorem 3.3 we know that  $\varphi$  is coercive, hence it satisfies the nonsmooth PS-condition (see Kourogenis-Papageorgiou [15]). As before we consider the direct sum decomposition

$$W_{\text{per}}^{1,p}(T, \mathbb{R}^{\mathbb{N}}) = \mathbb{R}^{\mathbb{N}} \oplus V$$

with  $V = \{v \in W_{\text{per}}^{1,p}(T, \mathbb{R}^{\mathbb{N}}) : \int_0^b v(t) dt = 0\}$ . By virtue of hypothesis  $H(j)_2(vi)$  given  $\varepsilon > 0$ , we can find  $\delta > 0$  such that for almost all  $t \in T$  and all  $\|x\| \leq \delta$  we have  $-\frac{\varepsilon}{p}\|x\|^p \leq j(t, x)$ . Let  $v \in V$  with  $\|v'\|_p \leq \frac{\delta}{b^{\frac{1}{q}}}$ . From the Poincare-Wirtinger

inequality we have that  $\|v\|_\infty \leq b^{\frac{1}{q}} \|v'\|_p \leq \delta$ . So if  $v \in V$  with  $\|v'\|_p \leq \frac{\delta}{b^{\frac{1}{q}}} = \delta_1$ , we have  $\|v\|_\infty \leq \delta$  and so

$$\begin{aligned} \varphi(v) &= \frac{1}{p} \|v'\|_p^p + \int_0^b j(t, v(t)) dt \\ &\geq \frac{1}{p} \|v'\|_p^p + \frac{\varepsilon}{p} \|v\|_p^p \\ &\geq \frac{1}{p} \left(1 - \frac{\varepsilon}{\beta_1}\right) \|v'\|_p^p \quad \text{for some } \beta_1 > 0, \end{aligned}$$

from the Poincare-Wirtinger inequality. Choose  $\varepsilon \leq \beta_1$ , to infer that for  $\|v\| \leq \delta_1$  we have  $\varphi(v) \geq 0$ .

Also if  $y \in \mathbb{R}^N$  and  $\|y\| \leq r_0$ , then by hypothesis  $H(j)_2(vi)$  we have that

$$\varphi(y) = \int_0^b j(t, y) dt \leq 0.$$

Note that  $\varphi$  being coercive, it is bounded below. If  $\inf \varphi < 0$ , then using  $r = \min \{\delta_1, r_0\} > 0$  we can apply Theorem 4.1 and obtain two nontrivial critical points of  $\varphi$ , which we can check are two distinct nontrivial solutions of (3.1) in  $C^1(T, \mathbb{R}^N)$ .

If  $\inf \varphi = 0$ , then by virtue of hypothesis  $H(j)_2(vi)$  for all  $y \in \mathbb{R}^N$  with  $b^{\frac{1}{p}} \|y\|_{\mathbb{R}^N} \leq \delta_1$  we have  $\inf \varphi = \varphi(y) = 0$  and so we conclude that  $\varphi$  has an infinity of critical points, therefore problem (3.1) has an infinity of solutions in  $C^1(T, \mathbb{R}^N)$ . □

The nonsmooth locally Lipschitz potential function

$$j(t, x) = \begin{cases} -\|x\|^p \ln(1 + \|x\|^p) & \text{if } \|x\| \leq 1 \\ \chi_E(t) \ln \|x\| + \chi_{E^c}(t) \sin \pi \|x\| - \ln 2 & \text{if } \|x\| \geq 1 \end{cases},$$

with  $|E| > 0$ , satisfies hypotheses  $H(j)_2$ .

### REFERENCES

- [1] Adams, R., *Sobolev Spaces*, Pure and Applied Mathematics **65**, Academic Press, New York/London 1975.
- [2] Berger, M. Schechter, M., *On the solvability of semilinear gradient operator equations*, Adv. Math. **25** (1977), 97–132.
- [3] Brezis, H., Nirenberg, L., *Remarks on finding critical points*, Comm. Pure Appl. Math. **44** (1991), 939–963.
- [4] Chang, K. C., *Variational methods for nondifferentiable functionals and their applications to partial differential equations*, J. Math. Anal. Appl. **80** (1981), 102–129.
- [5] Clarke, F., *Optimization and Nonsmooth Analysis*, Wiley, New York 1983.
- [6] Dang, H., Oppenheimer, S., *Existence and uniqueness results for some nonlinear boundary value problems*, J. Math. Anal. Appl. **198**, (1996) 35–48.
- [7] del Pino, M., Manasevich, R., Murua, A., *Existence and multiplicity of solutions with prescribed period for a second order quasilinear ODE*, Nonlinear Anal. **18**, (1992) 79–92.



- [8] Fabry, C., Fayyad, D., *Periodic solutions of second order differential equations with a  $p$ -Laplacian and asymmetric nonlinearities*, Rend. Istit. Mat. Univ. Trieste **24** (1992), 207–227.
- [9] Gasinski, L., Papageorgiou, N. S., *A multiplicity result for nonlinear second order periodic equations with nonsmooth potential*, Bull. Belg. Math. Soc. Simon Stevin **9** (2002a), 245–258.
- [10] Guo, Z., *Boundary value problems for a class of quasilinear ordinary differential equations*, Differential Integral Equations **6** (1993), 705–719.
- [11] Hu, S., Papageorgiou, N. S., *Handbook of Multivalued Analysis. Volume I: Theory*, Kluwer, Dordrecht, The Netherlands 1997.
- [12] Hu, S., Papageorgiou, N. S., *Handbook of Multivalued Analysis. Volume II: Applications*, Kluwer, Dordrecht, The Netherlands 2000.
- [13] Kandilakis, D., Kourogenis, N., Papageorgiou, N., *Two nontrivial critical point for nonsmooth functional via local linking and applications*, J. Global Optim., to appear.
- [14] Kourogenis, N., Papageorgiou, N. S., *Nonsmooth critical point theory and nonlinear elliptic equations at resonance*, J. Austral. Math. Soc. Ser. A **69** (2000), 245–271.
- [15] Kourogenis, N., Papageorgiou, N. S., *A weak nonsmooth Palais-Smale condition and coercivity*, Rend. Circ. Mat. Palermo **49** (2000), 521–526.
- [16] Manasevich, R., Mawhin, J., *Periodic solutions for nonlinear systems with  $p$ -Laplacian-like operators*, J. Differential Equations **145** (1998), 367–393.
- [17] Mawhin, J., Willem, M., *Critical Point Theory and Hamiltonian Systems*, Vol. 74 of Applied Mathematics Sciences, Springer-Verlag, New York 1989.
- [18] Tang, C. L., Wu, X. P., *Periodic solutions for second order systems with not uniformly coercive potential*, J. Math. Anal. Appl. **259** (2001), 386–397.

DEPARTMENT OF MATHEMATICS  
SCHOOL OF APPLIED MATHEMATICS AND NATURAL SCIENCES  
NATIONAL TECHNICAL UNIVERSITY, ZOGRAFOU CAMPUS  
ATHENS 15780, GREECE  
*E-mail:* mfilip@math.ntua.gr