

Lutz Heindorf

Another note on countable Boolean algebras

Commentationes Mathematicae Universitatis Carolinae, Vol. 37 (1996), No. 4, 815--819

Persistent URL: <http://dml.cz/dmlcz/118889>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Another note on countable Boolean algebras

LUTZ HEINDORF

Abstract. We prove that a Boolean algebra is countable iff its subalgebra lattice admits a continuous complementation.

Keywords: Boolean algebra, subalgebra lattice, metrizable

Classification: Primary 06E05; Secondary 54E35

The title refers to [4], where T. Jech proved that the subalgebra lattice $Sub A$ of a countable Boolean algebra A is complemented, i.e., for each $B \leq A$ there exists $B^* \leq A$ such that $B \cap B^* = \{0, 1\}$ and $B \cup B^*$ generates A . Independently this and stronger results were proved at about the same time by J.B. Remmel [6] and S. Todorčević (unpublished, his proof and many more facts on $Sub A$ are given in the survey [1]).

In Section 1 we describe a construction of complements B^* with the additional feature that for each $a \in A$, whether or not a belongs to B^* depends only on the intersection of B with a finite subalgebra of A . In other words, the mapping $B \mapsto B^*$ is continuous with respect to the natural topology on $Sub A$, a subbase of which, by definition, consists of all sets

$$\{C \in Sub A : a \in C\} \text{ and } \{C \in Sub A : a \notin C\},$$

where a runs through A . Notice that all sets $\{C \in Sub A : B \cap F = C \cap F\}$ constitute a base at the point $B \in Sub A$, when F runs through all finite subalgebras of A .

In Section 2 we prove that countability is necessary for continuous complementation. Thus

Theorem 1. *A Boolean algebra is countable iff its subalgebra lattice admits a continuous complementation.*

Our notation is in accordance with [5], with the exception that we use \vee , \wedge , and $-$ for the lattice-theoretic Boolean operations of join, meet and complementation and reserve $+$ for symmetric difference: $a + b = (a \wedge -b) \vee (b \wedge -a)$. In connection with $+$, meets are sometimes called products and denoted by \cdot instead of \wedge . Recall that each Boolean algebra is a ring with unit under this addition and multiplication. Moreover, $a + a = 0$ for all a .

1. The construction of complements

Let A be the given countable Boolean algebra. We can assume that A is infinite, for, in the finite case continuity is for free and the existence of complements guaranteed by the above mentioned results. We use the well-known fact (cf. 15.10 in [5]) that A has an ordered base, i.e., a set of generators K , say, which is a chain under the Boolean partial order. We can and will assume that $0, 1 \notin K$. Then there is the following normal form assertion, where $\langle M \rangle$ denotes the subalgebra of A generated by M .

- (1) If $L \subseteq K$, then each non-zero element of $\langle L \rangle$ can be uniquely written as $l_1 + l_2 + \dots + l_q$, where $q \geq 1$ and $l_1 < l_2 < \dots < l_q$ all belong to $L \cup \{1\}$.

It is well-known that, for arbitrary $M \subseteq A$, the subalgebra $\langle M \rangle$ consists of 0 and all finite sums of products (= meets) of elements of $M \cup \{1\}$. Being a chain, $L \cup \{1\}$ is closed under products, which yields the existence of the desired representations.

Assuming that the element a has two different representations $a = l'_1 + \dots + l'_p = l''_1 + \dots + l''_q$ we get $0 = a + a = l'_1 + \dots + l'_p + l''_1 + \dots + l''_q$ and, after cancellation of possible pairs $l'_i = l''_j$ and rearrangement, $0 = l_1 + \dots + l_r$, with $1 \leq r \leq p + q$ and $l_1 < l_2 < \dots < l_r$. The number of terms is at least two, for $0 \notin L \cup \{1\}$. But then

$$l_r = l_1 + \dots + l_{r-1} \leq l_1 \vee \dots \vee l_{r-1} = l_{r-1} < l_r,$$

which is a contradiction.

As A is countable, we can fix an injective enumeration $(k_n)_{n < \omega}$ of K . For a given subalgebra B of A we define subsets L_n^B of $\{k_i : i < n\}$ in the following inductive way.

$$L_0^B = \emptyset \quad \text{and} \quad L_{n+1}^B = \begin{cases} L_n^B, & \text{if there are } l_1, \dots, l_p \in L_n^B \ (p \geq 0!) \\ & \text{such that } k_n + l_1 + \dots + l_p \in B. \\ L_n^B \cup \{k_n\}, & \text{otherwise.} \end{cases}$$

We are now going to show that by letting B^* be the subalgebra generated by $L^B = \bigcup_{n < \omega} L_n^B$ we get the desired continuous complementation. Notice first that, by construction,

- (2) $k_n \in L^B \iff k_n \in L_{n+1}^B$.

To prove $B^* \cap B = \{0, 1\}$, we assume the contrary and consider some alleged $b \in B^* \cap B \setminus \{0, 1\}$. Passing to $1 + b$ if necessary, claim (1) yields a representation $b = k_{n_1} + \dots + k_{n_q}$ with all $k_{n_i} \in L^B$. Let n_t be maximal among the n_i . Then, by construction, $k_{n_t} \notin L_{n_t+1}^B$, hence, by (2), $k_{n_t} \notin L^B$, a contradiction.

To prove that $B \cup B^*$ generates A , it is clearly sufficient to express each k_n in the form $b + b^*$. This is trivial if $k_n \in L^B$. But otherwise $k_n \notin L_{n+1}^B$ and there is some finite (possibly zero) sum $f \in \langle L_n^B \rangle \subseteq B^*$ such that $k_n + f \in B$. So, $k_n = (k_n + f) + f$ is the desired representation.

It remains to check continuity. If a equals 0 or 1, every B^* contains it. Otherwise, according to claim (1), a can be uniquely written as

$$a = k_{n_1} + \dots + k_{n_q} \quad \text{or} \quad a = k_{n_1} + \dots + k_{n_q} + 1,$$

with $k_{n_1}, \dots, k_{n_q} \in K$. By uniqueness, $a \in B^* \Leftrightarrow k_{n_1}, \dots, k_{n_q} \in L^B$. By construction, whether or not k_n belongs to L^B depends only on the intersection of B with $\langle \{k_i : i \leq n\} \rangle$. So, whether or not a belongs to B^* depends only on the intersection of B with the finite subalgebra $\langle \{k_i : i \leq \max\{n_1, \dots, n_q\}\} \rangle$ of A .

2. The converse

In order to keep the argument short, we use Stone duality and conceive the given Boolean algebra A as $Clop X$, the algebra of all clopen (= closed and open) subsets of some compact and zero-dimensional topological space X . Accordingly, we use the set-theoretic notation for the Boolean operations. We assume that $B \mapsto B^*$ is a continuous complementation $Sub A \rightarrow Sub A$. Our aim is to show that X is metrizable.

For $x, y \in X$ we let $B(x, y)$ denote the subalgebra $\{a \in A : x \in a \Leftrightarrow y \in a\}$ of $A = Clop X$ and use $B^*(x, y)$ for its complement. Obviously

$$(3) \quad B(x, x) = A, \text{ hence } B^*(x, x) = \{\emptyset, X\}.$$

Let x and y be distinct now. If $a \in B^*(x, y)$ does not separate x and y , then it belongs to $B^*(x, y) \cap B(x, y) = \{\emptyset, X\}$ and equals \emptyset or X . Repeated application of this observation yields that

$$(4) \quad \text{if } x \neq y, \text{ then } B^*(x, y) \text{ is a four-element subalgebra of } A.$$

Indeed, consider $a, b \in B^*(x, y) \setminus \{\emptyset, X\}$. Then both a and b must separate x and y , so their symmetric difference does not and equals, therefore, \emptyset or X . So $a = b$ or $a = X \setminus b$. This shows that $B^*(x, y)$ has at most four elements. But $B^*(x, y)$ cannot be the two-element subalgebra, for, otherwise, $B(x, y) \cup B^*(x, y) = B(x, y)$ could not generate the whole of A .

Next we observe that

$$(5) \quad \text{the assignment } (x, y) \mapsto B(x, y) \text{ defines a continuous mapping } X^2 \rightarrow Sub A.$$

To check this, it is sufficient to consider the preimages of subbasic sets in the space $Sub A$. Well, $\{(x, y) : a \in B(x, y)\} = a^2 \cup (X \setminus a)^2$ and $\{(x, y) : a \notin B(x, y)\} = a \times (X \setminus a) \cup (X \setminus a) \times a$ are both clopen for each $a \in A$.

It follows that the mapping $(x, y) \mapsto B^*(x, y)$ is also continuous. Finally, we need the following claim.

$$(6) \quad \text{For every four-element subalgebra } B \text{ of } A \text{ the set}$$

$$W_B = \{(x, y) \in X^2 : B^*(x, y) = B\}$$

is clopen in X^2 and does not intersect the diagonal $\Delta = \{(x, x) : x \in X\}$.

To see this, we fix $b \in B \setminus \{\emptyset, X\}$. We already know that $B^*(x, y)$ has at most four elements. Therefore,

$$B^*(x, y) = B \iff b \in B^*(x, y).$$

It follows that W_B is clopen, being the preimage of the subbasic clopen set $\{C \in \text{Sub } A : b \in C\}$ under the continuous mapping B^* . By (3), $B^*(x, x)$ is always two-element, so W_B cannot intersect Δ .

Assertions (4) and (6) yield a representation of $X^2 \setminus \Delta$ as the disjoint union $\bigcup_B W_B$ of clopen (hence compact) subsets of X^2 . It follows that $X^2 \setminus \Delta$ is paracompact. To end our argument it remains to apply the taylormade metrization theorem of G. Gruenhage (2.6 in [3]): *The compact space X is metrizable iff $X^2 \setminus \Delta$ is paracompact.*

3. The topological version

Recall (cf. [2, 2.7.20]) that the exponential or hyperspace $\text{exp } X$ of a topological space X is the set of all non-empty closed subsets of X equipped with the Vietoris topology. A subbase of this topology consists of all

$$\{F \in \text{exp } X : F \subseteq U\} \quad \text{and} \quad \{F \in \text{exp } X : F \cap U \neq \emptyset\},$$

where U runs through all open subsets of X .

Theorem 2. *A compact and zero-dimensional space X is metrizable iff there is a continuous mapping $f : X \times \text{exp } X \rightarrow X$ such that*

$$f(x, F) \in F \quad \text{and} \quad f(x, F) = x \text{ if } x \in F.$$

We just sketch the proof. If $F \subseteq X$, then $B(F) = \{b \in \text{Clop } X : F \subseteq b \text{ or } b \cap F = \emptyset\}$ is a subalgebra of $\text{Clop } X$. Assuming that X is metrizable, the Boolean algebra $\text{Clop } X$ is countable so Theorem 1 applies and each $B(F)$ has a continuously chosen complement $B^*(F)$. A somewhat lengthy verification then shows that the desired mapping $f : X \times \text{exp } X \rightarrow X$ can be defined by

$$\{f(x, F)\} = \bigcap \{b^* \Delta b : x \in b^* \in B^*(F); b \cap F = \emptyset\}.$$

Here Δ means symmetric difference, the set-theoretic version of $+$. The reader who wants to fill in the details is advised to prove the following claim first.

- (7) Every $a \in \text{Clop } X$ can be uniquely written in the form $b^* \Delta b$, where $b \in B^*(F)$ and $b \cap F = \emptyset$.

For the other direction, one can mimic the proof in Section 2. Let the mapping $f : X \times \text{exp } X \rightarrow X$ be given. For $x, y \in X$ we let $b(x, y)$ denote the clopen set $\{z \in X : f(z, \{x, y\}) = x\}$. Then $b(x, x) = X$ and $\emptyset \neq b(x, y) \neq X$ for $x \neq y$. A routine but tedious verification shows that $W_b = \{(x, y) \in X^2 : b(x, y) = b\}$ is clopen for all $b \in \text{Clop } X \setminus \{\emptyset, X\}$. Then the decomposition

$$X^2 \setminus \Delta = \bigcup \{W_b : b \neq X\}$$

witnesses the paracompactness of $X^2 \setminus \Delta$ and Gruenhage's theorem can be applied as before.

REFERENCES

- [1] Bonnet R., *Subalgebras*, Chapter 10 in vol. 2 of: J. D. Monk (ed.) *Handbook of Boolean algebras*, North-Holland, Amsterdam, 1989.
- [2] Engelking R., *General Topology*, PWN, Warsaw, 1977.
- [3] Gruenhage G., *Covering properties on $X^2 \setminus \Delta$, W -sets, and compact subsets of Σ -products*, *Topology and its Applications* **17** (1978), 287–304.
- [4] Jech T., *A note on countable Boolean algebras*, *Algebra Universalis* **14** (1982), 257–262.
- [5] Koppelberg S., *General theory of Boolean algebras*, vol 1. of: J. D. Monk (ed.) *Handbook of Boolean algebras*, North-Holland, Amsterdam, 1989.
- [6] Remmel J.B., *Complementation in the lattice of subalgebras of a Boolean algebra*, *Algebra Universalis* **10** (1980), 48–64.

FREIE UNIVERSITÄT BERLIN, 2. MATHEMATISCHES INSTITUT, ARNIMALLEE 3,
D-14195 BERLIN, GERMANY

E-mail: heindorf@math.fu-berlin.de

(Received November 6, 1995)