

Wiktor Bartol; Francesc Rosselló
The weak hereditary class of a variety

Czechoslovak Mathematical Journal, Vol. 56 (2006), No. 2, 697–710

Persistent URL: <http://dml.cz/dmlcz/128098>

Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE WEAK HEREDITARY CLASS OF A VARIETY

WIKTOR BARTOL, Warszawa, and FRANCESC ROSSELLÓ, Palma de Mallorca

(Received December 18, 2003)

Abstract. We study the *weak hereditary class* $S_w(\mathcal{K})$ of all weak subalgebras of algebras in a total variety \mathcal{K} . We establish an algebraic characterization, in the sense of Birkhoff's HSP theorem, and a syntactical characterization of these classes. We also consider the problem of when such a weak hereditary class is weak equational.

Keywords: partial algebras, varieties, weak subalgebras, weak equations

MSC 2000: 08A55, 08B99

1. INTRODUCTION

The usefulness of partial algebras derives mostly from their extendibility to total algebras belonging to some specific variety or satisfying some other specific conditions. This feature has been frequently used in the literature, just to mention the original proof of the Grätzer-Schmidt theorem on the characterization of congruence lattices [3]. In this paper we intend to study the class $S_w(\mathcal{K})$ of all partial algebras embeddable in a variety of total algebras \mathcal{K} , and we call this class the *weak hereditary class* of \mathcal{K} . Thus, $S_w(\mathcal{K})$ can be seen as the class of all partial structural approximations of algebras in \mathcal{K} ; the notation reflects the obvious fact that a partial algebra is embeddable in \mathcal{K} if and only if it is (up to isomorphism) a weak subalgebra of an algebra in \mathcal{K} .

Clearly any algebra \mathbf{A} in $S_w(\mathcal{K})$ must have the entire equational theory $\text{Eq}(\mathcal{K})$ among the equations that are weakly valid in \mathbf{A} , i.e., $S_w(\mathcal{K}) \subseteq \text{Mod}_w(\text{Eq}(\mathcal{K}))$. As we will see several times in this paper, this condition is by no means sufficient,

This work has been partially supported by the Spanish DGES, project BFM2000-1113-C02-01.

and the converse inclusion is not generally true. It may be interesting however to distinguish those varieties of total algebras \mathcal{K} for which $S_w(\mathcal{K})$ is a weak equational class, be it for the simple reason that weak equational theories are decidable [8]. In this paper we prove some general conditions for this to hold true, and in a subsequent paper we plan to investigate exhaustively this question for unary varieties.

To end this introduction, let us fix some notations and conventions to be used throughout this paper.

Once a type of algebras Σ is fixed, by a partial algebra (resp. an equation) we mean a partial algebra (resp. an equation) of type Σ . In this paper we only consider finitary homogeneous types.

Given a partial algebra denoted by a capital letter in boldface type (\mathbf{A} , \mathbf{B} , etc.), we denote its carrier by the same capital letter in slanted type (A , B , etc.) and the operations by superscripting the operation symbol with the algebra's name ($\varphi^{\mathbf{A}}$, $\psi^{\mathbf{B}}$, ...).

Given a set \mathcal{E} of Σ -equations, we denote by $\text{Mod}_t(\mathcal{E})$ the *total variety* defined by \mathcal{E} , i.e., the class of all total Σ -algebras that satisfy \mathcal{E} . In a similar way, we denote by $\text{Mod}_w(\mathcal{E})$ the *weak equational class* defined by \mathcal{E} , i.e., the class of all partial Σ -algebras that satisfy \mathcal{E} weakly.

Given a total variety \mathcal{K} , $S_w(\mathcal{K})$ is the class of all weak subalgebras of algebras in \mathcal{K} .

All terms are supposed to have their variables in a fixed, countably infinite set of variables $\mathcal{X} = \{x_i : i \geq 1\}$. Given a term \mathbf{p} , we denote its set of explicit variables by $\text{var}(\mathbf{p})$.

2. WEAK HEREDITARY CLASSES

In the sequel, and unless otherwise stated, let $\Sigma = (\Omega, \eta)$ be a fixed type of algebras.

Definition 1. A class \mathcal{M} of partial algebras is *weak hereditary* if and only if $\mathcal{M} = S_w(\mathcal{K})$ for some total variety \mathcal{K} ; in particular, we say that $S_w(\mathcal{K})$ is the *weak hereditary class* of \mathcal{K} .

Example 2. Let \mathcal{E} be a set of equations and $\mathcal{K} = \text{Mod}_t(\mathcal{E})$ the variety defined by \mathcal{E} . If a partial algebra \mathbf{A} satisfies \mathcal{E} as existential equations, then any *inner completion* of \mathbf{A} (i.e., any completion of \mathbf{A} on its carrier set) satisfies \mathcal{E} ; thus, any existential model of \mathcal{E} belongs to $S_w(\mathcal{K})$. (Actually, it is proved in [2] that any existential model of \mathcal{E} is a *relative* subalgebra of a total model of \mathcal{E} .)

Example 3. Let now \mathcal{E} be a set of *regular* equations and $\mathcal{K} = \text{Mod}_t(\mathcal{E})$. If a partial algebra \mathbf{A} satisfies \mathcal{E} in the strong (or Kleene's) sense, then the one-point

completion of \mathbf{A} satisfies \mathcal{E} (see [7]). Therefore, any strong model of such a set \mathcal{E} belongs to $S_w(\mathcal{K})$.

More generally, if \mathcal{E} is any set of equations closed under Birkhoff's equational logic rules, then any strong model of \mathcal{E} belongs to $S_w(\mathcal{K})$: see Corollary 11 below. Thus, if \mathbf{A} satisfies strongly the *whole* equational theory $\text{Eq}(\mathcal{K})$ of a total variety \mathcal{K} , then $\mathbf{A} \in S_w(\mathcal{K})$. However, when \mathcal{E} is not closed in the Birkhoff sense, strong models of \mathcal{E} need not be weak subalgebras of total models of \mathcal{E} , as the following simple counterexample shows.

Example 4. Let Σ be a type of unary algebras with at least three different operation symbols φ_0, φ_1 and φ_2 , and let \mathbf{A} be a partial algebra with carrier $A = \{a, b\}$ and with $\varphi_0^{\mathbf{A}}(a) = \varphi_0^{\mathbf{A}}(b) = a, \varphi_1^{\mathbf{A}}(a) = \varphi_1^{\mathbf{A}}(b) = b$, and $\varphi_2^{\mathbf{A}}$ discrete. Then \mathbf{A} is a strong model of the equations

$$\varphi_0(x_1) \approx \varphi_0(x_2), \quad \varphi_1(x_1) \approx \varphi_1(x_2), \quad \varphi_0\varphi_2(x_1) \approx \varphi_1\varphi_2(x_2)$$

but no completion of \mathbf{A} can satisfy them. Indeed, any total model of these equations also satisfies $\varphi_0(x_1) \approx \varphi_1(x_2)$, while $\varphi_0^{\mathbf{A}}(a) \neq \varphi_1^{\mathbf{A}}(b)$.

This last example entails in particular that, although any algebra in $S_w(\text{Mod}_t(\mathcal{E}))$ is a weak model of \mathcal{E} because weak subalgebras preserve weak satisfaction of equations, the converse implication is in general false.

Thus, a weak model of the whole equational theory of a total variety need not belong to its weak hereditary class. Indeed, this algebra \mathbf{A} satisfies weakly the whole equational theory of $\text{Mod}_t(\{\varphi^2(x) \approx x\})$, but it does not belong to $S_w(\{\varphi^2(x) \approx x\})$.

Remark 5. A weak hereditary class is closed under isomorphisms, weak subalgebras and arbitrary products. Hence, it is quasi-equational: actually, by [1, § 3, Theorem 1], $\mathbf{A} \in S_w(\mathcal{K})$ if and only if

$$\mathbf{A} \models \bigwedge_{j \in J} \mathbf{p}_j \approx \mathbf{q}_j \rightarrow x \approx y$$

for every such quasi-equation holding in \mathcal{K} (here “ \approx ” represents an existential equation). In Theorem 12 below we give another syntactical characterization of $S_w(\mathcal{K})$.

The next result provides an algebraic characterization of weak hereditary classes in the sense of Birkhoff's HSP Theorem. Here and in the sequel, H, S, S_t and P stand for the “closed homomorphic image”, “closed subalgebra”, “total (closed) subalgebra” and “direct product” algebraic operators, respectively. Moreover, given a class \mathcal{M} of partial algebras, \mathcal{M}^* denotes the subclass of all total algebras in \mathcal{M} . Notice that $S_w(\mathcal{K})^* = \mathcal{K}$ for every total variety \mathcal{K} .

Theorem 6. Let \mathcal{M} be a class of partial algebras. The following assertions on \mathcal{M} are equivalent:

- i) \mathcal{M} is a weak hereditary class.
- ii) $\mathcal{M} = S_w HSP(\mathcal{M}^*)$.
- iii) $\mathcal{M} = S_w HS_t P(\mathcal{M})$.

Proof. (i) \implies (ii): If $\mathcal{M} = S_w(\mathcal{K})$ for some variety \mathcal{K} of total algebras, then $\mathcal{M}^* = \mathcal{K} = HSP(\mathcal{K}) = HSP(\mathcal{M}^*)$ and therefore $\mathcal{M} = S_w(HSP(\mathcal{M}^*))$.

(ii) \implies (iii): Assume $\mathcal{M} = S_w HSP(\mathcal{M}^*)$. Then

$$P(\mathcal{M}) = PS_w HSP(\mathcal{M}^*) \subseteq S_w PHSP(\mathcal{M}^*) \subseteq S_w HSP(\mathcal{M}^*) = \mathcal{M}$$

and

$$S_t(\mathcal{M}) = S_t S_w HSP(\mathcal{M}^*) \subseteq S_w HSP(\mathcal{M}^*) = \mathcal{M}.$$

Therefore $S_t P(\mathcal{M}) \subseteq \mathcal{M}$, and, in fact, $S_t P(\mathcal{M}) \subseteq \mathcal{M}^*$. Consequently,

$$S_w HS_t P(\mathcal{M}) \subseteq S_w H(\mathcal{M}^*) \subseteq S_w HSP(\mathcal{M}^*) = \mathcal{M}.$$

On the other hand, the inclusion $\mathcal{M} \subseteq S_w HS_t P(\mathcal{M})$ follows from the fact that $SP(\mathcal{M}^*) \subseteq S_t P(\mathcal{M})$ for any class \mathcal{M} (if \mathbf{A} is a closed subalgebra of a total algebra, \mathbf{A} is total too), and therefore $\mathcal{M} = S_w HSP(\mathcal{M}^*) \subseteq S_w HS_t P(\mathcal{M})$.

(iii) \implies (i): Assume now that $\mathcal{M} = S_w HS_t P(\mathcal{M})$. We will prove that, under this assumption, $HS_t P(\mathcal{M}) = HSP(\mathcal{M}^*)$. Indeed, on the one hand, the inclusion $SP(\mathcal{M}^*) \subseteq S_t P(\mathcal{M})$ used in the last step of the proof of the previous implication entails that $HSP(\mathcal{M}^*) \subseteq HS_t P(\mathcal{M})$ for any class \mathcal{M} . And on the other hand, if $\mathbf{B} \in HS_t P(\mathcal{M})$ then \mathbf{B} is total and belongs to \mathcal{M} because $\mathcal{M} = S_w HS_t P(\mathcal{M})$, and therefore $\mathbf{B} \in \mathcal{M}^* \subseteq HSP(\mathcal{M}^*)$; this proves $HS_t P(\mathcal{M}) \subseteq HSP(\mathcal{M}^*)$.

Thus, $HS_t P(\mathcal{M})$ is a variety of total algebras and \mathcal{M} is its weak hereditary class. □

Remark 7. Notice that the equality $HS_t P(\mathcal{M}) = HSP(\mathcal{M}^*)$ need not hold for each class \mathcal{M} of partial algebras. For instance, let \mathbf{A} be a non-total algebra containing a total and non-trivial closed subalgebra \mathbf{A}' , and let $\mathcal{M} = \{\mathbf{A}\}$. Then $\mathbf{A}' \in HS_t P(\mathcal{M})$, while $HSP(\mathcal{M}^*)$ only contains trivial total algebras.

In the sequel we provide an alternative syntactical characterization of weak hereditary classes. Recall that given a class \mathcal{K} of similar algebras and an algebra \mathbf{A} of the same type, a \mathcal{K} -reflection of \mathbf{A} is a homomorphism $e: \mathbf{A} \rightarrow \mathbf{B}$ with $\mathbf{B} \in \mathcal{K}$ such that every homomorphism of \mathbf{A} to an algebra in \mathcal{K} factorizes in a unique way through e . Since such an algebra \mathbf{B} , when it exists, is determined uniquely (up to isomorphism)

by \mathbf{A} , we call it the \mathcal{K} -reflection of \mathbf{A} whenever e is understood or nonrelevant. The first lemma is an easy consequence of the universal property of \mathcal{K} -reflections.

Lemma 8. *Let \mathcal{K} be a total variety and \mathbf{A} a partial algebra. Then, $\mathbf{A} \in S_w(\mathcal{K})$ if and only if \mathbf{A} is (up to isomorphism) a weak subalgebra of its \mathcal{K} -reflection.*

The next lemma describes in a simple (and suitable for our purposes) way the reflection of a partial algebra with respect to a variety of total algebras.

Lemma 9. *Let \mathbf{A} be a partial algebra and $i_{\mathbf{A}}: \mathbf{A} \hookrightarrow \mathbf{F}$ its free completion (given by the embedding of \mathbf{A} into \mathbf{F} as a relative subalgebra), and let \mathcal{K} be a total variety.*

Let $\theta(\mathcal{K})$ be the transitive closure in \mathbf{F} of the relation

$$\varepsilon(\mathcal{K}) = \{(\mathbf{p}^{\mathbf{F}}(v), \mathbf{q}^{\mathbf{F}}(v)) : \mathbf{p} \approx \mathbf{q} \in \text{Eq}(\mathcal{K}), \mathbf{v} \in A^{\mathcal{X}}\}.$$

Then $\theta(\mathcal{K})$ is a congruence on \mathbf{F} such that $\mathbf{F}/\theta(\mathcal{K}) \in \mathcal{K}$, and the composition

$$\text{nat}_{\theta(\mathcal{K})} \circ i_{\mathbf{A}}: \mathbf{A} \rightarrow \mathbf{F}/\theta(\mathcal{K})$$

is the \mathcal{K} -reflection of \mathbf{A} .

Proof. It is clear that $\theta(\mathcal{K})$ is an equivalence relation: it is reflexive because \mathbf{F} is generated by A , it is symmetric because $\text{Eq}(\mathcal{K})$ is symmetric, and it is transitive by construction. Now, let $\varphi \in \Omega$ with $\eta(\varphi) = n \geq 1$, and let $(a_1, b_1), \dots, (a_n, b_n) \in \theta(\mathcal{K})$. Without any loss of generality, we may assume that there exist an integer $m \geq 1$ and pairs $(a_{i,j}, a_{i,j+1}) \in \varepsilon(\mathcal{K})$, $i = 1, \dots, n$, $j = 0, \dots, m-1$, such that $a_{i,0} = a_i$ and $a_{i,m} = b_i$.

Then, for every $j = 0, \dots, m-1$, there exist equations $\mathbf{p}_{i,j} \approx \mathbf{q}_{i,j} \in \text{Eq}(\mathcal{K})$ and valuations $v_{i,j}: \mathcal{X} \rightarrow A$ such that $a_{i,j} = \mathbf{p}_{i,j}^{\mathbf{F}}(v_{i,j})$ and $a_{i,j+1} = \mathbf{q}_{i,j}^{\mathbf{F}}(v_{i,j})$, $i = 1, \dots, n$; $\text{Eq}(\mathcal{K})$ being closed under substitution, we may assume that all these equations have pairwise disjoint sets of explicit variables. But then, since $\text{Eq}(\mathcal{K})$ is closed under composition,

$$\varphi(\mathbf{p}_{1,j}, \dots, \mathbf{p}_{n,j}) \approx \varphi(\mathbf{q}_{1,j}, \dots, \mathbf{q}_{n,j}) \in \text{Eq}(\mathcal{K}).$$

Therefore, taking a valuation $v: \mathcal{X} \rightarrow A$ such that its restriction to every $\text{var}(\mathbf{p}_{i,j}) \cup \text{var}(\mathbf{q}_{i,j})$ is equal to the corresponding restriction of $v_{i,j}$, we have

$$\begin{aligned} &(\varphi^{\mathbf{F}}(a_{1,j}, \dots, a_{n,j}), \varphi^{\mathbf{F}}(a_{1,j+1}, \dots, a_{n,j+1})) \\ &= (\varphi(\mathbf{p}_{1,j}, \dots, \mathbf{p}_{n,j})^{\mathbf{F}}(v), \varphi(\mathbf{q}_{1,j}, \dots, \mathbf{q}_{n,j})^{\mathbf{F}}(v)) \in \varepsilon(\mathcal{K}) \end{aligned}$$

for every $j = 0, \dots, m-1$. Then, finally,

$$(\varphi^{\mathbf{F}}(a_1, \dots, a_n), \varphi^{\mathbf{F}}(b_1, \dots, b_m)) = (\varphi^{\mathbf{F}}(a_{1,0}, \dots, a_{n,0}), \varphi^{\mathbf{F}}(a_{1,m}, \dots, a_{n,m})) \in \theta(\mathcal{K}).$$

Thus, $\theta(\mathcal{K})$ is indeed a congruence on \mathbf{F} .

We show now that $\mathbf{F}/_{\theta(\mathcal{K})}$ satisfies $\text{Eq}(\mathcal{K})$. Let $\mathbf{p} \approx \mathbf{q} \in \text{Eq}(\mathcal{K})$, say, with $\text{var}(\mathbf{p}) \cup \text{var}(\mathbf{q}) = \{x_1, \dots, x_n\}$, and let $([c_1]_{\theta}, \dots, [c_n]_{\theta})$ be any n -tuple of elements of $\mathbf{F}/_{\theta(\mathcal{K})}$.

Since \mathbf{F} is generated by A , for every c_i there exists a term \mathbf{p}_i and a valuation $v_i: \mathcal{X} \rightarrow A$ such that $c_i = \mathbf{p}_i^{\mathbf{F}}(v_i)$; as before, we may assume that the sets of variables of the terms \mathbf{p}_i are pairwise disjoint. Then

$$\mathbf{p}(x_1/\mathbf{p}_1, \dots, x_n/\mathbf{p}_n) \approx \mathbf{q}(x_1/\mathbf{p}_1, \dots, x_n/\mathbf{p}_n) \in \text{Eq}(\mathcal{K}).$$

But now, taking a valuation $v: \mathcal{X} \rightarrow A$ such that its restriction to every $\text{var}(\mathbf{p}_i)$ is equal to the corresponding restriction of v_i , we have

$$\begin{aligned} & (\mathbf{p}^{\mathbf{F}}(c_1, \dots, c_n), \mathbf{q}^{\mathbf{F}}(c_1, \dots, c_n)) \\ &= (\mathbf{p}(x_1/\mathbf{p}_1, \dots, x_n/\mathbf{p}_n)^{\mathbf{F}}(v), \mathbf{q}(x_1/\mathbf{p}_1, \dots, x_n/\mathbf{p}_n)^{\mathbf{F}}(v)) \in \varepsilon(\mathcal{K}), \end{aligned}$$

and therefore

$$\mathbf{p}^{\mathbf{F}/\theta}([c_1]_{\theta}, \dots, [c_n]_{\theta}) = \mathbf{q}^{\mathbf{F}/\theta}([c_1]_{\theta}, \dots, [c_n]_{\theta}).$$

This implies that $\mathbf{F}/_{\theta(\mathcal{K})}$ satisfies $\mathbf{p} \approx \mathbf{q}$.

Finally, let $f: \mathbf{F} \rightarrow \mathbf{B}$ be any homomorphism with $\mathbf{B} \in \mathcal{K}$. Then obviously $\theta(\mathcal{K}) \subseteq \ker f$, and therefore there exists a unique homomorphism $\tilde{f}: \mathbf{F}/_{\theta(\mathcal{K})} \rightarrow \mathbf{B}$ such that $f = \tilde{f} \circ \text{nat}_{\theta(\mathcal{K})}$. This means that $\text{nat}_{\theta(\mathcal{K})}: \mathbf{F} \rightarrow \mathbf{F}/_{\theta(\mathcal{K})}$ is a \mathcal{K} -reflection of \mathbf{F} . Since \mathbf{F} is a free completion of \mathbf{A} , $\text{nat}_{\theta(\mathcal{K})} \circ i_{\mathbf{A}}: \mathbf{A} \rightarrow \mathbf{F}/_{\theta(\mathcal{K})}$ is a \mathcal{K} -reflection of \mathbf{A} . \square

The next two corollaries are simple consequences of the previous two lemmas.

Corollary 10. *With the notation and assumptions of Lemma 9, $\mathbf{A} \in S_w(\mathcal{K})$ if and only if $\theta(\mathcal{K}) \cap A^2 = \Delta_A$ (the diagonal on A).*

Corollary 11. *If \mathcal{K} is a total variety and \mathbf{A} satisfies $\text{Eq}(\mathcal{K})$ strongly, then $\mathbf{A} \in S_w(\mathcal{K})$.*

Proof. If \mathbf{A} is a strong model of $\text{Eq}(\mathcal{K})$, then, for every $(a, b) \in \varepsilon(\mathcal{K})$, $b = a$ whenever $a \in A$. This clearly implies that $\theta(\mathcal{K}) \cap A^2 = \Delta_A$, and the last corollary applies. \square

In order to simplify the notation in the statement of the next result, we introduce the concept of *relative prefixes* of two Σ -terms. A term $\mathbf{t} = \mathbf{t}(x_1, \dots, x_n) \in \text{T}_{\Sigma}(\mathcal{X})$

(with explicit variables x_1, \dots, x_n) is a *common suffix* of two terms $\mathbf{p}, \mathbf{q} \in T_\Sigma(\mathcal{X})$ when there exist terms $\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{q}_1, \dots, \mathbf{q}_n$ such that

$$\mathbf{p} = \mathbf{t}(x_1/\mathbf{p}_1, \dots, x_n/\mathbf{p}_n), \quad \mathbf{q} = \mathbf{t}(x_1/\mathbf{q}_1, \dots, x_n/\mathbf{q}_n).$$

In this case, $(\mathbf{p}_1, \mathbf{q}_1), \dots, (\mathbf{p}_n, \mathbf{q}_n)$ are called *pairs of relative prefixes* of the pair (\mathbf{p}, \mathbf{q}) with respect to \mathbf{t} .

Theorem 12. *Let \mathbf{A} be a partial Σ -algebra and \mathcal{K} a variety of total Σ -algebras. Then, $\mathbf{A} \in S_w(\mathcal{K})$ if and only if \mathbf{A} satisfies all quasi-equations of the form*

$$\left(\mathbf{p}_0 \approx \mathbf{q}_0 \wedge \mathbf{q}_k \approx \mathbf{p}_k \wedge \bigwedge_{j=0}^{k-1} \bigwedge_{l=1}^{n_j} \bar{\mathbf{q}}_{j,l} \approx \bar{\mathbf{p}}_{j+1,l} \right) \rightarrow \mathbf{p}_0 \approx \mathbf{q}_k,$$

where, for some sequence of equations

$$\mathbf{p}_0 \approx \mathbf{q}_0, \mathbf{p}_1 \approx \mathbf{q}_1, \dots, \mathbf{p}_k \approx \mathbf{q}_k \in \text{Eq}(\mathcal{K}), \quad k \geq 0,$$

with pairwise disjoint sets of explicit variables (i.e., such that $(\text{var}(\mathbf{p}_i) \cup \text{var}(\mathbf{q}_i)) \cap (\text{var}(\mathbf{p}_j) \cup \text{var}(\mathbf{q}_j)) = \emptyset$ for every $i \neq j$),

$$(\bar{\mathbf{q}}_{j,1}, \bar{\mathbf{p}}_{j+1,1}), \dots, (\bar{\mathbf{q}}_{j,n_j}, \bar{\mathbf{p}}_{j+1,n_j})$$

are the pairs of relative prefixes of $(\mathbf{q}_j, \mathbf{p}_{j+1})$ with respect to some common suffix, for every $j = 0, \dots, k-1$.

Proof. The “only if” implication is straightforward, because every algebra in \mathcal{K} satisfies all quasi-equations of the form given in the statement, and the satisfaction of such quasi-equations is preserved by weak subalgebras.

As for the “if” implication, by Corollary 10 it is enough to prove that if \mathbf{A} satisfies all quasi-equations of the form given in the statement, then $\theta(\mathcal{K}) \cap A^2 = \Delta_A$, where $\theta(\mathcal{K})$ is the congruence on the free completion \mathbf{F} of \mathbf{A} defined in Lemma 9.

So, let $(a, a') \in \theta(\mathcal{K})$ with $a, a' \in A$. Then, there exist

$$(b_0, b_1), (b_1, b_2), \dots, (b_k, b_{k+1}) \in \varepsilon(\mathcal{K}), \quad k \geq 0,$$

such that $a = b_0$ and $a' = b_{k+1}$; without any loss of generality we assume that no other b_i in this sequence of pairs belongs to A .

By the definition of $\varepsilon(\mathcal{K})$, there exist equations $\mathbf{p}_0 \approx \mathbf{q}_0, \mathbf{p}_1 \approx \mathbf{q}_1, \dots, \mathbf{p}_k \approx \mathbf{q}_k$ in $\text{Eq}(\mathcal{K})$ (and, as always, we shall assume their sets of explicit variables to be pairwise disjoint) and valuations $v_0, \dots, v_k: \mathcal{X} \rightarrow A$ such that for every $i = 0, \dots, k$,

$$(b_i, b_{i+1}) = (\mathbf{p}_i^{\mathbf{F}}(v_i), \mathbf{q}_i^{\mathbf{F}}(v_i)).$$

Since \mathbf{A} is an initial segment of \mathbf{F} , $b_0, b_{k+1} \in A$ implies $b_0 = \mathbf{p}_0^{\mathbf{A}}(v_0)$ and $b_{k+1} = \mathbf{q}_k^{\mathbf{A}}(v_k)$. Moreover, from the other Peano axioms satisfied by the free completion \mathbf{F} of \mathbf{A} (see [2] or [6]), we deduce that if $\mathbf{q}_j^{\mathbf{F}}(v_j) = \mathbf{p}_{j+1}^{\mathbf{F}}(v_{j+1}) = b_{j+1} \notin A$, then there is a common suffix \mathbf{t}_j of \mathbf{q}_j and \mathbf{p}_{j+1} such that, if

$$(\bar{\mathbf{q}}_{j,1}, \bar{\mathbf{p}}_{j+1,1}), \dots, (\bar{\mathbf{q}}_{j,n_j}, \bar{\mathbf{p}}_{j+1,n_j})$$

are the pairs of relative prefixes of $(\mathbf{q}_j, \mathbf{p}_{j+1})$ with respect to \mathbf{t}_j , then $\bar{\mathbf{q}}_{j,l}^{\mathbf{A}}(v_j)$ and $\bar{\mathbf{p}}_{j+1,l}^{\mathbf{A}}(v_{j+1})$ are defined and equal for every $l = 1, \dots, n_j$ (*grosso modo*, if $\mathbf{q}_j^{\mathbf{F}}(v_j) = \mathbf{p}_{j+1}^{\mathbf{F}}(v_{j+1}) \notin A$, then both must be equal to the same term, a common suffix \mathbf{t}_j of \mathbf{q}_j and \mathbf{p}_{j+1} , applied in \mathbf{F} to the same elements of A , given by the application in \mathbf{A} of the relative prefixes of \mathbf{q}_j and \mathbf{p}_{j+1} with respect to this common suffix to v_j and v_{j+1} , respectively).

Let now $v: \mathcal{X} \rightarrow A$ be a valuation such that its restriction to every $\text{var}(\mathbf{p}_i) \cup \text{var}(\mathbf{q}_i)$ is equal to the restriction of v_i to this set. We have

$$(\mathbf{A}, v) \models \mathbf{p}_0 \approx \mathbf{p}_0 \wedge \mathbf{q}_k \approx \mathbf{q}_k \wedge \bigwedge_{j=0}^{k-1} \bigwedge_{l=1}^{n_j} \bar{\mathbf{q}}_{j,l} \approx \bar{\mathbf{p}}_{j+1,l}.$$

Then, by hypothesis,

$$(\mathbf{A}, v) \models \mathbf{p}_0 \approx \mathbf{q}_k$$

and therefore

$$a = \mathbf{p}_0^{\mathbf{A}}(v_0) = \mathbf{p}_0^{\mathbf{A}}(v) = \mathbf{q}_k^{\mathbf{A}}(v) = \mathbf{q}_k^{\mathbf{A}}(v_k) = a',$$

as we wanted to prove. □

3. WHEN IS $S_w(\mathcal{K})$ WEAK EQUATIONAL?

In this section we discuss some conditions for a class $S_w(\mathcal{K})$ of all “approximations” of algebras in a total variety \mathcal{K} to be weakly equational. As we have seen in Section 1, all members of $S_w(\mathcal{K})$ are weak models of the equational theory $\text{Eq}(\mathcal{K})$ of \mathcal{K} . However, Example 4 show that, in general, not every weak model of $\text{Eq}(\mathcal{K})$ belongs to $S_w(\mathcal{K})$.

Lemma 13. *Let \mathcal{K} be a variety of total algebras. Then $\text{Mod}_w(\text{Eq}(\mathcal{K})) = HS_w(\mathcal{K})$.*

Proof. By Andr eka-N emeti's general characterization theorem of model classes [1, § 3, Th. 1], $\mathbf{A} \in HS_w(\mathcal{K})$ if and only if

$$\mathbf{A} \models \left(\mathbf{p} \approx \mathbf{q} \wedge \mathbf{q} \approx \mathbf{p} \wedge \bigwedge_{j \in J} \mathbf{p}_j \approx \mathbf{q}_j \right) \rightarrow \mathbf{p} \approx \mathbf{q}$$

for every such quasi-equation satisfied in \mathcal{K} , which in this case means, \mathcal{K} being a total variety, simply that $\mathbf{p} \approx \mathbf{q} \in \text{Eq}(\mathcal{K})$. Since the weak satisfaction of an equation $\mathbf{p} \approx \mathbf{q}$ is nothing but the satisfaction of the quasi-equation

$$(\mathbf{p} \approx \mathbf{p} \wedge \mathbf{q} \approx \mathbf{q}) \rightarrow \mathbf{p} \approx \mathbf{q},$$

it is clear that $\mathbf{A} \in HS_w(\mathcal{K})$ if and only if \mathbf{A} satisfies weakly all equations in $\text{Eq}(\mathcal{K})$. □

This result yields a general necessary and sufficient condition on $S_w(\mathcal{K})$ to be weak equational.

Proposition 14. *Let \mathcal{K} be a variety of total algebras. The following conditions are equivalent.*

- i) $S_w(\mathcal{K})$ is a weak equational class.
- ii) $S_w(\mathcal{K}) = \text{Mod}_w(\text{Eq}(\mathcal{K}))$.
- iii) $S_w(\mathcal{K})$ is closed under closed homomorphic images.

Proof. Notice that if \mathcal{E} is the weak equational theory of $S_w(\mathcal{K})$ then $\mathcal{E} = \text{Eq}(\mathcal{K})$ (since $\mathcal{K} \subseteq S_w(\mathcal{K})$, we have $\mathcal{E} \subseteq \text{Eq}(\mathcal{K})$, and since weak satisfaction is preserved by weak subalgebras, $\text{Eq}(\mathcal{K}) \subseteq \mathcal{E}$). Therefore, $S_w(\mathcal{K})$ is weak equational if and only if $S_w(\mathcal{K}) = \text{Mod}_w(\text{Eq}(\mathcal{K}))$, which proves the equivalence between (i) and (ii). The equivalence between (ii) and (iii) is now a consequence of Lemma 13. □

This proposition, together with Theorem 12, can be used to show that weak hereditary classes of some specific total varieties are not weak equational.

It is known that every total variety \mathcal{K} of monounary algebras is induced by a single equation (see [4]). If \mathcal{K} is regular, then $\mathcal{K} = \text{Mod}_t(\varphi^r(x_1) \approx \varphi^s(x_1))$ for some $r \geq s \geq 0$, and if \mathcal{K} is irregular, then $\mathcal{K} = \text{Mod}_t(\varphi^r(x_1) \approx \varphi^r(x_2))$ for some $r \geq 0$.

Proposition 15. *Let \mathcal{K} be a variety of total monounary algebras. Then*

- i) *if \mathcal{K} is an irregular variety, then $S_w(\mathcal{K})$ is a weak equational class.*
- ii) *if \mathcal{K} is a regular variety induced by the equation $\varphi^r(x_1) \approx \varphi^s(x_1)$ with $r \geq s$, then $S_w(\mathcal{K})$ is weak equational iff $r = s$ or $r = s + 1$.*

Proof. i) Let \mathcal{K} be an irregular variety, i.e. $\mathcal{K} = \text{Mod}_t(\varphi^r(x_1) \approx \varphi^r(x_2))$ for some $r \geq 0$. Then any weak model \mathbf{A} of $\varphi^r(x) \approx \varphi^r(y)$ can be extended to a total algebra $\bar{\mathbf{A}}$ satisfying this equation. To prove it, we distinguish three cases:

- If $r = 0$, then \mathbf{A} is a trivial algebra and \mathcal{K} is the class of all trivial total algebras, and then any inner completion of \mathbf{A} belongs to \mathcal{K} .
 - If $r > 0$ and $(\varphi^{\mathbf{A}})^r(a)$ is defined for some $a \in A$, then add $\varphi^{\bar{\mathbf{A}}}(a') = (\varphi^{\mathbf{A}})^r(a)$ for every $a' \notin \text{dom } \varphi^{\mathbf{A}}$, and the resulting $\bar{\mathbf{A}}$ satisfies $\varphi^r(x_1) \approx \varphi^r(x_2)$.
 - If $r > 0$ and $\text{dom}(\varphi^{\mathbf{A}})^r = \emptyset$, then take any $a_0 \notin \text{dom } \varphi^{\mathbf{A}}$ and add $\varphi^{\bar{\mathbf{A}}}(a') = a_0$ for every $a' \notin \text{dom } \varphi^{\mathbf{A}}$ (including a_0 itself). Again, the resulting $\bar{\mathbf{A}}$ satisfies $\varphi^r(x_1) \approx \varphi^r(x_2)$.
- ii) Assume now \mathcal{K} is a regular variety and $\mathcal{K} = \text{Mod}_t(\varphi^r(x_1) \approx \varphi^s(x_1))$ for some $r \geq s \geq 0$. To prove the claim we again consider three cases.

- Assume first that $r > s + 1$. Consider the relative subalgebra \mathbf{A} of the Σ -term algebra $\mathbf{T}_\Sigma(\{x, y\})$ supported on the initial segment determined by $\{\varphi^{r-1}(x), \varphi^{r-1}(y)\}$; it is easily seen to be a weak subalgebra of the \mathcal{K} -free algebra generated by $\{x, y\}$. Let θ denote the equivalence relation on A identifying $\varphi^{r-1}(x)$ and $\varphi^{r-1}(y)$. It is a closed congruence on \mathbf{A} , and thus $\mathbf{A}/\theta \in HS_w(\mathcal{K})$. But $\mathbf{A}/\theta \notin S_w(\mathcal{K})$, because it does not satisfy the quasi-equation

$$\begin{aligned} (\varphi^s(x_1) \approx \varphi^s(x_1) \wedge \varphi^s(x_2) \approx \varphi^s(x_2) \wedge \varphi^{r-1}(x_1) \approx \varphi^{r-1}(x_2)) \\ \rightarrow \varphi^s(x_1) \approx \varphi^s(x_2), \end{aligned}$$

obtained from the equations $\varphi^s(x_1) \approx \varphi^r(x_1)$ and $\varphi^r(x_2) \approx \varphi^s(x_2)$ as in Theorem 12, with respect to any valuation $v: \mathcal{X} \rightarrow A/\theta$ such that $v(x_1) = [x]_\theta$ and $v(x_2) = [y]_\theta$.

So, if $r > s + 1$, $S_w(\mathcal{K})$ is not closed under closed homomorphic images, and therefore it is not weak equational.

- If $r = s$, then \mathcal{K} is the class of all Σ -algebras, and therefore $S_w(\mathcal{K})$ is the (weak equational) class of all partial Σ -algebras.
- Finally, if $r = s + 1$, then completing any weak model \mathbf{A} of $\varphi^{s+1}(x) \approx \varphi^s(x)$ by adding a φ -loop to every point where $\varphi^{\mathbf{A}}$ is not defined, yields a total Σ -algebra satisfying $\varphi^{s+1}(x_1) \approx \varphi^s(x_1)$, i.e. an algebra in \mathcal{K} . So, in this case every weak model of $\text{Eq}(\mathcal{K})$ is in $S_w(\mathcal{K})$, which implies that the latter is a weak equational class.

This completes the proof of the proposition. □

In a subsequent paper we plan to characterize those varieties of total algebras of an arbitrary unary type which have a weak equational weak hereditary class.

Example 16. Let Σ be a type of algebras with only two operation symbols $*$ and $+$, both binary, and let \mathcal{K} be the total variety of Σ -algebras defined by the absorption equation $x_1 + (x_1 * x_2) \approx x_1$. The weak hereditary class $S_w(\mathcal{K})$ is weak equational. Indeed, for every partial Σ -algebra \mathbf{A} , let $\bar{\mathbf{A}}$ be its completion defined in the following way:

- $\bar{A} = A \cup \{a_0\}$ for some $a_0 \notin A$.
- For every $x, y \in \bar{A}$, if $x * y$ is not defined in \mathbf{A} , then $x * y = a_0$ in $\bar{\mathbf{A}}$.
- For every $x, y \in \bar{A}$, if $x + y$ is not defined in \mathbf{A} , then $x + y = x$ in $\bar{\mathbf{A}}$.

It is straightforward to check that if \mathbf{A} satisfies weakly $x_1 + (x_1 * x_2) \approx x_1$, then $\bar{\mathbf{A}} \in \mathcal{K}$ and hence $\mathbf{A} \in S_w(\mathcal{K})$.

Let now \mathcal{K}' be the total variety of Σ -algebras defined by *both* absorption equations

$$x_1 + (x_1 * x_2) \approx x_1, \quad x_1 * (x_1 + x_2) \approx x_1.$$

Then, $S_w(\mathcal{K}')$ is not weak equational. Indeed, let \mathbf{A} be a partial Σ -algebra with carrier $A = \{a, b\}$, operation $+$ totally defined by

$$a + a = a + b = b, \quad b + a = b + b = a,$$

and operation $*$ discrete. It is clear then that \mathbf{A} is a weak model of $\text{Eq}(\mathcal{K}')$. But no completion of \mathbf{A} belongs to \mathcal{K}' , because in any extension of \mathbf{A} satisfying the equation $x_1 * (x_1 + x_2) \approx x_1$ the equality $a * b = a * (a + b) = a$ is true, and then this extension does not satisfy the other absorption equation, since then $a + (a * b) = a + a = b$.

In the next example we will use the following straightforward result.

Proposition 17. *Let Σ be a type of algebras and $\Sigma_0 \subseteq \Sigma$ a subtype of it. Let \mathcal{K} be a variety of total Σ -algebras,*

$$\mathcal{E}_0 = \{\mathbf{p} \approx \mathbf{q} \in \text{Eq}(\mathcal{K}) : \mathbf{p}, \mathbf{q} \text{ } \Sigma_0\text{-terms}\},$$

and $\mathcal{K}_0 = \text{Mod}_t(\mathcal{E}_0)$ the variety of total Σ_0 -algebras defined by \mathcal{E}_0 . If $S_w(\mathcal{K})$ is weak equational, then $S_w(\mathcal{K}_0)$ is weak equational.

Proof. Notice that, \mathcal{E}_0 being closed under Birkhoff's equational logic rules, $\mathcal{E}_0 = \text{Eq}(\mathcal{K}_0)$. Let now \mathbf{A} be a weak model of \mathcal{E}_0 , and let $\hat{\mathbf{A}}$ be the partial Σ -algebra whose Σ_0 -reduct is \mathbf{A} and where all realizations of operation symbols not belonging to Σ_0 are discrete. This Σ -algebra $\hat{\mathbf{A}}$ is a weak model of $\text{Eq}(\mathcal{K})$ and therefore, by the assumption, it is a weak subalgebra of a total Σ -algebra \mathbf{B} in \mathcal{K} . Then, \mathbf{A} is a weak subalgebra of the Σ_0 -reduct of \mathbf{B} , which belongs to \mathcal{K}_0 . □

Example 18. Let Σ be a type of algebras with only one operation symbol $*$, which is binary, and let \mathcal{K} be the class of all semigroups: i.e., the total variety of Σ -algebras defined by the equation $x_1 * (x_2 * x_3) \approx (x_1 * x_2) * x_3$. Then $S_w(\mathcal{K})$ is not weak equational.

Consider the relative subalgebra \mathbf{A} of the Σ -term algebra $\mathbf{T}_\Sigma(\{x_1, x_2, x_3, y_2, y_3\})$ supported on the initial segment A generated by

$$(x_1 * x_2) * x_3, (x_1 * y_2) * y_3, x_2 * x_3, y_2 * y_3,$$

and let θ be the equivalence relation on A that identifies $x_2 * x_3$ with $y_2 * y_3$. It is straightforward to prove that \mathbf{A} is a weak subalgebra of the free semigroup generated by $\{x_1, x_2, x_3, y_2, y_3\}$, and that θ is a closed congruence on \mathbf{A} . But $\mathbf{A}/\theta \notin S_w(\mathcal{K})$, because it does not satisfy the quasi-equation

$$\begin{aligned} ((x_1 * x_2) * x_3 \approx (x_1 * x_2) * x_3 \wedge (x_4 * x_5) * x_6 \approx (x_4 * x_5) * x_6 \wedge x_2 * x_3 \approx x_5 * x_6) \\ \rightarrow (x_1 * x_2) * x_3 \approx (x_4 * x_5) * x_6 \end{aligned}$$

w.r.t. any valuation $v: \mathcal{X} \rightarrow \mathbf{A}/\theta$ satisfying

$$x_1, x_4 \mapsto [x_1]_\theta, x_2 \mapsto [x_2]_\theta, x_3 \mapsto [x_3]_\theta, x_5 \mapsto [y_2]_\theta, x_6 \mapsto [y_3]_\theta.$$

Therefore, $S_w(\mathcal{K})$ is not closed under closed homomorphic images and thus it is not weak equational.

The last proposition also implies that the weak hereditary classes of the varieties of monoids and groups are not weak equational. To cover the cases of abelian groups, rings, lattices, etc., we must prove that the weak hereditary class of the variety of abelian semigroups

$$\mathcal{K}' = \text{Mod}_t(x_1 * (x_2 * x_3) \approx (x_1 * x_2) * x_3, x_1 * x_2 \approx x_2 * x_1)$$

is not weak equational either. To do this, we can use an argument similar to the previous one.

Let $\mathcal{K}_{ab} = \text{Mod}_t(x_1 * x_2 \approx x_2 * x_1)$: its weak hereditary class is clearly weak equational. Let \mathbf{F}_{ab} be the \mathcal{K}_{ab} -free algebra generated by the set $G = \{x_1, x_2, x_3, y_2, y_3\}$, let \mathbf{A} be the relative subalgebra of \mathbf{F}_{ab} supported on the set A consisting of G and the elements

$$x_1 * x_2, x_1 * y_2, x_2 * x_3, y_2 * y_3, (x_1 * x_2) * x_3, (x_1 * y_2) * y_3,$$

and let θ be the equivalence relation on A that identifies $x_2 * x_3$ with $y_2 * y_3$. It is straightforward to prove that \mathbf{A} is a weak subalgebra of the free abelian semigroup

generated by $\{x_1, x_2, x_3, y_2, y_3\}$, and that θ is a closed congruence on \mathbf{A} . But $\mathbf{A}/\theta \notin S_w(\mathcal{K}')$, because it does not satisfy the quasi-equation

$$\begin{aligned} ((x_1 * x_2) * x_3 \approx (x_1 * x_2) * x_3 \wedge (x_4 * x_5) * x_6 \approx (x_4 * x_5) * x_6 \wedge x_2 * x_3 \approx x_5 * x_6) \\ \rightarrow (x_1 * x_2) * x_3 \approx (x_4 * x_5) * x_6 \end{aligned}$$

with respect to any valuation $v: \mathcal{X} \rightarrow \mathbf{A}/\theta$ such that

$$x_1, x_4 \mapsto [x_1]_\theta, \quad x_2 \mapsto [x_2]_\theta, \quad x_3 \mapsto [x_3]_\theta, \quad x_5 \mapsto [y_2]_\theta, \quad x_6 \mapsto [y_3]_\theta.$$

Therefore, $S_w(\mathcal{K}')$ is not closed under closed homomorphic images and hence it is not weak equational.

To close this paper, let us point out the following straightforward consequence of Corollary 19.

Corollary 19. *Let \mathcal{K} be a total variety. Then, with the notations of Lemma 9, $S_w(\mathcal{K})$ is a weak equational class if and only if for every partial algebra \mathbf{A} , if $\varepsilon(\mathcal{K}) \cap A^2 = \Delta_A$, then $\theta(\mathcal{K}) \cap A^2 = \Delta_A$.*

This result suggests that some sort of congruence extension property may be relevant in the characterization of those total varieties whose weak hereditary class is weak equational. The investigation of other congruence extension properties for partial algebras has found applications in other, quite different contexts [5].

References

- [1] *H. Andr eka, I. N emeti*: Generalization of the concept of variety and quasivariety to partial algebras through category theory. *Dissertationes Math. (Rozpr. Matem.) 204*, (1983). [Zbl 0518.08007](#)
- [2] *P. Burmeister*: A Model Theoretic Approach to Partial Algebras. *Math. Research 32*. Akademie-Verlag, Berlin, 1986. [Zbl 0598.08004](#)
- [3] *G. Gr atzer, E. T. Schmidt*: Characterizations of congruence lattices of abstract algebras. *Acta Sci. Math. (Szeged) 24* (1963), 34–59. [Zbl 0117.26101](#)
- [4] *D. Jakubikova-Studenovska*: On completions of partial monounary algebras. *Czechoslovak Math. J. 38* (1988), 256–268. [Zbl 0659.08003](#)
- [5] *M. Llabr es, F. Rossell o*: Pushout complements for arbitrary partial algebras. In: *Proc. 6th International Workshop on Theory and Application of Graph Transformation TAGT'98. Lect. Notes in Comp. Sc. 1764*. 2000, pp. 131–144. [Zbl 0957.08004](#)
- [6] *J. Schmidt*: A homomorphism theorem for partial algebras. *Colloq. Math. 21* (1970), 5–21. [Zbl 0199.32402](#)
- [7] *B. Staruch, B. Staruch*: Strong regular varieties of partial algebras. *Alg. Univ. 31* (1994), 157–176. [Zbl 0805.08002](#)

- [8] *R. Szymański*: Decidability of weak equational theories. Czechoslovak Math. J. 46 (1996), 629–664. [Zbl 0879.08005](#)

Authors' addresses: Wiktor Bartol, Inst. Matematyki, Uniw. Warszawski, ul. Banacha 2, 02-097 Warszawa, Poland, e-mail: bartol@mimuw.edu.pl; Francesc Rosselló, Dept. Matemàtiques i Informàtica, Univ. Illes Balears, 07122 Palma de Mallorca, Spain, e-mail: rossello@uib.es.