

Florian Luca

On the Euler function of repdigits

Czechoslovak Mathematical Journal, Vol. 58 (2008), No. 1, 51–59

Persistent URL: <http://dml.cz/dmlcz/128245>

Terms of use:

© Institute of Mathematics AS CR, 2008

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE EULER FUNCTION OF REPDIGITS

FLORIAN LUCA, Morelia

(Received November 24, 2005)

Dedicated to William D. Banks on his $\sqrt{\varphi(2005)}$ th birthday.

Abstract. For a positive integer n we write $\varphi(n)$ for the Euler function of n . In this note, we show that if $b > 1$ is a fixed positive integer, then the equation

$$\varphi\left(x\frac{b^n-1}{b-1}\right) = y\frac{b^m-1}{b-1}, \quad \text{where } x, y \in \{1, \dots, b-1\},$$

has only finitely many positive integer solutions (x, y, m, n) .

Keywords: Euler function, prime, divisor

MSC 2000: 11A25

1. INTRODUCTION

For a positive integer n we write $\varphi(n)$ for the Euler function of n . In this paper, we prove the following result.

Theorem 1.1. *If $b > 1$ is given, then the equation*

$$(1) \quad \varphi\left(x\frac{b^n-1}{b-1}\right) = y\frac{b^m-1}{b-1},$$

with $x, y \in \{1, \dots, b-1\}$ has only finitely many positive integer solutions (x, y, m, n) .

Some equations of a similar flavor have been treated in [3], [4], [5] and [6].

We use the Vinogradov symbols \ll and \gg , and the Landau symbol O with their regular meanings. The constants implied by them may depend on our parameter b . We use p, q and P with or without subscripts to denote prime numbers. For a positive

real number x we use $\log x$ for the maximum between 2 and the natural logarithm of x . Note that with this convention, the function \log is sub-multiplicative; i.e., $\log(xy) \leq \log x \log y$ holds for all positive real numbers x and y . For a positive integer n , we write $P(n)$, $p(n)$, $\omega(n)$, $\Omega(n)$ and $\tau(n)$ for the largest prime factor of n , smallest prime factor of n , the number of distinct prime factors of n , the number of prime power divisors (> 1) of n , and the total number of divisors of n , respectively. We put $u_n = (b^n - 1)/(b - 1)$. Finally, we use c_0, c_1, \dots for positive constants depending on b which are labeled increasingly throughout the paper.

2. THE PROOF

Since b is fixed, and (x, y) can take only $(b - 1)^2$ values, we may assume that both x and y are fixed. Let $N = x(b^n - 1)/(b - 1)$. If $m > n$, then

$$\varphi(N) = y \frac{b^m - 1}{b - 1} \geq \frac{b^{n+1} - 1}{b - 1} > b^n - 1 \geq N,$$

which is a contradiction. If $m = n$, then $\varphi(N)/N = y/x$. Since $P(N)$ divides the denominator of the rational number $\varphi(N)/N$ in reduced form, it follows that $P(N) \leq b - 1$. In particular, $P(u_n) \leq b - 1$. Since for $n > 6$, u_n always has a *primitive divisor*, which, in particular, is a prime congruent to 1 modulo n , we get that $n \leq \max\{6, b - 2\}$ (see [1] and [2] for the existence and properties of primitive divisors).

From now on, we assume that $n > m$. We will first show that $n - m$ is bounded. Let $k = \gcd(m, n)$. Then k divides $\lambda = n - m$, therefore

$$(2) \quad b^k \leq b^\lambda \ll \frac{N}{\varphi(N)} = \prod_{P|N} \left(1 + \frac{1}{P-1}\right) \ll \prod_{\substack{P|N \\ P > b}} \left(1 + \frac{1}{P-1}\right).$$

Let $P \mid N$ such that $P > b$. Then P does not divide x and there exists a divisor l_P of n minimal with the property that $P \mid u_{l_P}$. The number l_P is called *the order of apparition of P in the sequence $(u_n)_{n \geq 1}$* and P is certainly primitive for u_{l_P} . Furthermore, $P \equiv 1 \pmod{l_P}$. We now fix $d \mid n$ and consider

$$(3) \quad S_d = \sum_{l_P=d} \frac{1}{P} \quad \text{and} \quad \omega_d = \#\{P: l_P = d\}.$$

Clearly,

$$b^d \gg u_d \geq \prod_{l_P=d} P \geq d^{\omega_d},$$

giving

$$(4) \quad \omega_d \ll \frac{d}{\log d}.$$

Using estimate (4), we can estimate the sum \mathcal{S}_d defined in (3) as follows

$$(5) \quad \mathcal{S}_d \leq \sum_{\substack{l_P=d \\ P < d^2}} \frac{1}{P} + \sum_{\substack{l_P=d \\ P \geq d^2}} \frac{1}{P} \ll \sum_{\substack{P \equiv 1 \pmod{d} \\ P \leq d^2}} \frac{1}{P} + \frac{\omega_d}{d^2} \ll \frac{\log \log d}{\varphi(d)},$$

where in the above inequalities (5) we used the estimate (4), together with the Brun-Titchmarsh Theorem which asserts that the estimate

$$\sum_{\substack{p \equiv a \pmod{b} \\ p < t}} \frac{1}{p} \ll \frac{\log \log t}{p}$$

holds for all coprime integers $1 \leq a \leq b$ and all positive real numbers t (see, for example, Lemma 6.3 in [7] or Theorem 1 in [8]). Let c_0 be an upper bound for the constant implied by the Vinogradov symbol appearing in (5), and assume that $c_0 > 1$.

Taking logarithms in the inequality (2) and using the inequality $1 + t < e^t$ which is valid for all positive real numbers t , we get

$$(6) \quad k \leq \lambda \leq O(1) + \sum_{\substack{P|u_n \\ P > b}} \frac{1}{P-1} \leq \sum_{\substack{d|n \\ d > 1}} \mathcal{S}_d + O\left(1 + \sum_{P \geq 2} \frac{1}{P^2}\right) \\ \leq c_0 \sum_{\substack{d|n \\ d > 1}} \frac{\log \log d}{\varphi(d)} + O(1).$$

Since the function $\log \log(\cdot)$ is sub-multiplicative, it follows that the function $c_0 \log \log n / \varphi(n)$ satisfies

$$\frac{c_0 \log \log(ab)}{\varphi(ab)} \leq \frac{c_0 \log \log a}{\varphi(a)} \cdot \frac{c_0 \log \log b}{\varphi(b)}, \quad \text{whenever } \gcd(a, b) = 1.$$

Hence, writing $n = p_1^{\nu_1} \dots p_s^{\nu_s}$, with $p(n) = p_1 < \dots < p_s = P(n)$, we have

$$\sum_{\substack{d|n \\ d > 1}} \frac{c_0 \log \log d}{\varphi(d)} \leq \prod_{i=1}^s \left(1 + \sum_{\nu=1}^{\nu_i} \frac{c_0 \log \log(p_i^\nu)}{p_i^{\nu-1}(p_i-1)}\right) - 1.$$

Since obviously

$$\sum_{\nu \geq 1} \frac{\log \log(p^\nu)}{p^{\nu-1}(p-1)} \ll \frac{\log \log p}{p},$$

we get that there exists a positive constant c_1 such that

$$(7) \quad \sum_{\substack{d|n \\ d>1}} \frac{c_0 \log \log d}{\varphi(d)} \leq \prod_{p|n} \left(1 + \frac{c_1 \log \log p}{p}\right) - 1.$$

Combining (7) with the estimate (6), we get

$$k \leq \lambda \ll \prod_{p|n} \left(1 + \frac{c_1 \log \log p}{p}\right),$$

and therefore

$$k \leq \lambda \ll \prod_{p|n} \left(1 + \frac{c_1 \log \log p}{p}\right) \ll \exp\left(c_1 \sum_{p|n} \frac{\log \log p}{p}\right),$$

which, after taking logarithms, gives

$$(8) \quad \log k \leq \log \lambda \ll 1 + \sum_{p|n} \frac{\log \log p}{p}.$$

We now bound the sum

$$\mathcal{T} = \sum_{p|n} \frac{\log \log p}{p}.$$

Assume first that $p \mid k$. Clearly, $\omega(k) = O(\log k / \log \log k)$, therefore, by the Prime Number Theorem, there exists an absolute constant c_2 , such that

$$(9) \quad \begin{aligned} \mathcal{T}_1 &= \sum_{p|k} \frac{\log \log p}{p} \ll \sum_{q \leq c_2 \log k} \frac{\log \log q}{q} \\ &\ll \log \log(c_2 \log k) \sum_{q \leq c_2 \log k} \frac{1}{q} \ll (\log \log \log k)^2. \end{aligned}$$

Assume now that $p \nmid k$. Then $p \nmid m$. Thus,

$$(10) \quad \text{ord}_p(bu_m) = \text{ord}_p(b) + \text{ord}_p(u_m) \ll 1 + \text{ord}_p(l_p) \ll 1 + \frac{p}{\log p}.$$

Here, for a positive integer n and a prime p we use $\text{ord}_p(n)$ for the exact order at which p divides n , together with the well-known facts that $p \mid u_m$ if and only if $l_p \mid m$, that $l_p \mid p - 1$, and that if $p \nmid m$, then

$$\text{ord}_p(u_m) = \text{ord}_p(u_{l_p}) \leq \text{ord}_p(b^{p-1} - 1) \leq \frac{\log(b^{p-1})}{\log p} \ll \frac{p}{\log p}.$$

Now let t be any positive integer and let us count the contribution to the sum \mathcal{T} from primes in $\mathcal{I}_t = [2^t, 2^{t+1}]$. Let p be a prime in \mathcal{I}_t and let n_t be the number of prime factors of n in \mathcal{I}_t which do not divide k . Then n has at least 2^{n_t-1} distinct divisors which are multiples of p . For each one of these divisors d except $O(1)$ of them (actually, for each one of these divisors except, possibly, the values less than or equal to 6), u_d has a primitive divisor; i.e., a prime $q \mid u_d$ such that $q \nmid u_{d'}$ for any $d' < d$, and $q \equiv 1 \pmod{d}$. This argument shows that u_n has at least $2^{n_t-1} - 6$ distinct divisors congruent to 1 modulo p , giving $\text{ord}_p(\varphi(N)) \geq 2^{n_t-1} - 6$. Combining this argument with the estimate (10), we get

$$2^{n_t-1} \ll 1 + \frac{p}{\log p} \ll 1 + \frac{2^t}{t},$$

giving $n_t \ll t$. Thus,

$$(11) \quad \mathcal{T}_2 = \sum_{\substack{p \mid n \\ p \nmid k}} \frac{\log \log p}{p} \ll \sum_{t \geq 1} \frac{n_t \log \log(2^{t+1})}{2^t} \ll \sum_{t \geq 1} \frac{t \log t}{2^t} \ll 1.$$

Inserting the estimates (9) and (11) into the estimate (8), we get

$$\log k \leq \log \lambda \ll 1 + (\log \log \log k)^3,$$

leading to the conclusion that k (hence, also λ) is bounded. We may therefore assume that both k and λ are fixed. Furthermore, by replacing now b by b^k , x by $x(b^k - 1)/(b - 1)$, and y by $y(b^k - 1)/(b - 1)$, we may assume that m and n are coprime; i.e., that $k = 1$.

To finish, we shall show in what follows first that $p_1 = p(n)$ is bounded, then that $s = \Omega(n)$ is bounded, and finally that n itself is bounded.

Assume that $p(n) = p_1$ can get arbitrarily large. In particular, we may assume that $p_1 > \min\{6, b\}$. Then the smallest prime factor of u_n is congruent to 1 modulo p_i for some $i \geq 1$, therefore it is $\geq 2p_1 + 1 > b > x$. Hence, the equation (1) can be written as

$$\varphi(u_n) = \frac{y}{\varphi(x)} u_m,$$

therefore

$$(12) \quad \frac{\varphi(u_n)}{u_n} = \frac{yu_m}{\varphi(x)u_n} \leq \frac{yu_{n-1}}{\varphi(x)u_n} \leq \frac{(b-1)(b^{n-1}-1)}{b^n-1}.$$

The limit of the expression appearing on the right hand side of the above inequality (12) when $n \rightarrow \infty$ is $1 - 1/b$. Hence, if $n > c_3$, then the right-hand side of the above inequality is $\leq c_4 = 1 - 1/(2b)$. Thus,

$$c_4^{-1} \leq \frac{u_n}{\varphi(u_n)} = \prod_{P|n} \left(1 + \frac{1}{P-1}\right) \leq \exp \left(\sum_{\substack{d|n \\ d>1}} \mathcal{S}_d + O \left(\sum_{p \geq p_1} \frac{1}{p^2} \right) \right),$$

giving

$$c_5 \leq \sum_{d|n} \mathcal{S}_d + O \left(\frac{1}{p_1} \right),$$

where $c_5 = \log(c_4^{-1}) > 0$. Thus, if c_6 is the constant implied by the above Landau symbol, and if $p_1 > c_7 = 2c_6c_5^{-1}$, then we get

$$1 \ll \sum_{\substack{d|n \\ d>1}} \mathcal{S}_d,$$

where the constant implied in the above Vinogradov symbol is $c_8 = 2c_5^{-1}$. Using the estimates (5) and (7), we get that

$$(13) \quad 1 \ll \sum_{\substack{d|n \\ d>1}} \mathcal{S}_d \leq \sum_{\substack{d|n \\ d>1}} \frac{c_0 \log \log d}{\varphi(d)} \leq \prod_{i=1}^s \left(1 + \frac{c_1 \log \log p_i}{p_i}\right) - 1.$$

The same argument employed to bound the number of prime factors of n in the interval \mathcal{I}_t which do not divide k , shows that n has at least $2^{s-1} - 6$ prime factors which are congruent to 1 modulo p_1 . Hence, $\text{ord}_{p_1}(\varphi(N)) \geq 2^{s-1} - 6$, while by the inequality (10), the number $\text{ord}_{p_1}(\varphi(N))$ cannot exceed $\text{ord}_{p_1}(b^{p_1-1} - 1) = O(p_1/\log p_1)$. This shows that $s = \omega(n) \leq c_9 \log p_1$. Hence,

$$(14) \quad \prod_{i=1}^s \left(1 + \frac{c_1 \log \log p_i}{p_i}\right) - 1 \leq \left(1 + \frac{c_1 \log \log p_1}{p_1}\right)^{c_9 \log p_1} - 1 \\ \leq \exp \left(c_{10} \frac{\log p_1 \log \log p_1}{p_1} \right) - 1.$$

Here, $c_{10} = c_1 c_9$. Since the function $(\log p_1 \log \log p_1)/p_1$ is bounded, we conclude that there exists a constant c_{11} such that

$$(15) \quad \exp\left(c_{10} \frac{\log p_1 \log \log p_1}{p_1}\right) - 1 \leq c_{11} \frac{\log p_1 \log \log p_1}{p_1}.$$

The combination of the inequalities (13), (14) and (15) leads to the conclusion that

$$p_1 \ll \log p_1 \log \log p_1,$$

which shows that p_1 is bounded. Now n has at least $\tau(n/p_1) - 6$ divisors which are multiples of p_1 and which are > 6 . For each such divisor, u_n has a primitive divisor which is congruent to 1 modulo p_1 , which shows that $\text{ord}_{p_1}(\varphi(N)) \geq \tau(n/p_1) - 6$. Since by the estimate (10) this p_1 -adic order is $\ll 1 + p_1/\log p_1 \ll 1$, we get that $\tau(n/p_1) \ll 1$, therefore $\tau(n) \ll 1$. In particular, $\Omega(n)$ is bounded.

To finish the proof, it suffices to show that for each $i \leq s$, p_i is bounded. We proceed by induction on i , the case $i = 1$ being obvious. Fix s , $1 \leq i \leq s - 1$, and assume inductively that p_i is bounded. Since the numbers ν_j for $j = 1, \dots, s$ are also bounded, we may assume that the first i distinct primes as well as their multiplicities are all fixed. Write $n_1 = \prod_{j=1}^i p_j^{\nu_j}$. Then $p_{i+1} = p(n/n_1)$. Assume that p_{i+1} can get arbitrarily large. Suppose, in particular, that it is larger than $\min\{6, b^{n_1} - 1\}$. Then writing $u_n = (b^{n_1} - 1)/(b - 1) \cdot (b^n - 1)/(b^{n_1} - 1)$, and observing that every prime factor of $(b^n - 1)/(b^{n_1} - 1)$ is congruent to 1 modulo p_j for some $j \geq i + 1$; hence, larger than $b^{n_1} - 1$, we get that

$$yu_m = \varphi(N) = \varphi(xu_{n_1})\varphi\left(\frac{b^n - 1}{b^{n_1} - 1}\right),$$

so writing $N_1 = (b^n - 1)/(b^{n_1} - 1)$, we get

$$\frac{\varphi(N_1)}{N_1} = \frac{yu_m}{\varphi(xu_{n_1})N_1} = \frac{y(b^n - 1)(b^{n_1} - 1)}{(b - 1)\varphi(xu_{n_1})(b^n - 1)}.$$

The left-hand side of the above-equality is < 1 , while the right hand side tends to (assuming that $n \rightarrow \infty$)

$$L = \frac{y(b^{n_1} - 1)b^{-\lambda}}{(b - 1)\varphi(xu_{n_1})}.$$

Note that the above number is < 1 , for if it were equal to 1, we would then get the equation

$$(b - 1)\varphi(xu_{n_1}) = \frac{y(b^{n_1} - 1)}{b^\lambda},$$

which is impossible since its left-hand side is an integer and its right-hand side is not. Hence, $L < 1$. Thus, choosing c_{12} to be some constant in the interval $(L, 1)$, we get that

$$(16) \quad c_{12}^{-1} \leq \frac{N_1}{\varphi(N_1)} = \prod_{P|N_1} \left(1 + \frac{1}{P-1}\right).$$

It is clear that $P | N_1$ if and only if $P | u_n, n_1 | l_P$ and $l_P > n_1$. Hence, using again the fact that $1 + t < e^t$ for all $t > 0$, and the estimate (5), we get

$$\prod_{P|N_1} \left(1 + \frac{1}{P-1}\right) \leq \exp \left(c_0 \sum_{\substack{n_1|d \\ d > n_1}} \frac{\log \log d}{\varphi(d)} + O \left(\sum_{P \geq p_{i+1}} \frac{1}{P^2} \right) \right),$$

which together with the estimates (16) and (7) leads to

$$\begin{aligned} c_{13} &\leq c_0 \sum_{\substack{n_1|d \\ d > n_1}} \frac{\log \log d}{\varphi(d)} + O \left(\frac{1}{p_{i+1}} \right) \\ &\leq c_0 \mathcal{S}_{n_1} \left(\prod_{j=i+1}^s \left(1 + \frac{c_1 \log \log p_j}{p_j} \right) - 1 \right) + O \left(\frac{1}{p_{i+1}} \right), \end{aligned}$$

where $c_{13} = \log(c_{12}^{-1}) > 0$. Writing c_{14} for an upper bound for $c_0 \mathcal{S}_{n_1}$, and c_{15} for the constant implied by the above Landau symbol, we get that if $p_{i+1} > 2c_{15}c_{13}^{-1}$, then

$$1 \ll \prod_{j=i+1}^s \left(1 + \frac{c_1 \log \log p_j}{p_j} \right) - 1 \leq \left(1 + \frac{c_1 \log \log p_{i+1}}{p_{i+1}} \right)^s - 1,$$

where the constant implied in the above Vinogradov symbol is $c_{16} = 2c_{14}c_{13}^{-1}$. The above inequality certainly implies that

$$1 \ll \frac{s \log \log p_{i+1}}{p_{i+1}},$$

which leads to $p_{i+1} \ll 1$, thus completing the induction and finishing the proof of the theorem.

3. COMMENTS AND REMARKS

If one replaces the condition that x and y belong to $\{1, \dots, b-1\}$ with the weaker condition that x and y are fixed (or bounded), then it is perhaps not true that the equation (1) has only finitely many such solutions (m, n) . For example, taking $b = 2$, $x = 1$, $y = 2$, we note that the equation (1) is always satisfied when $m = n - 1$ and $2^n - 1$ is prime. Of course, we do not know that there are infinitely many *Mersenne primes*; i.e., primes of the form $2^n - 1$, but the general belief is that this is indeed so. Note further that when $m = n = 1$, then the equation (1) is trivially satisfied with $y = \varphi(x)$. It would be interesting to study the nontrivial solutions of the equation (1) in all five variables (x, y, b, m, n) ; i.e., where the base b is also variable. We conjecture that there exists an absolute constant n_0 such that all such solutions have $n \leq n_0$. We leave this conjecture as an open problem for the reader.

Acknowledgements. This paper was written during a very enjoyable visit of the author to the Laboratoire de Mathématiques Nicolas Oresme of the University of Caen; he wishes to express his thanks to that institution for its hospitality and support. This work was also supported in part by research grants SEP-CONACYT 46755, PAPIIT IN104505 and a Guggenheim Fellowship.

References

- [1] *Yu. Bilu, G. Hanrot and P. M. Voutier*: Existence of primitive divisors of Lucas and Lehmer numbers (with an appendix by M. Mignotte). *J. Reine Angew. Math.* *539* (2001), 75–122. zbl
- [2] *R. D. Carmichael*: On the numerical factors of the arithmetic forms $\alpha^n \pm \beta^n$. *Ann. Math.* *15* (1913), 30–70. zbl
- [3] *F. Luca*: On the equation $\varphi(|x^m + y^m|) = |x^n + y^n|$. *Indian J. Pure Appl. Math.* *30* (1999), 183–197. zbl
- [4] *F. Luca*: On the equation $\varphi(x^m - y^m) = x^n + y^n$. *Irish Math. Soc. Bull.* *40* (1998), 46–55. zbl
- [5] *F. Luca*: Euler indicators of binary recurrent sequences. *Collect. Math.* *53* (2002), 133–156. zbl
- [6] *F. Luca*: Problem 10626. *Amer. Math. Monthly* *104* (1997), 871.
- [7] *K. K. Norton*: On the number of restricted prime factors of an integer I. *Illinois J. Math.* *20* (1976), 681–705 Zbl 0329.10035.
- [8] *C. Pomerance*: On the distribution of amicable numbers. *J. Reine Angew. Math.* *293/294* (1977), 217–222. zbl

Author's address: Florian Luca, Instituto de Matemáticas, Universidad Nacional Autónoma de México, C.P. 58089, Morelia, Michoacán, México, e-mail: fluca@matmor.unam.mx.