

František Machala

Contexts and sublattices of concept lattices

Czechoslovak Mathematical Journal, Vol. 44 (1994), No. 4, 757–762

Persistent URL: <http://dml.cz/dmlcz/128487>

Terms of use:

© Institute of Mathematics AS CR, 1994

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CONTEXTS AND SUBLATTICES OF CONCEPT LATTICES

FRANTIŠEK MACHALA, Olomouc

(Received December 31, 1992)

Any context \mathcal{J} can be uniquely assigned a complete concept lattice $L_{\mathcal{J}}$ (see e.g. [3]). In this paper we describe substructures in \mathcal{J} such that their concept lattices are all complete sublattices in $L_{\mathcal{J}}$. As a consequence a characterization of contexts with distributive or modular concept lattices is obtained. Another characterization for distributive lattices is given in [1].

Definition 1. A context is a triple $\mathcal{J} = (G, M, I)$ where G and M are sets and $I \subseteq G \times M$. For $B \subseteq M, B \neq \emptyset$, we put $B^\downarrow = \{g \in G \mid g I m \ \forall m \in B\}$ and $\emptyset^\downarrow = G$. For $A \subseteq G, A \neq \emptyset$, we put $A^\uparrow = \{m \in M \mid g I m \ \forall g \in A\}$ and $\emptyset^\uparrow = M$. Let further $A^{\uparrow\downarrow} = (A^\uparrow)^\downarrow, B^{\downarrow\uparrow} = (B^\downarrow)^\uparrow$.

Remark 1. From Definition 1 we have: $B_1 \subseteq B_2 \Rightarrow B_2^\downarrow \subseteq B_1^\downarrow$ for $B_1, B_2 \subseteq M$, and $A_1 \subseteq A_2 \Rightarrow A_2^\uparrow \subseteq A_1^\uparrow$ for $A_1, A_2 \subseteq G$.

$$A \subseteq G, \quad A = A^{\uparrow\downarrow} \Leftrightarrow \exists B \subseteq M, \quad B^\downarrow = A, \\ \bigcap_{i \in I} B_i^\downarrow = \left(\bigcup_{i \in I} B_i \right)^\downarrow, \quad B_i \subseteq M \quad \forall i \in I.$$

Theorem 1. If $\mathcal{J} = (G, M, I)$ is a context and $Q = \{A \subseteq G \mid A = A^{\uparrow\downarrow}\}$, then (Q, \subseteq) is a partially ordered set with unit element G . If we put $\bigwedge_{i \in I} A_i = \bigcap_{i \in I} A_i$, $\left(\bigcap_{i \in I} A_i^\downarrow \right)^\downarrow = \bigvee_{i \in I} A_i$ for $A_i \in Q \ \forall i \in I$, then $L_{\mathcal{J}} = (Q, \wedge, \vee)$ is a complete lattice.

Remark 2. The lattice from Theorem 1 is called the *concept lattice* of \mathcal{J} . Suppose $A_i \in Q \ \forall i \in I$ and let $B_i \subseteq M, B_i^\downarrow = A_i \ \forall i \in I$. Then $\bigwedge_{i \in I} A_i = \bigcap_{i \in I} B_i^\downarrow = \left(\bigcup_{i \in I} B_i \right)^\downarrow, \bigvee_{i \in I} A_i = \left(\bigcap_{i \in I} B_i^{\downarrow\uparrow} \right)^\downarrow$.

Let $\mathcal{J} = (G, M, I)$ be a context. For any subset $B \subseteq M$ we put $\bar{B} = \{C \subseteq M \mid C^\downarrow = B^\downarrow\}$ and $D_B = \bigcup_{C \in \bar{B}} C$. Then $D_B^\downarrow = \left(\bigcup_{C \in \bar{B}} C \right)^\downarrow = \bigcap_{C \in \bar{B}} C^\downarrow = B^\downarrow$ holds and hence $D_B \in \bar{B}$.

Definition 2. Let $\mathcal{J} = (G, M, I)$ be a context. A non-empty set $\mathcal{M} \subseteq 2^M$ is *admissible* in \mathcal{J} if for any non-empty subset $\mathcal{B} \subseteq \mathcal{M}$

- (1) there exists $X \in \mathcal{M}$ such that $\bigcup_{B \in \mathcal{B}} B \in \bar{X}$,
- (2) there exists $Y \in \mathcal{M}$ such that $\bigcap_{B \in \mathcal{B}} B^{\uparrow} \in \bar{Y}$.

Remark 3. Examples of admissible sets in a context \mathcal{J} :

- a) $\mathcal{M} = 2^M$,
- b) $\mathcal{M} = \{B\}$, where $B \subseteq M$.
- c) If \mathcal{M} is an admissible set in \mathcal{J} , then $\bar{\mathcal{M}} = \bigcup_{B \in \mathcal{M}} \bar{B}$, $D_{\mathcal{M}} = \{D_B \mid B \in \mathcal{M}\}$ are also admissible sets in \mathcal{J} , and $\mathcal{M}, D_{\mathcal{M}} \subseteq \bar{\mathcal{M}}$.

Theorem 2. Let $\mathcal{J} = (G, M, I)$ be a context.

1. If \mathcal{M} is an admissible subset in \mathcal{J} , then $L_1 = \{B^\downarrow \mid B \in \mathcal{M}\}$ is a complete sublattice of $L_{\mathcal{J}}$.
2. Let L_1 be a complete sublattice of the lattice $L_{\mathcal{J}}$. Let us consider a subset $\mathcal{M} \subseteq 2^M$ such that $B^\downarrow \in L_1 \forall B \in \mathcal{M}$ and for any $x \in L_1$ there exists $X \in \mathcal{M}$ such that $X^\downarrow = x$. Then \mathcal{M} is an admissible subset in \mathcal{J} .

Proof. 1. Evidently $L_1 \subseteq L_{\mathcal{J}}$. Let $\mathcal{A} \subseteq L_1$, $\mathcal{A} \neq \emptyset$. Then there exists a (non-empty) subset $\mathcal{B} \subseteq \mathcal{M}$ such that $\mathcal{A} = \{B^\downarrow \mid B \in \mathcal{B}\}$. We get $\bigwedge \mathcal{A} = \bigwedge_{B \in \mathcal{B}} B^\downarrow =$

$\left(\bigcup_{B \in \mathcal{B}} B \right)^\downarrow$. By (1), there is $X \in \mathcal{M}$ such that $\bigcup_{B \in \mathcal{B}} B \in \bar{X}$, i.e. $\bigwedge \mathcal{A} \in L_1$. Further, $\bigvee \mathcal{A} = \left(\bigcap_{B \in \mathcal{B}} B^{\uparrow} \right)^\downarrow$ and by (2) there is $Y \in \mathcal{M}$ with $\bigcap_{B \in \mathcal{B}} B^{\uparrow} \in \bar{Y}$, which means $\bigvee \mathcal{A} \in L_1$.

2. Consider $\mathcal{B} \subseteq \mathcal{M}$, $\mathcal{B} \neq \emptyset$. We have $\bigwedge_{B \in \mathcal{B}} B^\downarrow \left(\bigcup_{B \in \mathcal{B}} B \right)^\downarrow \in L_1$. Hence there exists $X \in \mathcal{M}$ such that $X^\downarrow = \left(\bigcup_{B \in \mathcal{B}} B \right)^\downarrow$ and thus $\bigcup_{B \in \mathcal{B}} B \in \bar{X}$. Simultaneously we have $\bigvee_{B \in \mathcal{B}} B^\downarrow = \left(\bigcap_{B \in \mathcal{B}} B^{\uparrow} \right)^\downarrow \in L_1$. Similarly as in the previous case there exists $Y \in \mathcal{M}$ such that $\bigcap_{B \in \mathcal{B}} B^{\uparrow} \in \bar{Y}$. □

Remark 4. If \mathcal{M}_1 is an admissible set in a context \mathcal{J} , L_1 the sublattice in $L_{\mathcal{J}}$ corresponding to \mathcal{M}_1 by 1 of Theorem 2, and \mathcal{M}_2 the admissible set in \mathcal{J} by 2

of Theorem 2, then $\overline{\mathcal{M}}_1 = \overline{\mathcal{M}}_2$ (see Remark 3c). If L_1 is a complete sublattice in $L_{\mathcal{J}}$, \mathcal{M} the admissible set in \mathcal{J} corresponding to L_1 by 2 of Theorem 2 and L_2 the lattice corresponding to \mathcal{M} by 1 of Theorem 2, then $L_1 = L_2$.

Definition 3. Let $\mathcal{J} = (G, M, I)$ be a context, G_1, M_1 subsets of G, M and $I_1 \subseteq G_1 \times M_1$. If $I_1 \subseteq I$, then the context $\mathcal{J}_1 = (G_1, M_1, I_1)$ is *embedded* in \mathcal{J} . If $I_1 = I \cap (G_1 \times M_1)$, then \mathcal{J}_1 is a *subcontext* of the context \mathcal{J} .

Definition 4. Let $\mathcal{J} = (G, M, I)$ be a context and $\mathcal{M} \subseteq 2^M$ an admissible set in \mathcal{J} . Put $M_1 = \bigcup_{B \in \mathcal{M}} B, G_1 = \bigcup_{B \in \mathcal{M}} B^\downarrow$ and let for $g \in G_1, m \in M_1: g I_1 m \Leftrightarrow \exists B \in \mathcal{M}, m \in B, g \in B^\downarrow$. The context $\mathcal{J}_{\mathcal{M}} = (G_1, M_1, I_1)$ is \mathcal{M} -*embedded* in \mathcal{J} .

Remark 5. In Definition 4 we have $M_1 \subseteq M, G_1 \subseteq G$ and $g I_1 m \Rightarrow g I m$. Hence the context $\mathcal{J}_{\mathcal{M}}$ is embedded in \mathcal{J} by Definition 3.

Remark 6. Let $\mathcal{J}_{\mathcal{M}} = (G_1, M_1, I_1)$ be a context \mathcal{M} -embedded in a context $\mathcal{J} = (G, M, I)$. By (2), there exists $X \in \mathcal{M}$ such that $\bigcap_{B \in \mathcal{M}} B^{\downarrow\uparrow} \in \overline{X}$. Moreover, $X \in M_1, G_1 = X^\downarrow$ holds.

Theorem 3. If $\mathcal{J}_{\mathcal{M}} = (G_1, M_1, I_1)$ is a context \mathcal{M} -embedded in a context $\mathcal{J} = (G, M, I)$, then the lattice $L_{\mathcal{J}_{\mathcal{M}}}$ is a complete sublattice of the lattice $L_{\mathcal{J}}$ and $L_{\mathcal{J}_{\mathcal{M}}} = \{B^\downarrow \mid B \in \mathcal{M}\}$.

Proof. The symbol \downarrow from Definition 1 will be written in $\mathcal{J}_{\mathcal{M}}$ on the left and in \mathcal{J} on the right (as usual). Hence for any sets $C \subseteq M_1: \downarrow C = \{g \in G_1 \mid g I_1 c \forall c \in C\}$ and $L_{\mathcal{J}_{\mathcal{M}}} = \{\downarrow C \mid C \subseteq M_1\}$. By Remark 5, $\downarrow C \subseteq G_1 \cap C^\downarrow$. Let $\mathcal{B} \subseteq \mathcal{M}, \mathcal{B} \neq \emptyset$, and put $D = \bigcup_{B \in \mathcal{B}} B$. If $g \in D^\downarrow$, then $g I d \forall d \in D$. Moreover, for any $d \in D$ there is $B \in \mathcal{B}$, i.e. $B \in \mathcal{M}$, such that $d \in B$ and since $g I d$, we have $g \in B^\downarrow$. Consequently, $g I_1 d$ and $g \in \downarrow D$. Hence $D^\downarrow \subseteq \downarrow D$ and therefore $D^\downarrow = \downarrow D$.

Consider any set $C \subseteq M_1$ and let $g \in \downarrow C$. Then $g I_1 m \forall m \in C$. For any $m \in C$ there exists $B_m \in \mathcal{M}$ such that $m \in B_m$ and $g \in B_m^\downarrow$. If we put $D = \bigcup_{m \in C} B_m$, then $C \subseteq D$. Because $D \subseteq M_1$, we get, by Remark 1, $\downarrow D \subseteq \downarrow C$ and hence $D^\downarrow \subseteq \downarrow C$. Moreover, $g \in \bigcap_{m \in C} B_m^\downarrow = \left(\bigcup_{m \in C} B_m\right)^\downarrow = D^\downarrow$ and $\downarrow C \subseteq D^\downarrow$. That means $\downarrow C = D^\downarrow$. Since \mathcal{M} is an admissible set in \mathcal{J} , by (1) there exists a set $X \in \mathcal{M}$ such that $D = \bigcup_{m \in C} B_m \in \overline{X}$, so $D^\downarrow = X^\downarrow$ and $\downarrow C = X^\downarrow$.

By Theorem 2, $L_1 = \{B^\downarrow \mid B \in \mathcal{M}\}$ is a complete sublattice in $L_{\mathcal{J}}$. By the preceding, $L_1 = L_{\mathcal{J}_{\mathcal{M}}}$. The lattice operations in L_1 and $L_{\mathcal{J}_{\mathcal{M}}}$ are the same, and hence the lattices L_1 and $L_{\mathcal{J}_{\mathcal{M}}}$ coincide. \square

Theorem 4. Let $\mathcal{J} = (G, M, I)$ be a context and L_1 a complete sublattice of the lattice $L_{\mathcal{J}}$. Let us consider a set $\mathcal{M} \subseteq 2^M$ such that $B^\downarrow \in L_1 \forall B \in \mathcal{M}$ and for any element $x \in L_1$ there exists $B \in \mathcal{M}$ such that $B^\downarrow = x$. If $\mathcal{J}_{\mathcal{M}}$ is a context \mathcal{M} -embedded in \mathcal{J} , then the lattices L_1 and $L_{\mathcal{J}_{\mathcal{M}}}$ coincide.

Proof. By Theorem 2, \mathcal{M} is an admissible set in \mathcal{J} . If we consider a context $\mathcal{J}_{\mathcal{M}}$ \mathcal{M} -embedded in \mathcal{J} , then, by Theorem 3, $L_{\mathcal{J}_{\mathcal{M}}} = \{B^\downarrow \mid B \in \mathcal{M}\} = L_1$. \square

Remark 7. The equality $L_{\mathcal{J}_{\mathcal{M}_1}} = L_{\mathcal{J}_{\mathcal{M}_2}}$ may hold for different sets $\mathcal{M}_1, \mathcal{M}_2$ in \mathcal{J} (see Remark 4 and e.g. [2]).

Theorem 5. Let $\mathcal{J}_{\mathcal{M}} = (G_1, M_1, I_1)$ be a context \mathcal{M} -embedded in a context $\mathcal{J} = (G, M, I)$. The following conditions are equivalent (the mapping \downarrow is denoted in the context \mathcal{J} on the right and in the context $\mathcal{J}_{\mathcal{M}}$ on the left):

1. $\mathcal{J}_{\mathcal{M}}$ is a subcontext in \mathcal{J} .
2. For $g \in G_1, m \in M_1$ we have $g I m \Rightarrow g I_1 m$.
3. For any $m \in M_1$ we have $\downarrow\{m\} = \{m\}^\downarrow \cap G_1$.
4. For any sets $B \subseteq M_1$ we have $\downarrow B = B^\downarrow \cap G_1$.

Proof of this theorem is easy. \square

By using well-known theorems of the lattice theory we get

Theorem 6. Consider a context \mathcal{J} . The concept lattice $L_{\mathcal{J}}$ is distributive (modular) if and only if there is no context \mathcal{M} -embedded in \mathcal{J} with the concept lattice isomorphic to the lattices in Figs. 1, 2 (Fig. 2).




Fig. 1




Fig. 2

Examples. 1. Consider the context $\mathcal{J} = (G, M, I)$ in Fig. 3, where G and M are sets of points and the relation I is denoted by segments connecting the corresponding points. Then $\mathcal{M} = \{\{m_2\}, \{m_2, m_3\}, \{m_2, m_6\}, \{m_2, m_3, m_5\}, \{m_2, m_3, m_6\}, \{m_2, m_3, m_5, m_6\}\}$ is an admissible set in \mathcal{J} . Fig. 4 shows the \mathcal{M} -embedded context $\mathcal{J}_{\mathcal{M}}$ while in Fig. 5 we see the lattice $L_{\mathcal{J}_{\mathcal{M}}}$ ($\mathcal{J}_{\mathcal{M}}$ is a subcontext in \mathcal{J}). By Theorem 6 the lattice $L_{\mathcal{J}}$ is not modular. Fig. 6 shows the lattice $L_{\mathcal{J}}$ and the sublattice $L_{\mathcal{J}_{\mathcal{M}}}$ (marked).




Fig. 3




Fig. 4




Fig. 5




Fig. 6

2. Let us consider the context $\mathcal{J} = (G, M, I)$, where M is the set of planes of the extended three-dimensional Euclidean space E_3 , G is the set of points of this space and I the usual incidence relation. Hence for $m \in M$, $\{m\}^\perp$ is the set of points of the plane m . In Fig. 7 a sublattice L_1 of the lattice $L_{\mathcal{J}}$ is shown. The unit element is




Fig. 7

the set of all points of E_3 , i.e. $1 = G = \emptyset^\perp$. The elements a, b are the sets of points of some planes m_1, m_2 , i.e. $a = \{m_1\}^\perp, b = \{m_2\}^\perp$. An element c is a set of all points of the line r which is a meet of planes m_1, m_2 , i.e. $c = \{m_1, m_2\}^\perp$. The elements p, q, \dots are one-point sets of c (points of the line r). We have $\{p\} = \{m_1, m_2, m\}^\perp$, where m is a plane such that $p \in \{m\}^\perp, r \not\subseteq \{m\}^\perp$. Evidently $\{m_1, m_2, m, n\}^\perp = \emptyset$ where $m,$

n are the planes defined by the preceding two different points. By Theorem 2 the set $\mathcal{M} = \{B \subseteq M \mid B^\downarrow \in L_1\}$ is admissible in \mathcal{J} and determines the \mathcal{M} -embedded context $\mathcal{J}_{\mathcal{M}} = (G_1, M_1, I_1)$ in \mathcal{J} . Then the relation I_1 satisfies

$$\begin{aligned} g \in \{m_1\}^\downarrow &\Rightarrow g I_1 m_1, \\ g \in \{m_2\}^\downarrow &\Rightarrow g I_1 m_2, \\ g \in \{m_1, m_2\}^\downarrow \wedge g \in \{m\}^\downarrow, \quad m \in M_1 &\Rightarrow g I_1 m. \end{aligned}$$

For other points $g \in G$ and planes $m \in M$ the relation I_1 is not defined. By Theorem 4 we obtain $L_{\mathcal{J}_{\mathcal{M}}} = L_1$. The context $\mathcal{J}_{\mathcal{M}}$ is embedded in \mathcal{J} , but it is not a subcontext in \mathcal{J} with a concept lattice L_1 .

References

- [1] *M. Erné*: Distributive Laws for Concept Lattices. Fachbereich Mathematik, Technische Hochschule Darmstadt, 1989, pp. 28.
- [2] *F. Machala*: Isomorphismen von Kontexten und Konzeptualverbänden. Acta Univ. Pal. Olomouensis 110 (1993), 123–139.
- [3] *R. Wille*: Restructuring lattice theory: an approach based on hierarchies of concepts. I. Rival, Ordered sets. Reidel, Dordrecht-Boston, 1982, pp. 445–470.

Author's address: Department of Algebra and Geometry, Palacký University, Tomkova 38, 772 00 Olomouc, Czech Republic.