

M. S. Santos

Compactness theorems for the Bakry-Emery Ricci tensor on semi-Riemannian manifolds

Commentationes Mathematicae Universitatis Carolinae, Vol. 58 (2017), No. 1, 79–86

Persistent URL: <http://dml.cz/dmlcz/146029>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2017

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Compactness theorems for the Bakry-Emery Ricci tensor on semi-Riemannian manifolds

M.S. SANTOS

Abstract. In this manuscript we provide new extensions for the Myers theorem in weighted Riemannian and Lorentzian manifolds. As application we obtain a closure theorem for spatial hypersurfaces immersed in some time-like manifolds.

Keywords: Bakry-Emery Ricci curvature tensor; closure theorem; Riccati equation

Classification: 53C20

1. Introduction

Given a semi-Riemannian manifold (M^n, g) and a smooth function f on M^n , the *weighted manifold* M_f^n associated to M^n and f is the triple $(M^n, g, d\mu = e^{-f} dv)$, where dv is the volume element of M . In the literature it is also called *smooth metric space*. The analysis of the weighted geometry in M_f is deeply related with a family of Ricci tensors introduced by Bakry and Emery in [1] given by

$$\text{Ric}_f^k = \text{Ric} + \text{Hess } f - \frac{1}{k} df \otimes df,$$

where $k \in (0, \infty)$.

In the Riemannian context, the classical Myers compactness theorem [15] was extended for weighted manifolds in many ways. See for instance, [2], [5], [11], [12], [13], [14], [16], [17], [21], [24]. Exploring a Riccati inequality obtained from the Bochner formula we get the following improved version of weighted Myers theorem in the spirit of the Sprouse theorem in [20].

Theorem 1.1. *Let M_f^n be a weighted complete Riemannian manifold. Then for any $\delta > 0$, and $a > 0$, there exists an $\epsilon = \epsilon(n, a, \delta)$ satisfying the following:*

If there is a point p such that along each geodesic γ emanating from p , the Ric_f^k curvature satisfies

$$(1.1) \quad \int_0^\infty \max \{ (n-1)a - \text{Ric}_f^k(\gamma', \gamma'), 0 \} dt < \epsilon(n, a, \delta)$$

then M is compact with $\text{diam}(M) \leq \frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}} + \delta$.

Recently, the geometry of weighted Lorentzian manifold has been subject of great interest. For instance in [4], J. Case has shown that certain aspects of the Bakry-Emery comparison theory can be adapted to a Lorentzian manifold, allowing him to prove a Bakry-Emery version of the Hawking-Penrose singularity theorem for general relativity. This result leads to a Hawking-Penrose theorem for scalar-tensor gravitation theories. In this same context, we can mention Rupert and Woolgar in [19], Woolgar in [22] and Galloway and Woolgar in [9].

Motivated by the ideas of Yun in [23] we obtain an extension of Theorem 1.1 for *weighted globally hyperbolic space-time* (see Section 2.2). The main tool in our approach is a Raychaudhuri type inequality raised by Case in [4]. Precisely, we obtain the following result:

Theorem 1.2. *Let M_f^n be a weighted globally hyperbolic space-time. Then for any $\delta > 0$, $a > 0$, there exists an $\epsilon = \epsilon(n, a, \delta)$ satisfying the following:*

If there is a point p such that along each future directed time-like geodesic γ emanating from p , with $l(\gamma) = \sup\{t \geq 0, d(p, \gamma(t)) = t\}$ the Ric_f^k curvature satisfies

$$\int_0^{l(\gamma)} \max\{(n-1)a - \text{Ric}_f^k(\gamma', \gamma'), 0\} dt < \epsilon(n, a, \delta)$$

then the time-like diameter satisfies $\text{diam}(M) \leq \frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}} + \delta$.

Theorem 1.1 can be applied in the context of space-time manifolds. Namely, suppose that \bar{M}^{n+1} is a space-time and M^n is a spatial hypersurface of \bar{M} . It is an interesting problem to inquire under which conditions M is compact. In this sense, the works of Galloway in [8] and Galloway and Frankel in [7] are very successful. Recently, Cavalcante, Oliveira and the author [5] generalized the closure theorems of [7] and [8] in the weighted case. Here, using Theorem 1.1 we are able to improve Theorem 4.3 of [5] as follows:

Theorem 1.3. *Let M^n be a spatial hypersurface in \bar{M}_f^{n+1} and assume that M is complete in the induced metric. Then for any $\delta > 0$, $a > 0$, there exists an $\epsilon = \epsilon(n, a, \delta)$ satisfying the following:*

If there is a point $p \in M$ such that along each geodesic γ in M emanating from p , the condition

$$\int_0^\infty \max\{(n-1)a - \overline{\text{Ric}}_f^k(X, X) - \langle v(X), X \rangle H_f + \langle a(X), X \rangle, 0\} dt < \epsilon(n, a, \delta)$$

is satisfied, where $X = \gamma'$, then M is compact and $\text{diam}(M) \leq \frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}} + \delta$.

2. Weighted Myers theorems

2.1 The Riemannian case. Given a weighted manifold M_f^n the *weighted Laplacian operator* can be defined by $\Delta_f u = \Delta u - \langle \nabla f, \nabla u \rangle$, for functions u of class

C^2 on M . It is a remarkable fact that the weighted Laplacian satisfies a Bochner type inequality (see [10] or [21]).

Fixed $p \in M$ let $r(x) = d(x, p)$ denote the distance function and let $m_f(r)$ be the weighted Laplacian of the distance function. From the Bochner inequality for the weighted Laplacian one can verify that m_f satisfies the following Riccati inequality (see Appendix A of [21]):

$$(2.1) \quad m'_f + \frac{m_f^2}{n+k-1} \leq -\text{Ric}_f^k(\gamma', \gamma'), \quad \text{for all } t \geq 0,$$

where γ is a geodesic emanating from p and m'_f stands for the derivative of m_f with respect to r . The inequality (2.1) is the main tool in the proof of our first result.

PROOF OF THEOREM 1.1: For any small positive $\epsilon < a^2$ to be determined later, consider the following sets

$$E_1 = \{t \in [0, \infty); \text{Ric}_f^k(\gamma'(t), \gamma'(t)) \geq (n-1)(a - \sqrt{\epsilon})\}$$

and

$$E_2 = \{t \in [0, \infty); \text{Ric}_f^k(\gamma'(t), \gamma'(t)) < (n-1)(a - \sqrt{\epsilon})\}.$$

From the inequality (2.1) we have on E_1

$$(2.2) \quad \frac{\frac{m'_f}{n+k-1}}{\left(\frac{m_f}{n+k-1}\right)^2 + \frac{(n-1)(a-\sqrt{\epsilon})}{n+k-1}} \leq -1.$$

On the other hand, on E_2 we have

$$(2.3) \quad \frac{\frac{m'_f}{n+k-1}}{\left(\frac{m_f}{n+k-1}\right)^2 + \frac{(n-1)(a-\sqrt{\epsilon})}{n+k-1}} \leq \frac{a - \sqrt{\epsilon} - \text{Ric}_f^k(\gamma', \gamma')/(n-1)}{a - \sqrt{\epsilon}}.$$

Now, using the assumption (1.1) on the Bakry-Emery Ricci tensor we get

$$\begin{aligned} \epsilon &> \int_0^\infty \max \{(n-1)a - \text{Ric}_f^k(\gamma'(t), \gamma'(t)), 0\} \\ &> \int_{E_2} \{(n-1)a - \text{Ric}_f^k(\gamma'(t), \gamma'(t))\} dt \\ &> \int_{E_2} \{(n-1)a - (n-1)(a - \sqrt{\epsilon})\} dt \\ &= \mu(E_2)(n-1)\sqrt{\epsilon}. \end{aligned}$$

That is, we have

$$(2.4) \quad \mu(E_2) < \frac{\sqrt{\epsilon}}{n-1},$$

where μ is the Lebesgue measure on \mathbb{R} .

Using inequalities (2.2)–(2.4) we obtain

$$\begin{aligned}
 \int_0^r \frac{\frac{m'_f}{n+k-1}}{\left(\frac{m_f}{n+k-1}\right)^2 + \frac{(n-1)(a-\sqrt{\epsilon})}{n+k-1}} dt &\leq \int_{[0,r] \cap E_1} \frac{\frac{m'_f}{n+k-1}}{\left(\frac{m_f}{n+k-1}\right)^2 + \frac{(n-1)(a-\sqrt{\epsilon})}{n+k-1}} dt \\
 &\quad + \int_{[0,r] \cap E_2} \frac{\frac{m'_f}{n+k-1}}{\left(\frac{m_f}{n+k-1}\right)^2 + \frac{(n-1)(a-\sqrt{\epsilon})}{n+k-1}} dt \\
 &\leq -\mu\{[0,r] \cap E_1\} + \frac{\epsilon}{(n-1)(a-\sqrt{\epsilon})} \\
 &\leq -r + \mu\{[0,r] \cap E_2\} + \frac{\epsilon}{(n-1)(a-\sqrt{\epsilon})} \\
 &\leq -r + \frac{\sqrt{\epsilon}}{n-1} + \frac{\epsilon}{(n-1)(a-\sqrt{\epsilon})}.
 \end{aligned}$$

Define $\tau(\epsilon) = \frac{\sqrt{\epsilon}}{n-1} + \frac{\epsilon}{(n-1)(a-\sqrt{\epsilon})}$. The integral of the left hand side can be computed explicitly and therefore we get

$$\arctan\left(\frac{m_f(r)}{(n+k-1)a(\epsilon)}\right) \leq a(\epsilon)(-r + \tau(\epsilon)) + \frac{\pi}{2},$$

where $a(\epsilon) = \sqrt{\frac{(n-1)(a-\sqrt{\epsilon})}{n+k-1}}$.

So we get

$$m_f(r) \leq -(n+k-1)a(\epsilon) \cot(a(\epsilon)(-r + \tau(\epsilon))),$$

for any r such that $\tau(\epsilon) < r < \frac{\pi}{a(\epsilon)} + \tau(\epsilon)$.

In particular, $m_f(\gamma(r))$ goes to $-\infty$ as $r \rightarrow (\frac{\pi}{a(\epsilon)} + \tau(\epsilon))^+$. It implies that γ cannot be minimal beyond $\frac{\pi}{a(\epsilon)} + \tau(\epsilon)$. Otherwise m_f would be a smooth function at $r = \frac{\pi}{a(\epsilon)} + \tau(\epsilon)$. Taking ϵ explicitly so that $\frac{\pi}{a(\epsilon)} + \tau(\epsilon) = \frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}} + \delta$ and using the completeness of M we have the desired result. \square

2.2 The Lorentzian case. Now let us discuss the Lorentzian version of the Theorem 1.1. Let M be a time-oriented Lorentzian manifold. Given $p \in M$ we set

$$J^+(p) = \{q \in M : \text{there exist a future pointing causal curve from } p \text{ to } q\},$$

called the *causal future* of p . The *causal past* $J^-(p)$ is defined similarly. We say that M is *globally hyperbolic* if the set $J(p, q) := J^+(p) \cap J^-(q)$ is compact for all p and q joined by a causal curve (see [3]). Mathematically, global hyperbolicity often plays a role analogous to geodesic completeness in Riemannian geometry.

Let $\gamma : [a, b] \rightarrow M$ be a future-directed time-like unit-speed geodesic. Given $\{E_1, E_2, \dots, E_n\}$ an orthonormal frame field along γ , and for each $i \in \{1, \dots, n\}$

we let J_i be the unique Jacobi field along c such that $J_i(a) = 0$ and $J'_i(0) = E_i$. Denote by A the matrix $A = [J_1 J_2 \dots J_n]$, where each column is just the vector for J_i in the basis defined by $\{E_i\}$. In this situation, we have that $A(t)$ is invertible if and only if $\gamma(t)$ is not conjugate to $\gamma(a)$.

Now we define $B_f = A'A^{-1} - \frac{1}{n-1}(f \circ \gamma)'E$ wherever A is invertible, where $E(t)$ is the identity map on $(\gamma'(t))^\perp$. The *f-expansion function* is a smooth function defined by $\theta_f = \text{tr } B_f$ (see [4, Definition 2.6]). Note that, if $|\theta_f| \rightarrow \infty$ as $t \rightarrow t_0$, where $t_0 \in [a, b]$, then $\gamma(t_0)$ is conjugate to $\gamma(a)$.

Recently, Case in [4] obtained the following relation between the Bakry-Emery Ricci tensor and the *f-expansion function* θ_f :

Lemma 2.1. *Under the above notations,*

$$(2.5) \quad \theta'_f \leq -\text{Ric}_f^k(\gamma', \gamma') - \frac{\theta_f^2}{k+n-1}.$$

The inequality (2.5) is called *(k, f)-Raychaudhuri inequality*. This inequality is a generalization of the well known Raychaudhuri inequality (see for instance [6]).

The *distance* between two time-like related points is the supremum of lengths of causal curves joining the points. It follows that the distance between any two time-like related points in a globally hyperbolic space-time is the length of such a maximal time-like geodesic. The *time-like diameter*, $\text{diam}(M)$, of a Lorentzian manifold is defined to be the supremum of distances $d(p, q)$ between points of M .

PROOF OF THEOREM 1.2: Let $\epsilon(n, a, \delta)$ be the explicit constant in the previous theorem. Assume by contradiction that there are two points p and q with $d(p, q) > \frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}} + \delta$. On the other hand, since M is globally hyperbolic, there exists a maximal time-like geodesic γ joining p and q such that $\ell(\gamma) = d(p, q)$. Following the steps of the proof of Theorem 1.1 using the *(k, f)-Raychaudhuri inequality* (2.5) we get

$$\lim_{t \rightarrow t_0^+} \theta_f(t) = -\infty,$$

where $t_0 = \frac{\pi}{a(\epsilon)} + \tau(\epsilon)$. So, we conclude that γ cannot be maximal beyond $\frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}} + \delta$ which is a contradiction. □

An immediate consequence of Theorem 1.2 is the Lorentzian version of the original weighted Myers theorem obtained by Qian in [16]. Namely:

Corollary 2.2. Let M_f^n be a weighted globally hyperbolic space-time. Let a be a positive constant and assume that

$$\text{Ric}_f^k(v, v) \geq (n-1)a,$$

for all unit time-like vector field $v \in TM$. Then the time-like diameter satisfies $\text{diam}(M) \leq \frac{\pi}{\sqrt{\frac{(n-1)a}{n+k-1}}}$.

3. Application

Let \bar{M}^{n+1} be an $(n+1)$ -dimensional space-time manifold. Let M^n be a spatial hypersurface immersed in \bar{M}^{n+1} . That is, the metric induced in M^n by \bar{M}^{n+1} is a Riemannian metric. Then the unit tangent vectors to the future directed time-like geodesics orthogonal to M define a smooth unit time-like vector field \mathbf{u} in a neighborhood of M . For the sake of simplicity we will omit the dimension super index and we always use a *bar* for geometric objects related to \bar{M} .

Let X be a vector field tangent to M . Extend X along the flow lines generated by \mathbf{u} , and suppose that X is invariant under the flow generated by \mathbf{u} , i.e, $[X, \mathbf{u}] = 0$, where $[\cdot, \cdot]$ is the Lie bracket. The *velocity* and *acceleration* of X along the flow generated by \mathbf{u} are given respectively by

$$v(X) = \bar{\nabla}_{\mathbf{u}}X \quad \text{and} \quad a(X) = \bar{\nabla}_{\mathbf{u}}\bar{\nabla}_{\mathbf{u}}X.$$

The second fundamental form of M as a hypersurface of \bar{M} is defined by $b(X) = -\bar{\nabla}_X\mathbf{u}$. Let H denote its *mean curvature* function. We point out that $H = -\bar{\text{div}}\mathbf{u}$, that is, the averaged *Hubble expansion parameter* at points of M in relativistic cosmology (see [18, §3.3.1] or [6, p. 161]).

The *f-mean curvature* or *weighted mean curvature* of M is defined by

$$H_f = H - (\bar{\nabla}f)^\perp,$$

where $(\bar{\nabla}f)^\perp$ is a normal projection of $\bar{\nabla}f$ on M .

We will use the following technical lemma obtained in [7] (see also [5]).

Lemma 3.1. *Under the above conditions, let $\gamma(s)$ be a normalized geodesic in M and set $X = \gamma'$. Then*

$$(3.1) \quad \begin{aligned} \text{Ric}(X, X) &= \overline{\text{Ric}}(X, X) - \langle a(X), X \rangle + \langle v(X), X \rangle H + \langle v(X), X \rangle^2 \\ &+ \sum_{j=2}^n \langle v(X), e_j \rangle^2, \end{aligned}$$

where $\{e_1 = X, e_2, \dots, e_n\}$ is an orthonormal basis of T_pM .

Finally, we are in position to prove our closure theorem.

PROOF OF THEOREM 1.3: Taking into account that X is invariant under the flow, we get $v(X) = -b(X)$. Then

$$(3.2) \quad \begin{aligned} \overline{\text{Hess}} f(X, X) &= \langle \bar{\nabla}_X \bar{\nabla}f, X \rangle \\ &= \text{Hess } f(X, X) + \langle \bar{\nabla}f, \mathbf{u} \rangle \langle b(X), X \rangle \\ &= \text{Hess } f(X, X) - \langle \bar{\nabla}f, \mathbf{u} \rangle \langle v(X), X \rangle. \end{aligned}$$

From equations (1), (3.1) and (3.2) we get easily

$$\text{Ric}_f^k(X, X) \geq \overline{\text{Ric}}_f^k(X, X) - \langle a(X), X \rangle + \langle v(X), X \rangle H_f,$$

for all $k > 0$. Thus

$$(n-1)a - \text{Ric}_f^k(X, X) \leq (n-1)a - \overline{\text{Ric}}_f^k(X, X) + \langle a(X), X \rangle - \langle v(X), X \rangle H_f.$$

Applying Theorem 1.1 we have the desired result. \square

Acknowledgment. The results in this paper are part of the author's doctoral thesis. While the author was at Universidade Federal de Alagoas, he was fully supported by CAPES-Brazil. He wants to thank M. Cavalcante for the valuable suggestions and comments.

REFERENCES

- [1] Bakry D., Emery E., *Diffusions hypercontractives*, Séminaire de probabilités, XIX, 1983/84, Lecture Notes in Math., 1123, Springer, Berlin, 1985, pp. 177–206.
- [2] Bakry D., Ledoux M., *Sobolev inequalities and Myers diameter theorem for an abstract Markov generator*, Duke Math. J. **81** (1996), no. 1, 252–270.
- [3] Beem J., Ehrlich P., Easley K., *Global Lorentzian Geometry*, 2nd edn., Marcel Dekker, New York, 1996.
- [4] Case J., *Singularity theorems and the lorentzian splitting theorem for the Bakry-Emery-Ricci tensor*, J. Geom. Phys. **60** (2010), no. 3, 477–490.
- [5] Cavalcante M.P., Oliveira J.Q., Santos M.S., *Compactness in weighted manifolds and applications*, Results Math. **68** (2015), 143–156.
- [6] Frankel T., *Gravitation Curvature. An Introduction to Einstein's Theory*, W.H. Freeman and Co., San Francisco, Calif., 1979.
- [7] Frankel T., Galloway G., *Energy density and spatial curvature in general relativity*, J. Math. Phys. **22** (1981), no. 4, 813–817.
- [8] Galloway G.J., *A generalization of Myers theorem and an application to relativistic cosmology*, J. Differential Geom. **14** (1979), 105–116.
- [9] Galloway G.J., Woolgar E., *Cosmological singularities in Bakry-Émery space-times*, preprint, 2013.
- [10] Ledoux M., *The geometry of Markov diffusion generators*, Ann. Fac. Sci. Toulouse Math. **9** (2000), no. 2, 305–366.
- [11] Limoncu M., *The Bakry-Emery Ricci tensor and its applications to some compactness theorems*, Math. Z. **271** (2012), 715–722.
- [12] Limoncu M., *Modifications of the Ricci tensor and applications*, Arch. Math. (Basel) **95** (2010), 191–199.
- [13] Lott J., *Some geometric properties of the Bakry-Émery-Ricci tensor*, Comment. Math. Helv. **78** (2003), no. 4, 865–883.
- [14] Morgan F., *Myers' theorem with density*, Kodai Math. J. **29** (2006), no. 3, 454–461.
- [15] Myers S.B., *Riemannian manifolds with positive mean curvature*, Duke Math. J. **8** (1941) 401–404.
- [16] Qian Z., *Estimates for weighted volumes and applications*, Quart. J. Math. Oxford **48** (1997), 235–242.
- [17] Rimoldi M., *A remark on Einstein warped products*, Pacific J. Math. **252** (2011), no. 1, 207–218.
- [18] Ringström H., *On the Topology and Future Stability of the Universe*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2013.
- [19] Rupert M., Woolgar E., *Bakry-Émery black holes*, Classical Quantum Gravity **31** (2014), no. 2, 025008.

- [20] Sprouse S., *Integral curvature bounds and bounded diameter*, Comm. Anal. Geom. **8** (2000), 531–543.
- [21] Wei G., Wylie W., *Comparison geometry for the Bakry-Emery Ricci tensor*, J. Differential Geom. **83** (2009), no. 2, 377–405.
- [22] Woolgar E., *Scalar-tensor gravitation and the Bakry-Emery-Ricci tensor*, Classical Quantum Gravity **30** (2013) 085007.
- [23] Yun J.-G., *A note on the generalized Myers theorem*, Bull. Korean Math. Soc. **46** (2009), no. 1, 61–66.
- [24] Zhang S., *A theorem of Ambrose for Bakry-Emery Ricci tensor*, Ann. Global Anal. Geom. **45** (2014), no. 3, 233–238.

UNIVERSIDADE FEDERAL DO CEARÁ, CAMPUS RUSSAS, CE, CEP 62900-000, BRAZIL

E-mail: marciosantos@ufc.br

(Received February 2, 2016)