

Bui Duc Nam; Nguyen Huu Nhan; Le Thi Phuong Ngoc; Nguyen Thanh Long
On a system of nonlinear wave equations with the Kirchhoff-Carrier and
Balakrishnan-Taylor terms

Mathematica Bohemica, Vol. 147 (2022), No. 2, 237–270

Persistent URL: <http://dml.cz/dmlcz/150331>

Terms of use:

© Institute of Mathematics AS CR, 2022

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A SYSTEM OF NONLINEAR WAVE EQUATIONS
WITH THE KIRCHHOFF-CARRIER
AND BALAKRISHNAN-TAYLOR TERMS

BUI DUC NAM, NGUYEN HUU NHAN, Ho Chi Minh City,

LE THI PHUONG NGOC , Nha Trang City,

NGUYEN THANH LONG , Ho Chi Minh City

Received May 19, 2020. Published online July 8, 2021.

Communicated by Reinhard Farwig

Abstract. We study a system of nonlinear wave equations of the Kirchhoff-Carrier type containing a variant of the Balakrishnan-Taylor damping in nonlinear terms. By the linearization method together with the Faedo-Galerkin method, we prove the local existence and uniqueness of a weak solution. On the other hand, by constructing a suitable Lyapunov functional, a sufficient condition is also established to obtain the exponential decay of weak solutions.

Keywords: system of nonlinear wave equations of Kirchhoff-Carrier type; Balakrishnan-Taylor term; Faedo-Galerkin method; local existence; exponential decay

MSC 2020: 35L20, 35L70, 35Q74, 37B25

1. INTRODUCTION

We consider the initial-boundary value problem for the system of nonlinear wave equations with the Kirchhoff-Carrier and Balakrishnan-Taylor terms

$$(1.1) \quad \begin{aligned} u_{tt} - \lambda u_{xxt} - \mu_1(t, \langle u_x(t), u_{xt}(t) \rangle) u_{xx} &= f_1(x, t, u, v, u_x, v_x, u_t, v_t), \\ v_{tt} - \mu_2(t, \|v(t)\|^2, \|v_x(t)\|^2) v_{xx} &= f_2(x, t, u, v, u_x, v_x, u_t, v_t), \end{aligned}$$

This research was funded by the Vietnam National University Ho Chi Minh City (VNU-HCM) under Grant No. B2020-18-01.

$0 < x < 1, 0 < t < T$, associated with the Robin-Dirichlet conditions

$$(1.2) \quad u(0, t) = u(1, t) = v(1, t) = v_x(0, t) - \zeta v(0, t) = 0$$

and initial conditions

$$(1.3) \quad (u(x, 0), v(x, 0)) = (\tilde{u}_0(x), \tilde{v}_0(x)), \quad (u_t(x, 0), v_t(x, 0)) = (\tilde{u}_1(x), \tilde{v}_1(x)),$$

where $\lambda > 0, \zeta \geq 0$ are given constants and $\tilde{u}_i, \tilde{v}_i, \mu_i, f_i$ ($i = 1, 2$) are given functions satisfying conditions, which will be specified later. In (1.1), the nonlinear terms $\mu_1(t, \langle u_x(t), u_{xt}(t) \rangle), \mu_2(t, \|v(t)\|^2, \|v_x(t)\|^2)$ depend on the integrals $\langle u_x(t), u_{xt}(t) \rangle = \int_0^1 u_x(x, t) u_{xt}(x, t) dx, \|v(t)\|^2 = \int_0^1 v^2(x, t) dx$ and $\|v_x(t)\|^2 = \int_0^1 v_x^2(x, t) dx$.

The system (1.1) is regarded as the combination between the wave equation of the Kirchhoff-Carrier type and that with the Balakrishnan-Taylor damping, in which a related case of (1.1)₁ was proposed by Balakrishnan and Taylor in 1989, see [1]. They established the following new model for flight structures with viscous and nonlinear nonlocal damping in the one dimensional case

$$(1.4) \quad \rho u_{tt} + EI u_{xxxxx} - c u_{xxt} - \left(H + \frac{EA}{2L} \int_0^L |u_x|^2 dx + \tau \left| \int_0^L u_x u_{xt} dx \right|^{2(N+\eta)} \int_0^L u_x u_{xt} dx \right) u_{xx} = 0,$$

where $u = u(x, t)$ represents the transversal deflection of an extensible beam of length $2L > 0$ in the rest position, $\rho > 0$ is the mass density, E is Young's modulus of elasticity, I is the cross-sectional moment of inertia, H is the axial force (either traction or compression), A is the cross-sectional area, $c > 0$ is the coefficient of viscous damping, $\tau > 0$ is the Balakrishnan-Taylor damping coefficient, $0 \leq \tau < 1, 0 \leq \eta < \frac{1}{2}$ and $N \in \mathbb{N}$. Equation (1.4) seems to be related to the panel flutter equation and spillover problem, we refer the reader to [2] for more information. Since then, the equation with the Balakrishnan-Taylor damping was studied in many papers in which the properties of the solution such as stability, decay and blow-up in time are considered, see [3]–[6], [8], [9], [12]–[17], [20]–[22], [27], [31], [32], [35] and references therein. Some authors were interested in the effects of time-varying delay, which appear in many applications to science because physical, chemical, biological, thermal, and economical phenomena naturally not only depend on the present state but also on some past occurrences, see examples in [17], [20], [21].

In 2011, Emmrich and Thalhammer considered a class of integro-differential equations with applications in nonlinear elastodynamics. They proposed a general model for the description of nonlinear extensible beams incorporating the weak, viscous,

strong and Balakrishnan-Taylor damping as follows (see [8], equation (1.1))

$$(1.5) \quad u_{tt} + \alpha \Delta^2 u + \xi u + \kappa u_t - \lambda \Delta u_t + \mu \Delta^2 u_t - \left(\beta + \gamma \int_{\Omega} |\nabla u|^2 dx + \delta \left| \int_{\Omega} \nabla u \nabla u_t dx \right|^{q-2} \int_{\Omega} \nabla u \nabla u_t dx \right) \Delta u = h$$

in $\Omega \times (0, \infty)$, where $\Omega \subset \mathbb{R}^n$ is a bounded domain. The constants have physical meaning: $\alpha > 0$ is the elasticity coefficient, $\gamma > 0$ is the extensibility coefficient, $\lambda \geq 0$ is the viscous damping coefficient, $\mu \geq 0$ is the strong damping coefficient, $\delta \geq 0$ is the Balakrishnan-Taylor damping coefficient, $\beta \in \mathbb{R}$ is the axial force coefficient ($\beta > 0$ means traction or $\beta < 0$ compression), $\kappa \in \mathbb{R}$ is the weak damping coefficient (although without the sign condition), $\xi \in \mathbb{R}$ is source coefficient and the exponent q belongs to $[2, \infty)$. In [12], Tavares et al. worked with an alternative expression of the Balakrishnan-Taylor term

$$(1.6) \quad \Phi(u, u_t) = \int_{\Omega} \nabla u \nabla u_t dx = - \int_{\Omega} (\Delta u) u_t dx$$

to study the well-posedness and long-time dynamics of the class of extensible beams with the Balakrishnan-Taylor and frictional damping

$$(1.7) \quad u_{tt} + \Delta^2 u - \left(\beta + \gamma \int_{\Omega} |\nabla u|^2 dx + \delta |\Phi(u, u_t)|^{q-2} \Phi(u, u_t) \right) \Delta u + \kappa u_t + f(u) = h,$$

in $\Omega \times \mathbb{R}_+$, where Ω is a bounded domain in \mathbb{R}^n with the smooth boundary $\Gamma = \partial\Omega$. In [27], the authors proved the existence and uniqueness of the initial-boundary value problem

$$(1.8) \quad \begin{cases} u_{tt} - \lambda u_{xxt} - \mu(t, \langle u_x(t), u_{xt}(t) \rangle, \|u(t)\|^2, \|u_x(t)\|^2) u_{xx} \\ \quad = f(x, t, u, u_t, u_x, \langle u_x(t), u_{xt}(t) \rangle, \|u(t)\|^2, \|u_x(t)\|^2), & 0 < x < 1, 0 < t < T, \\ u(0, t) = u(1, t) = 0, \\ u(x, 0) = \tilde{u}_0(x), u_t(x, 0) = \tilde{u}_1(x), \end{cases}$$

where $\mu, f, \tilde{u}_0, \tilde{u}_1$ are given functions, $\lambda > 0$ is a given constant. When $\mu = B(\|u_x(t)\|^2) + \sigma(\langle u_x(t), u_{xt}(t) \rangle)$ and $f = -\lambda_1 u_t + f(u) + F(x, t)$, they put suitable hypotheses and sufficient conditions for the nonlinear Balakrishnan-Taylor damping $\sigma(\langle u_x(t), u_{xt}(t) \rangle)$ to get the exponential decay of solution. When $f = 0$ and μ depends only on $\|v(t)\|^2$ and $\|v_x(t)\|^2$, equation (1.8)₁ reduces to the equation of the Kirchhoff-Carrier type, describing nonlinear vibrations of an elastic string, which were studied in [7], [25], [34].

In [19], Jamil and Fetecau studied a mathematical model describing helical flows of the Maxwell fluid in an annular region between two infinite coaxial circular cylinders of radii 1 and $R > 1$ below

$$(1.9) \quad \begin{cases} \lambda u_{tt} + u_t = \nu \left(u_{xx} + \frac{1}{x} u_x - \frac{1}{x^2} u \right), & 1 < x < R, t > 0, \\ \lambda v_{tt} + v_t = \nu \left(v_{xx} + \frac{1}{x} v_x \right), & 1 < x < R, t > 0, \\ u_x(1, t) - u(1, t) = \frac{f}{\mu} t, \quad v_t(1, t) = \frac{g}{\mu} t, & t > 0, \\ u(R, t) = v(R, t) = 0, & t > 0, \\ u(x, 0) = u_t(x, 0) = 0, & 1 < x < R, \\ v(x, 0) = v_t(x, 0) = 0, & 1 < x < R, \end{cases}$$

where λ , μ , ν , f , and g are given constants. The authors obtained an exact solution for this problem by means of finite Hankel transforms and presented it in the series form in terms of Bessel functions $J_0(x)$, $Y_0(x)$, $J_1(x)$, $Y_1(x)$, $J_2(x)$ and $Y_2(x)$, satisfying all imposed initial and boundary conditions. Other works on helical flows for ordinary and fractional derivative models can be found in [28], [18] and [33]. Some recent works focus on studying the porous elastic systems with nonlinear damping, see [11], [10], [24], [29] and references therein. It seems that the first research about the system of equations of the Kirchhoff type with the Balakrishnan-Taylor damping is given in [26], where its model is described as

$$(1.10) \quad \begin{cases} u_{tt} - \left(a + b \|\nabla u\|^2 + \sigma \int_{\Omega} \nabla u \nabla u_t \, dx \right) \Delta u \\ \quad + \int_0^t g_1(t-s) \Delta u(s) \, ds = f_1(u, v), \quad t > 0, x \in \Omega, \\ v_{tt} - \left(a + b \|\nabla v\|^2 + \sigma \int_{\Omega} \nabla v \nabla v_t \, dx \right) \Delta v \\ \quad + \int_0^t g_2(t-s) \Delta v(s) \, ds = f_2(u, v), \quad t > 0, x \in \Omega, \\ u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x), & x \in \Omega, \\ v(x, 0) = v_0(x), \quad v_t(x, 0) = v_1(x), & x \in \Omega, \\ u(x, t) = v(x, t) = 0, & (x, t) \in \Gamma \times [0, \infty), \end{cases}$$

where Ω is a bounded domain in \mathbb{R}^n with a smooth boundary Γ . By the energy method, Mu and Ma in [26] obtained an arbitrary decay of solutions according to the relaxation functions. To the best of our knowledge, the system of equations of the Kirchhoff-Carrier type with the Balakrishnan-Taylor damping (1.1) has not

been extensively studied. Motivated by the above articles, we survey the unique existence and exponential decay of weak solutions of the problem (1.1). Our plan in this paper is as follows. In Section 2, we present some preliminaries. In Section 3, by applying the linearization method together with the Faedo-Galerkin method and the weak compact method, we prove the local existence and the uniqueness of a weak solution (Theorem 3.7). In Section 4, by constructing a suitable Lyapunov functional, we prove a condition sufficient to obtain the exponential decay of weak solutions (Theorem 4.2). The results obtained here are a relative generalization of [24], [27] and [34] by improving and developing these previous works essentially.

2. PRELIMINARIES

First, we put $\Omega = (0, 1)$, $Q_T = \Omega \times (0, T)$, $T > 0$ and denote the usual function spaces used in this paper by the notations $L^p = L^p(\Omega)$, $H^m = H^m(\Omega)$. Let $\langle \cdot, \cdot \rangle$ be either the scalar product in L^2 or the dual pairing of a continuous linear functional and an element of a function space. The notation $\|\cdot\|$ stands for the norm in L^2 and we denote by $\|\cdot\|_X$ the norm in the Banach space X . We call X' the dual space of X .

We denote by $L^p(0, T; X)$, $1 \leq p \leq \infty$, the Banach space of real functions $u: (0, T) \rightarrow X$, measurable and such that $\|u\|_{L^p(0, T; X)} < \infty$ with

$$\|u\|_{L^p(0, T; X)} = \begin{cases} \left(\int_0^T \|u(t)\|_X^p dt \right)^{1/p} & \text{if } 1 \leq p < \infty, \\ \text{ess sup}_{0 < t < T} \|u(t)\|_X & \text{if } p = \infty. \end{cases}$$

On $H^1 \equiv H^1(\Omega)$, we use the norm

$$\|v\|_{H^1} = \sqrt{\|v\|^2 + \|v_x\|^2}.$$

We put

$$(2.1) \quad V = \{v \in H : v(1) = 0\},$$

$$(2.2) \quad a(u, v) = \int_0^1 u_x(x)v_x(x) dx + \zeta u(0)v(0) \quad \forall u, v \in V.$$

The set V is a closed subspace of H^1 and on V , the three norms $\|v\|_{H^1}$, $\|v_x\|$ and $\|v\|_a = \sqrt{a(v, v)}$ are equivalent.

We state the following lemmas, the proofs of which are straightforward, and hence we omit the details.

Lemma 2.1. *The imbedding $H^1 \hookrightarrow C^0(\overline{\Omega})$ is compact and for all $v \in H^1$,*

$$\|v\|_{C^0(\overline{\Omega})} \leq \sqrt{2}\|v\|_{H^1}.$$

Lemma 2.2. *Let $\zeta \geq 0$. Then the imbedding $V \hookrightarrow C^0(\bar{\Omega})$ is compact and for all $v \in V$,*

$$\begin{aligned} \|v\|_{C^0(\bar{\Omega})} &\leq \|v_x\| \leq \|v\|_a, \\ \frac{1}{\sqrt{2}}\|v\|_{H^1} &\leq \|v_x\| \leq \|v\|_a \leq \sqrt{1+\zeta}\|v\|_{H^1}. \end{aligned}$$

Lemma 2.3. *Let $\zeta \geq 0$. Then the symmetric bilinear form $a(\cdot, \cdot)$ defined by (2.2) is continuous on $V \times V$ and coercive on V . Furthermore,*

- (i) $|a(u, v)| \leq (1 + \zeta)\|u_x\|\|v_x\|$ for all $u, v \in V$,
- (ii) $a(v, v) \geq \|v_x\|^2$ for all $v \in V$.

Lemma 2.4. *Let $\zeta \geq 0$. Then there exists a Hilbert orthonormal base $\{\tilde{\varphi}_j\}$ of L^2 consisting of the eigenfunctions $\tilde{\varphi}_j$ corresponding to the eigenvalue $\tilde{\lambda}_j$ such that*

$$\begin{aligned} 0 < \tilde{\lambda}_1 \leq \tilde{\lambda}_2 \leq \dots \leq \tilde{\lambda}_j \leq \dots, \quad \lim_{j \rightarrow \infty} \tilde{\lambda}_j = \infty; \\ a(\tilde{\varphi}_j, v) = \tilde{\lambda}_j \langle \tilde{\varphi}_j, v \rangle \quad \forall v \in V, \quad j = 1, 2, \dots \end{aligned}$$

Furthermore, the sequence $\{\tilde{\varphi}_j \tilde{\lambda}_j^{-1/2}\}$ is also a Hilbert orthonormal basis of V with respect to the scalar product $a(\cdot, \cdot)$.

On the other hand, we also have that $\tilde{\varphi}_j$ satisfy the boundary value problems

$$\begin{cases} -\Delta \tilde{\varphi}_j = \tilde{\lambda}_j \tilde{\varphi}_j & \text{in } (0, 1), \\ \tilde{\varphi}_{jx}(0) - \zeta \tilde{\varphi}_j(0) = \tilde{\varphi}_j(1) = 0, \quad \tilde{\varphi}_j \in C^\infty(\bar{\Omega}). \end{cases}$$

Proof. The proof of Lemma 2.4 can be found in [30], page 87, Theorem 7.7 with $H = L^2$ and $a(\cdot, \cdot)$ is defined by (2.2). \square

3. THE EXISTENCE AND UNIQUENESS THEOREM

Considering $T^* > 0$ fixed, we make the following assumptions:

- (H₁) $\tilde{u}_0, \tilde{u}_1 \in H_0^1 \cap H^2$, $\tilde{v}_0 \in V \cap H^2$, $\tilde{v}_1 \in V$, $\tilde{v}_{0x} - \zeta \tilde{v}_0 = 0$,
- (H₂) $\mu_1 \in C^1([0, T^*] \times \mathbb{R})$, $\mu_2 \in C^1([0, T^*] \times \mathbb{R}_+^2)$, and there exist positive constants μ_{1*}, μ_{2*} such that $\mu_1(t, y) \geq \mu_{1*}$ for all $(t, y) \in [0, T^*] \times \mathbb{R}$ and $\mu_2(t, y, z) \geq \mu_{2*}$ for all $(t, y, z) \in [0, T^*] \times \mathbb{R}_+^2$,
- (H₃) $f_i \in C^1([0, 1] \times [0, T^*] \times \mathbb{R}^6)$, $i = 1, 2$.

Let $\lambda > 0$, $\zeta \geq 0$. For every $T \in (0, T^*]$, we say that (u, v) is a weak solution of the problem (1.1)–(1.3) if

$$(u, v) \in \widetilde{W}_T = \{(u, v) \in L^\infty(0, T; (H_0^1 \cap H^2) \times (V \cap H^2)) : \\ (u', v') \in L^\infty(0, T; (H_0^1 \cap H^2) \times V), \\ u'' \in L^2(0, T; H_0^1) \cap L^\infty(0, T; L^2), v'' \in L^\infty(0, T; L^2)\}$$

and (u, v) satisfies the variational problem

$$(3.1) \quad \begin{cases} \langle u''(t), \varphi \rangle + \lambda \langle u'_x(t), \varphi_x \rangle + \mu_1[u](t) \langle u_x(t), \varphi_x \rangle = \langle f_1[u, v](t), \varphi \rangle, \\ \langle v''(t), \widetilde{\varphi} \rangle + \mu_2[v](t) a(v(t), \widetilde{\varphi}) = \langle f_2[u, v](t), \widetilde{\varphi} \rangle \end{cases}$$

for all $(\varphi, \widetilde{\varphi}) \in H_0^1 \times V$ and a.e. $t \in (0, T)$, together with the initial conditions

$$(3.2) \quad (u(0), u'(0)) = (\widetilde{u}_0, \widetilde{u}_1), \quad (v(0), v'(0)) = (\widetilde{v}_0, \widetilde{v}_1),$$

where

$$(3.3) \quad \begin{cases} \mu_1[u](t) = \mu_1(t, \langle u_x(t), u'_x(t) \rangle), \\ \mu_2[v](t) = \mu_2(t, \|v(t)\|^2, \|v_x(t)\|^2), \\ f_i[u, v](x, t) = f_i(x, t, u(x, t), v(x, t), u_x(x, t), v_x(x, t), u'(x, t), v'(x, t)), \quad i = 1, 2. \end{cases}$$

For each $M > 0$ given, we set the constants $K_M = K_M(f_1, f_2)$, $\widetilde{K}_M = \widetilde{K}_M(\mu_1, \mu_2)$ as

$$(3.4) \quad \begin{cases} K_M = K_M(f_1, f_2) = \sum_{j=1}^2 \|f_j\|_{C^1(A_M)} = \sum_{j=1}^2 \sum_{i=0}^8 \|D_i f_j\|_{C^0(A_M)}, \\ \widetilde{K}_M = \widetilde{K}_M(\mu_1, \mu_2) = \sum_{j=1}^2 \|\mu_j\|_{C^1(A_M^{(j)})} \\ = \|\mu_1\|_{C^0(A_M^{(1)})} + \sum_{i=1}^2 \|D_i \mu_1\|_{C^0(A_M^{(1)})} + \|\mu_2\|_{C^0(A_M^{(2)})} + \sum_{i=1}^3 \|D_i \mu_2\|_{C^0(A_M^{(2)})}, \end{cases}$$

where

$$(3.5) \quad \begin{cases} \|f_j\|_{C^0(A_M)} = \sup_{(x, t, y_1, \dots, y_6) \in A_M} |f_j(x, t, y_1, \dots, y_6)|, \quad j = 1, 2, \\ \|\mu_1\|_{C^0(A_M^{(1)})} = \sup_{(t, y) \in A_M^{(1)}} |\mu_1(t, y)|, \\ \|\mu_2\|_{C^0(A_M^{(2)})} = \sup_{(t, y, z) \in A_M^{(2)}} |\mu_2(t, y, z)|, \\ A_M = [0, 1] \times [0, T^*] \times [-M, M]^6, \\ A_M^{(1)} = [0, T^*] \times [-M^2, M^2], \\ A_M^{(2)} = [0, T^*] \times [0, M^2]^2. \end{cases}$$

For each $T \in (0, T^*]$, we put

$$(3.6) \quad V_T = \{(u, v) \in L^\infty(0, T; (H_0^1 \cap H^2) \times (V \cap H^2)) : \\ (u', v') \in L^\infty(0, T; (H_0^1 \cap H^2) \times V), (u'', v'') \in L^2(0, T; H_0^1 \times L^2)\};$$

it is a Banach space with respect to the norm

$$(3.7) \quad \|(u, v)\|_{V_T} = \max\{\|(u, v)\|_{L^\infty(0, T; (H_0^1 \cap H^2) \times (V \cap H^2))}, \\ \|(u', v')\|_{L^\infty(0, T; (H_0^1 \cap H^2) \times V)}, \|(u'', v'')\|_{L^2(0, T; H_0^1 \times L^2)}\}.$$

For every $M > 0$, we put

$$(3.8) \quad W(M, T) = \{(u, v) \in V_T : \|(u, v)\|_{V_T} \leq M\}, \\ W_1(M, T) = \{(u, v) \in W(M, T) : (u'', v'') \in L^\infty(0, T; L^2 \times L^2)\}.$$

Now, we establish the following recurrent sequence $\{(u_m, v_m)\}$. The first term is chosen as $(u_0, v_0) \equiv (0, 0)$; suppose that

$$(3.9) \quad (u_{m-1}, v_{m-1}) \in W_1(M, T),$$

then we associate (1.1)–(1.3) with the following problem.

Find $(u_m, v_m) \in W_1(M, T)$ ($m \geq 1$) which satisfies the linear variational problem

$$(3.10) \quad \begin{cases} \langle u_m''(t), \varphi \rangle + \lambda \langle u_m'(t), \varphi_x \rangle + \mu_{1m}(t) \langle u_m(t), \varphi_x \rangle = \langle F_{1m}(t), \varphi \rangle, \\ \langle v_m''(t), \tilde{\varphi} \rangle + \mu_{2m}(t) a(v_m(t), \tilde{\varphi}) = \langle F_{2m}(t), \tilde{\varphi} \rangle \end{cases}$$

for all $(\varphi, \tilde{\varphi}) \in H_0^1 \times V$ and a.e. $t \in (0, T)$, together with the initial conditions

$$(3.11) \quad (u_m(0), u_m'(0)) = (\tilde{u}_0, \tilde{u}_1), \quad (v_m(0), v_m'(0)) = (\tilde{v}_0, \tilde{v}_1),$$

where

$$(3.12) \quad \begin{aligned} \mu_{1m}(t) &= \mu_1[u_{m-1}](t) = \mu_1(t, \langle \nabla u_{m-1}(t), \nabla u_{m-1}'(t) \rangle), \\ \mu_{2m}(t) &= \mu_2[v_{m-1}](t) = \mu_2(t, \|v_{m-1}(t)\|^2, \|\nabla v_{m-1}(t)\|^2), \\ F_{im}(x, t) &= f_i[u_{m-1}, v_{m-1}](t) \\ &= f_i(x, t, u_{m-1}(x, t), v_{m-1}(x, t), \nabla u_{m-1}(x, t), \nabla v_{m-1}(x, t), \\ &\quad u_{m-1}'(x, t), v_{m-1}'(x, t)), \quad i = 1, 2. \end{aligned}$$

Then we have the following theorem.

Theorem 3.1. *Let (H₁)–(H₃) hold. Then there exist constants $M, T > 0$ such that, for $(u_0, v_0) \equiv (0, 0)$, there exists a recurrent sequence $\{(u_m, v_m)\} \subset W_1(M, T)$ defined by (3.9)–(3.12).*

Proof. The proof of Theorem 3.1 consists of three steps.

Step 1. The Faedo-Galerkin approximation. Let $\{\varphi_j\}$ be a basis of H_0^1 formed by the eigenfunction φ_j of the operator $-\Delta = -\frac{\partial^2}{\partial x^2}$ such that $-\Delta\varphi_j = \lambda_j\varphi_j$, $\varphi_j \in H_0^1 \cap C^\infty([0, 1])$, $\varphi_j(x) = \sqrt{2}\sin(j\pi x)$, $\lambda_j = (j\pi)^2$, $j = 1, 2, \dots$, and let $\{\tilde{\varphi}_j\}$ be the basis of V as in Lemma 2.4. Put

$$(3.13) \quad u_m^{(k)}(t) = \sum_{j=1}^k c_{mj}^{(k)}(t)\varphi_j, \quad v_m^{(k)}(t) = \sum_{j=1}^k d_{mj}^{(k)}(t)\tilde{\varphi}_j,$$

where the coefficients $c_{mj}^{(k)}(t)$, $d_{mj}^{(k)}(t)$ satisfy the system of linear differential equations

$$(3.14) \quad \begin{cases} \langle \ddot{u}_m^{(k)}(t), \varphi_j \rangle + \lambda \langle \dot{u}_{mx}^{(k)}(t), \varphi_{jx} \rangle + \mu_{1m}(t) \langle u_{mx}^{(k)}(t), \varphi_{jx} \rangle = \langle F_{1m}(t), \varphi_j \rangle, \\ \langle \ddot{v}_m^{(k)}(t), \tilde{\varphi}_j \rangle + \mu_{2m}(t) a(v_m^{(k)}(t), \tilde{\varphi}_j) = \langle F_{2m}(t), \tilde{\varphi}_j \rangle, \\ (u_m^{(k)}(0), \dot{u}_m^{(k)}(0)) = (\tilde{u}_{0k}, \tilde{u}_{1k}), \quad (v_m^{(k)}(0), \dot{v}_m^{(k)}(0)) = (\tilde{v}_{0k}, \tilde{v}_{1k}), \end{cases}$$

$1 \leq j \leq k$, in which

$$(3.15) \quad \begin{cases} (\tilde{u}_{0k}, \tilde{u}_{1k}) = \sum_{j=1}^k (\alpha_j^{(k)}, \beta_j^{(k)})\varphi_j \rightarrow (\tilde{u}_0, \tilde{u}_1) \text{ strongly in } (H_0^1 \cap H^2) \times (H_0^1 \cap H^2), \\ (\tilde{v}_{0k}, \tilde{v}_{1k}) = \sum_{j=1}^k (\tilde{\alpha}_j^{(k)}, \tilde{\beta}_j^{(k)})\tilde{\varphi}_j \rightarrow (\tilde{v}_0, \tilde{v}_1) \text{ strongly in } (V \cap H^2) \times V. \end{cases}$$

The system of (3.14) and (3.15) can be rewritten in the form

$$(3.16) \quad \begin{cases} \ddot{c}_{mj}^{(k)}(t) + \lambda\lambda_j \dot{c}_{mj}^{(k)}(t) + \lambda_j \mu_{1m}(t) c_{mj}^{(k)}(t) = \langle F_{1m}(t), \varphi_j \rangle, \\ \ddot{d}_{mj}^{(k)}(t) + \tilde{\lambda}_j \mu_{2m}(t) d_{mj}^{(k)}(t) = \langle F_{2m}(t), \tilde{\varphi}_j \rangle, \\ (c_{mj}^{(k)}(0), d_{mj}^{(k)}(0)) = (\alpha_j^{(k)}, \tilde{\alpha}_j^{(k)}), \\ (\dot{c}_{mj}^{(k)}(0), \dot{d}_{mj}^{(k)}(0)) = (\beta_j^{(k)}, \tilde{\beta}_j^{(k)}), \quad 1 \leq j \leq k. \end{cases}$$

Using Banach's contraction principle, it is not difficult to prove that the system (3.16) has a unique solution $c_{mj}^{(k)}(t)$, $d_{mj}^{(k)}(t)$, $1 \leq j \leq k$, on the interval $[0, T]$.

Step 2. A priori estimate. We put

$$(3.17) \quad \begin{aligned} S_m^{(k)}(t) &= \|\dot{u}_m^{(k)}(t)\|^2 + \|\dot{v}_m^{(k)}(t)\|^2 + \|\dot{u}_{mx}^{(k)}(t)\|^2 + \|\dot{v}_m^{(k)}(t)\|_a^2 + \lambda \|\Delta \dot{u}_m^{(k)}(t)\|^2 \\ &\quad + \mu_{1m}(t) (\|u_{mx}^{(k)}(t)\|^2 + \|\Delta u_m^{(k)}(t)\|^2) \\ &\quad + \mu_{2m}(t) (\|v_m^{(k)}(t)\|_a^2 + \|\Delta v_m^{(k)}(t)\|^2) \\ &\quad + 2\lambda \int_0^t (\|\dot{u}_{mx}^{(k)}(s)\|^2 + \|\Delta \dot{u}_m^{(k)}(s)\|^2) ds \\ &\quad + 2 \int_0^t \|\ddot{u}_{mx}^{(k)}(s)\|^2 ds + \int_0^t \|\ddot{v}_m^{(k)}(s)\|^2 ds. \end{aligned}$$

Then, it follows from (3.14), (3.17) that

$$\begin{aligned}
(3.18) \quad S_m^{(k)}(t) &= S_m^{(k)}(0) + 2\langle F_{1m}(0), \Delta\tilde{u}_{0k} + \Delta\tilde{u}_{1k} \rangle + 2\langle F_{2m}(0), \Delta\tilde{v}_{0k} \rangle \\
&\quad + 2\mu_{1m}(0)\langle \Delta\tilde{u}_{0k}, \Delta\tilde{u}_{1k} \rangle + 2\int_0^t \mu_{1m}(s)\|\Delta\dot{u}_m^{(k)}(s)\|^2 ds \\
&\quad + \int_0^t \mu'_{1m}(s)(\|u_{mx}^{(k)}(s)\|^2 + \|\Delta u_m^{(k)}(s)\|^2 + 2\langle \Delta u_m^{(k)}(s), \Delta\dot{u}_m^{(k)}(s) \rangle) ds \\
&\quad + \int_0^t \mu'_{2m}(s)(\|v_m^{(k)}(s)\|_a^2 + \|\Delta v_m^{(k)}(s)\|^2) ds \\
&\quad + 2\int_0^t (\langle F_{1m}(s), \dot{u}_m^{(k)}(s) \rangle + \langle F_{2m}(s), \dot{v}_m^{(k)}(s) \rangle) ds \\
&\quad + 2\int_0^t (\langle F'_{1m}(s), \Delta u_m^{(k)}(s) + \Delta\dot{u}_m^{(k)}(s) \rangle + \langle F'_{2m}(s), \Delta v_m^{(k)}(s) \rangle) ds \\
&\quad - 2\langle F_{1m}(t), \Delta u_m^{(k)}(t) + \Delta\dot{u}_m^{(k)}(t) \rangle - 2\langle F_{2m}(t), \Delta v_m^{(k)}(t) \rangle \\
&\quad - 2\mu_{1m}(t)\langle \Delta u_m^{(k)}(t), \Delta\dot{u}_m^{(k)}(t) \rangle + \int_0^t \|\ddot{v}_m^{(k)}(s)\|^2 ds \\
&= S_m^{(k)}(0) + 2\langle F_{1m}(0), \Delta\tilde{u}_{0k} + \Delta\tilde{u}_{1k} \rangle + 2\langle F_{2m}(0), \Delta\tilde{v}_{0k} \rangle \\
&\quad + 2\mu_{1m}(0)\langle \Delta\tilde{u}_{0k}, \Delta\tilde{u}_{1k} \rangle + \sum_{j=1}^9 I_j.
\end{aligned}$$

□

First, we need the following lemma whose proof is easy, hence we omit the details.

Lemma 3.2. *We have*

- (i) $|\mu_{im}(t)| \leq \tilde{K}_M, i = 1, 2,$
- (ii) $|\mu'_{1m}(t)| \leq \Psi_m(M, t),$
- (iii) $|\mu'_{2m}(t)| \leq \tilde{K}_M(1 + 4M^2),$
- (iv) $|F_{im}(x, t)| \leq K_M, i = 1, 2,$
- (v) $\|F'_{im}(t)\| \leq \Phi_m(M, t), i = 1, 2,$
- (vi) $\|F_{im}(t)\| \leq \|F_{im}(0)\| + \sqrt{T}K_M((1 + 4M)\sqrt{T} + 2M), i = 1, 2,$

where

$$\begin{aligned}
\Psi_m(M, t) &= \tilde{K}_M(1 + M^2 + M\|\nabla u''_{m-1}(t)\|), \\
\Phi_m(M, t) &= K_M(1 + 4M + \|u''_{m-1}(t)\| + \|v''_{m-1}(t)\|)
\end{aligned}$$

satisfy

- (vii) $\Psi_m(M, \cdot), \Phi_m(M, \cdot) \in L^2(0, T),$
- (viii) $\|\Psi_m(M, \cdot)\|_{L^2(0, T)} \leq \tilde{K}_M((1 + M^2)\sqrt{T} + M^2),$
- (ix) $\|\Phi_m(M, \cdot)\|_{L^2(0, T)} \leq K_M((1 + 4M)\sqrt{T} + 2M).$

Using Lemma 3.2 we estimate the terms I_j on the right-hand side of (3.18) as follows. Put $\lambda_* = \min\{\lambda, \mu_{1*}, \mu_{2*}\}$, by the inequality

$$(3.19) \quad \begin{aligned} S_m^{(k)}(t) &\geq \|\dot{u}_m^{(k)}(t)\|^2 + \|\dot{v}_m^{(k)}(t)\|^2 \\ &\quad + \lambda_* (\|\Delta \dot{u}_m^{(k)}(t)\|^2 + \|u_{mx}^{(k)}(t)\|^2 + \|\Delta u_m^{(k)}(t)\|^2 \\ &\quad + \|v_m^{(k)}(t)\|_a^2 + \|\Delta v_m^{(k)}(t)\|^2) \end{aligned}$$

we estimate I_1, I_2, I_3, I_4, I_5 , respectively, as

$$(3.20) \quad \begin{aligned} I_1 &= 2 \int_0^t \mu_{1m}(s) \|\Delta \dot{u}_m^{(k)}(s)\|^2 ds \leq \frac{2\tilde{K}_M}{\lambda_*} \int_0^t S_m^{(k)}(s) ds, \\ I_2 &= \int_0^t \mu'_{1m}(s) (\|u_{mx}^{(k)}(s)\|^2 + \|\Delta u_m^{(k)}(s)\|^2 + 2\langle \Delta u_m^{(k)}(s), \Delta \dot{u}_m^{(k)}(s) \rangle) ds \\ &\leq \frac{2}{\lambda_*} \int_0^t \Psi_m(M, s) S_m^{(k)}(s) ds, \\ I_3 &= \int_0^t \mu'_{2m}(s) (\|v_m^{(k)}(s)\|_a^2 + \|\Delta v_m^{(k)}(s)\|^2) ds \leq \frac{\tilde{K}_M(1+4M^2)}{\lambda_*} \int_0^t S_m^{(k)}(s) ds, \\ I_4 &= 2 \int_0^t (\langle F_{1m}(s), \dot{u}_m^{(k)}(s) \rangle + \langle F_{2m}(s), \dot{v}_m^{(k)}(s) \rangle) ds \leq 2TK_M^2 + \int_0^t S_m^{(k)}(s) ds, \\ I_5 &= 2 \int_0^t (\langle F'_{1m}(s), \Delta u_m^{(k)}(s) + \Delta \dot{u}_m^{(k)}(s) \rangle + \langle F'_{2m}(s), \Delta v_m^{(k)}(s) \rangle) ds \\ &\leq 2 \int_0^t (\|F'_{1m}(s)\| (\|\Delta u_m^{(k)}(s)\| + \|\Delta \dot{u}_m^{(k)}(s)\|) + \|F'_{2m}(s)\| \|\Delta v_m^{(k)}(s)\|) ds \\ &\leq 2\sqrt{\frac{3}{\lambda_*}} \int_0^t \Phi_m(M, s) \sqrt{S_m^{(k)}(s)} ds \\ &\leq \frac{3}{\lambda_*} \int_0^t \Phi_m(M, s) ds + \int_0^t \Phi_m(M, s) S_m^{(k)}(s) ds. \end{aligned}$$

Estimate of I_6 . By Lemma 3.2(v) and the inequality $2ab \leq \frac{1}{6}a^2 + 6b^2$ for all $a, b \geq 0$, we get

$$(3.21) \quad \begin{aligned} I_6 &= -2\langle F_{1m}(t), \Delta u_m^{(k)}(t) + \Delta \dot{u}_m^{(k)}(t) \rangle \\ &\leq 2\sqrt{\frac{2}{\lambda_*}} \|F_{1m}(t)\| \sqrt{S_m^{(k)}(t)} \\ &\leq \frac{1}{6} S_m^{(k)}(t) + \frac{12}{\lambda_*} \|F_{1m}(t)\|^2 \\ &\leq \frac{1}{6} S_m^{(k)}(t) + \frac{24}{\lambda_*} \|F_{1m}(0)\|^2 + \frac{24}{\lambda_*} TK_M^2 ((1+4M)\sqrt{T} + 2M)^2. \end{aligned}$$

Estimate of I_7 . Similarly,

$$\begin{aligned}
(3.22) \quad I_7 &= -2\langle F_{2m}(t), \Delta v_m^{(k)}(t) \rangle \\
&\leq \frac{\lambda_*}{6} \|\Delta v_m^{(k)}(t)\|^2 + \frac{6}{\lambda_*} \|F_{2m}(t)\|^2 \\
&\leq \frac{1}{6} S_m^{(k)}(t) + \frac{12}{\lambda_*} \|F_{2m}(0)\|^2 + \frac{12}{\lambda_*} TK_M^2((1+4M)\sqrt{T} + 2M)^2.
\end{aligned}$$

In order to estimate the term $I_8 = -2\mu_{1m}(t)\langle \Delta u_m^{(k)}(t), \Delta \dot{u}_m^{(k)}(t) \rangle$, we need the following lemma.

Lemma 3.3. *The term $\|\mu_{1m}(t)\Delta u_m^{(k)}(t)\|^2$ is estimated as*

$$(3.23) \quad \|\mu_{1m}(t)\Delta u_m^{(k)}(t)\|^2 \leq 2\|\mu_{1m}(0)\Delta \tilde{u}_{0k}\|^2 + \frac{2T}{\lambda_*} \int_0^t \Psi_m^2(M, s) S_m^{(k)}(s) \, ds.$$

Proof. First, we need to estimate $\|\frac{\partial}{\partial t}(\mu_{1m}(t)\Delta u_m^{(k)}(t))\|$. By Lemma 3.2(ii) and the inequality (3.19), we obtain

$$\begin{aligned}
(3.24) \quad \left\| \frac{\partial}{\partial t}(\mu_{1m}(t)\Delta u_m^{(k)}(t)) \right\| &\leq \|\mu'_{1m}(t)\Delta u_m^{(k)}(t)\| + \|\mu_{1m}(t)\Delta \dot{u}_m^{(k)}(t)\| \\
&\leq \Psi_m(M, t)(\|\Delta u_m^{(k)}(t)\| + \|\Delta \dot{u}_m^{(k)}(t)\|) \\
&\leq \sqrt{\frac{2}{\lambda_*}} \Psi_m(M, t) \sqrt{S_m^{(k)}(t)}.
\end{aligned}$$

The formula

$$\mu_{1m}(t)\Delta u_m^{(k)}(t) = \mu_{1m}(0)\Delta \tilde{u}_{0k} + \int_0^t \frac{\partial}{\partial s}(\mu_{1m}(s)\Delta u_m^{(k)}(s)) \, ds$$

implies that

$$\begin{aligned}
(3.25) \quad \|\mu_{1m}(t)\Delta u_m^{(k)}(t)\|^2 &\leq \left(\|\mu_{1m}(0)\Delta \tilde{u}_{0k}\| + \int_0^t \left\| \frac{\partial}{\partial s}(\mu_{1m}(s)\Delta u_m^{(k)}(s)) \right\| \, ds \right)^2 \\
&\leq 2\|\mu_{1m}(0)\Delta \tilde{u}_{0k}\|^2 + \frac{2T}{\lambda_*} \int_0^t \Psi_m^2(M, s) S_m^{(k)}(s) \, ds.
\end{aligned}$$

Lemma 3.3 is proved completely. \square

Using Lemma 3.3, the term I_8 is estimated as

$$\begin{aligned}
(3.26) \quad I_8 &= -2\mu_{1m}(t)\langle \Delta u_m^{(k)}(t), \Delta \dot{u}_m^{(k)}(t) \rangle \\
&\leq \frac{\lambda_*}{6} \|\Delta \dot{u}_m^{(k)}(t)\|^2 + \frac{6}{\lambda_*} \|\mu_{1m}(t)\Delta u_m^{(k)}(t)\|^2 \\
&\leq \frac{1}{6} S_m^{(k)}(t) + \frac{12}{\lambda_*} \|\mu_{1m}(0)\Delta \tilde{u}_{0k}\|^2 + \frac{12T}{\lambda_*^2} \int_0^t \Psi_m^2(M, s) S_m^{(k)}(s) \, ds.
\end{aligned}$$

Finally, we estimate $I_9 = \int_0^t \|\ddot{v}_m^{(k)}(s)\|^2 \, ds$ as follows. Equation (3.14)₂ can be rewritten as

$$\langle \ddot{v}_m^{(k)}(t), \tilde{\varphi}_j \rangle - \mu_{2m}(t)\langle \Delta v_m^{(k)}(t), \tilde{\varphi}_j \rangle = \langle F_{2m}(t), \tilde{\varphi}_j \rangle, \quad 1 \leq j \leq k.$$

Then, it follows after replacing $\tilde{\varphi}_j$ with $\ddot{v}_m^{(k)}(t)$ and integrating that

$$\begin{aligned}
(3.27) \quad I_9 &= \int_0^t \|\ddot{v}_m^{(k)}(s)\|^2 \, ds \\
&\leq 2 \int_0^t \mu_{2m}^2(s) \|\Delta v_m^{(k)}(s)\|^2 \, ds + 2 \int_0^t \|F_{2m}(s)\|^2 \, ds \\
&\leq 2\tilde{K}_M \int_0^t S_m^{(k)}(s) \, ds + 2TK_M^2.
\end{aligned}$$

It follows from (3.18), (3.20), (3.21), (3.22), (3.26) and (3.27) that

$$(3.28) \quad S_m^{(k)}(t) \leq \bar{S}_{m,k} + \sqrt{T}D_1(M) + \int_0^t \eta_m(M, s) S_m^{(k)}(s) \, ds,$$

where

$$\begin{aligned}
(3.29) \quad \bar{S}_{m,k} &= 2S_m^{(k)}(0) + 4\langle F_{1m}(0), \Delta \tilde{u}_{0k} + \Delta \tilde{u}_{1k} \rangle \\
&\quad + 4\langle F_{2m}(0), \Delta \tilde{v}_{0k} \rangle + 4\mu_{1m}(0)\langle \Delta \tilde{u}_{0k}, \Delta \tilde{u}_{1k} \rangle \\
&\quad + \frac{24}{\lambda_*} (\|\mu_{1m}(0)\Delta \tilde{u}_{0k}\|^2 + 2\|F_{1m}(0)\|^2 + \|F_{2m}(0)\|^2), \\
D_1(M) &= 8\left(1 + \frac{9}{\lambda_*} ((1 + 4M)\sqrt{T^*} + 2M)^2\right) \sqrt{T^*} K_M^2 \\
&\quad + \frac{6}{\lambda_*} ((1 + 4M)\sqrt{T^*} + 2M) K_M, \\
\eta_m(M, t) &= D_2(M) + 2\Phi_m(M, t) + \frac{4}{\lambda_*} \Psi_m(M, t) + \frac{24T}{\lambda_*^2} \Psi_m^2(M, t), \\
D_2(M) &= 2 + 2\left(2 + \frac{3 + 4M^2}{\lambda_*}\right) \tilde{K}_M.
\end{aligned}$$

On the other hand, the functions $F_{im}(x, 0) = f_i(x, 0, \tilde{u}_0(x), \tilde{v}_0(x), \tilde{u}_{0x}(x), \tilde{v}_{0x}(x), \tilde{u}_1(x), \tilde{v}_1(x))$, $\mu_{1m}(0) = \mu_1(0, \langle \tilde{u}_{0kx}, \tilde{u}_{1kx} \rangle)$ and $\mu_{2m}(0) = \mu_2(0, \|\tilde{v}_0\|^2, \|\tilde{v}_{0x}\|^2)$ are independent of m and the constant $\bar{S}_{m,k}$ is also independent of m , because

$$(3.30) \quad \begin{aligned} \bar{S}_{m,k} &= 2\|\tilde{u}_{1k}\|^2 + 2\|\tilde{v}_{1k}\|^2 + 2\|\tilde{u}_{1kx}\|^2 + 2\|\tilde{v}_{1k}\|_a^2 + 2\lambda\|\Delta\tilde{u}_{1k}\|^2 \\ &\quad + 2\mu_{1m}(0)(\|\tilde{u}_{0kx}\|^2 + \|\Delta\tilde{u}_{0k}\|^2) + 2\mu_{2m}(0)(\|\tilde{v}_{0k}\|_a^2 + \|\Delta\tilde{v}_{0k}\|^2) \\ &\quad + 4\langle F_{1m}(0), \Delta\tilde{u}_{0k} + \Delta\tilde{u}_{1k} \rangle + 4\langle F_{2m}(0), \Delta\tilde{v}_{0k} \rangle + 4\mu_{1m}(0)\langle \Delta\tilde{u}_{0k}, \Delta\tilde{u}_{1k} \rangle \\ &\quad + \frac{24}{\lambda_*}(\|\mu_{1m}(0)\Delta\tilde{u}_{0k}\|^2 + 2\|F_{1m}(0)\|^2 + \|F_{2m}(0)\|^2). \end{aligned}$$

By means of the convergences in (3.15), there exists a constant $M > 0$, independent of k and m , such that

$$(3.31) \quad \bar{S}_{m,k} \leq \frac{1}{2}M^2 \quad \forall m, k \in \mathbb{N}.$$

Now, we need the following lemmas.

Lemma 3.4. *We have*

- (i) $\eta_m(M, \cdot) \in L^1(0, T)$,
- (ii) $\int_0^T \eta_m(M, s) \, ds \leq \eta_M(T)$,

where

$$(3.32) \quad \begin{aligned} \eta_M(T) &= TD_2(M) + 2\sqrt{T}K_M((1 + 4M)\sqrt{T} + 2M) \\ &\quad + \frac{4}{\lambda_*}\sqrt{T}\tilde{K}_M((1 + M^2)\sqrt{T} + M^2) + \frac{24T}{\lambda_*^2}\tilde{K}_M^2((1 + M^2)\sqrt{T} + M^2)^2. \end{aligned}$$

Lemma 3.5. *For every $T \in (0, T^*]$ and $\delta > 0$, we put*

$$(3.33) \quad \begin{aligned} \bar{\gamma}_M(\delta, T) &= \left(3 + \frac{1}{\delta} + \frac{\tilde{K}_M(1 + 4M^2)}{\mu_*}\right)T + \frac{1}{\mu_*}\sqrt{T}\tilde{K}_M((1 + M^2)\sqrt{T} + M^2), \\ \theta_m(M, \delta, t) &= 3 + \frac{1}{\delta} + \frac{\tilde{K}_M(1 + 4M^2) + \Psi_{m+1}(M, t)}{\mu_*}. \end{aligned}$$

Then

$$(3.34) \quad \int_0^T \theta_m(M, \delta, t) \, dt \leq \bar{\gamma}_M(\delta, T).$$

The proofs of these lemmas are easy, hence we omit the details. □

Lemma 3.6. For every $T \in (0, T^*]$ and $\delta > 0$, we put

$$(3.35) \quad k_T(\delta) = \left(2 + \frac{1 + 2\sqrt{2}}{\sqrt{2\lambda_*}}\right) \sqrt{D_M(\delta, T) \exp(\overline{\gamma}_M(\delta, T))},$$

where

$$(3.36) \quad D_M(\delta, T) = 8TK_M^2 + (17T + \delta)M^4\tilde{K}_M^2.$$

Let $\delta > 0$ be such that

$$(3.37) \quad \left(2 + \frac{1 + 2\sqrt{2}}{\sqrt{2\lambda_*}}\right) M^2 \tilde{K}_M \sqrt{\delta} < 1.$$

Then, we can choose $T \in (0, T^*]$ such that

- (i) $(\frac{1}{2}M^2 + \sqrt{T}D_1(M)) \exp(\eta_M(T)) \leq M^2$,
- (ii) $k_T(\delta) < 1$.

Proof. The proof of Lemma 3.6 is easy, because

$$\lim_{T \rightarrow 0_+} \left(\frac{M^2}{2} + \sqrt{T}D_1(M)\right) \exp(\eta_M(T)) = \frac{M^2}{2} < M^2$$

and

$$\lim_{T \rightarrow 0_+} k_T(\delta) = \left(2 + \frac{2}{\sqrt{\lambda_*}} + \frac{1}{\sqrt{2\lambda_*}}\right) M^2 \tilde{K}_M \sqrt{\delta} < 1.$$

□

By (3.28), (3.31) and Lemma 3.6 (i)–(ii) we obtain

$$(3.38) \quad S_m^{(k)}(t) \leq M^2 \exp(-\eta_M(T)) + \int_0^t \eta_m(M, s) S_m^{(k)}(s) ds.$$

Using Gronwall's lemma, we deduce from Lemma 3.4 (ii) and (3.38) that

$$(3.39) \quad \begin{aligned} S_m^{(k)}(t) &\leq M^2 \exp(-\eta_M(T)) \exp\left(\int_0^t \eta_m(M, s) ds\right) \\ &\leq M^2 \exp(-\eta_M(T)) \exp(\eta_M(T)) \leq M^2 \end{aligned}$$

for all $t \in [0, T]$ and for all m and k . Therefore, we have

$$(3.40) \quad (u_m^{(k)}, v_m^{(k)}) \in W(M, T) \quad \forall m \text{ and } k.$$

Step 3. Limit process. From (3.40), we deduce the existence of a subsequence of $\{(u_m^{(k)}, v_m^{(k)})\}$ denoted identically and such that

$$(3.41) \quad \begin{cases} (u_m^{(k)}, v_m^{(k)}) \rightarrow (u_m, v_m) & \text{in } L^\infty(0, T; (H_0^1 \cap H^2) \times (V \cap H^2)) \text{ weak}^*, \\ (\dot{u}_m^{(k)}, \dot{v}_m^{(k)}) \rightarrow (u'_m, v'_m) & \text{in } L^\infty(0, T; (H_0^1 \cap H^2) \times V) \text{ weak}^*, \\ (\ddot{u}_m^{(k)}, \ddot{v}_m^{(k)}) \rightarrow (u''_m, v''_m) & \text{in } L^2(0, T; H_0^1 \times L^2) \text{ weak}, \\ (u_m, v_m) \in W(M, T). \end{cases}$$

Passing to limit in (3.14), we have (u_m, v_m) satisfying (3.10)–(3.12) in $L^2(0, T)$. On the other hand, it follows from (3.10) and (3.41)₄ that

$$\begin{aligned} u''_m &= \lambda \Delta u'_m + \mu_{1m}(t) \Delta u_m + F_{1m} \in L^\infty(0, T; L^2), \\ v''_m &= \mu_{2m}(t) \Delta v_m + F_{2m} \in L^\infty(0, T; L^2), \end{aligned}$$

hence $(u_m, v_m) \in W_1(M, T)$ and the proof of Theorem 3.1 is complete. \square

We note that

$$(3.42) \quad W_1(T) = \{(u, v) \in L^\infty(0, T; H_0^1 \times V) : \begin{aligned} &u' \in L^\infty(0, T; L^2) \cap L^2(0, T; H_0^1), \\ &v' \in L^\infty(0, T; L^2) \end{aligned}\}$$

is a Banach space with respect to the norm (see Lions [23])

$$(3.43) \quad \begin{aligned} \|(u, v)\|_{W_1(T)} &= \|u\|_{L^\infty(0, T; H_0^1)} + \|v\|_{L^\infty(0, T; V)} + \|u'\|_{L^\infty(0, T; L^2)} \\ &\quad + \|u'\|_{L^2(0, T; H_0^1)} + \|v'\|_{L^\infty(0, T; L^2)}. \end{aligned}$$

We use the result obtained in Theorem 3.1 and the compact imbedding theorems to prove the existence and uniqueness of a weak solution of the problem (1.1)–(1.3). Hence, we get the main result in this section as follows.

Theorem 3.7. *Let (H₁)–(H₃) hold. Then*

- (i) *the problem (1.1)–(1.3) has a unique weak solution $(u, v) \in W_1(M, T)$, where the constants $M > 0$ and $T > 0$ are chosen as in Theorem 3.1.*

Furthermore,

- (ii) *the linear recurrent sequence $\{(u_m, v_m)\}$ defined by (3.9)–(3.12) converges to the weak solution (u, v) of the problem (1.1)–(1.3) strongly in the space $W_1(T)$ and we have the estimate*

$$(3.44) \quad \|(u_m, v_m) - (u, v)\|_{W_1(T)} \leq C_T k_T^m \quad \forall m \in \mathbb{N},$$

where the constant $k_T = k_T(\delta) \in [0, 1)$ is defined as in Lemma 3.6 and C_T is a constant independent of m .

Proof. (a) Existence of the solution. We prove that $\{(u_m, v_m)\}$ is a Cauchy sequence in $W_1(T)$. Let $w_m = u_{m+1} - u_m$ and $\bar{w}_m = v_{m+1} - v_m$. Then (w_m, \bar{w}_m) satisfies the variational problem

$$(3.45) \quad \begin{cases} \langle w_m''(t), \varphi \rangle + \lambda \langle w_{mx}'(t), \varphi_x \rangle + \mu_{1,m+1}(t) \langle w_{mx}(t), \varphi_x \rangle \\ \quad = (\mu_{1,m+1}(t) - \mu_{1m}(t)) \langle \Delta u_m(t), \varphi \rangle + \langle F_{1,m+1}(t) - F_{1m}(t), \varphi \rangle \quad \forall \varphi \in H_0^1, \\ \langle \bar{w}_m''(t), \tilde{\varphi} \rangle + \mu_{2,m+1}(t) a(\bar{w}_m(t), \tilde{\varphi}) \\ \quad = (\mu_{2,m+1}(t) - \mu_{2m}(t)) \langle \Delta v_m(t), \tilde{\varphi} \rangle + \langle F_{2,m+1}(t) - F_{2m}(t), \tilde{\varphi} \rangle \quad \forall \tilde{\varphi} \in V, \\ w_m(0) = w_m'(0) = \bar{w}_m(0) = \bar{w}_m'(0) = 0, \end{cases}$$

where

$$(3.46) \quad \begin{aligned} F_{i,m+1}(t) - F_{im}(t) &= f_i[u_m, v_m](t) - f_i[u_{m-1}, v_{m-1}](t), \quad i = 1, 2, \\ \mu_{1,m+1}(t) - \mu_{1m}(t) &= \mu_1[u_m](t) - \mu_1[u_{m-1}](t), \\ \mu_{2,m+1}(t) - \mu_{2m}(t) &= \mu_2[v_m](t) - \mu_2[v_{m-1}](t). \end{aligned}$$

Taking $(\varphi, \tilde{\varphi}) = (w_m'(t), \bar{w}_m'(t))$ in (3.45), after integrating in t we get

$$(3.47) \quad \begin{aligned} S_m(t) &= \int_0^t (\mu'_{1,m+1}(s) \|w_{mx}(s)\|^2 + \mu'_{2,m+1}(s) \|\bar{w}_m(s)\|_a^2) ds \\ &\quad + 2 \int_0^t (\langle F_{1,m+1}(s) - F_{1m}(s), w_m'(s) \rangle \\ &\quad \quad + \langle F_{2,m+1}(s) - F_{2m}(s), \bar{w}_m'(s) \rangle) ds \\ &\quad + 2 \int_0^t (\mu_{1,m+1}(s) - \mu_{1m}(s)) \langle \Delta u_m(s), w_m'(s) \rangle ds \\ &\quad + 2 \int_0^t (\mu_{2,m+1}(s) - \mu_{2m}(s)) \langle \Delta v_m(s), \bar{w}_m'(s) \rangle ds \\ &= J_1 + J_2 + J_3 + J_4, \end{aligned}$$

where

$$(3.48) \quad \begin{aligned} S_m(t) &= \|w_m'(t)\|^2 + \|\bar{w}_m'(t)\|^2 \\ &\quad + \mu_{1,m+1}(t) \|w_{mx}(t)\|^2 + \mu_{2,m+1}(t) \|\bar{w}_m(t)\|_a^2 \\ &\quad + 2\lambda \int_0^t \|w_{mx}'(s)\|^2 ds \end{aligned}$$

and all the integrals on the right-hand side of (3.47) are estimated as follows.

Integral J_1 . By Lemma 3.2 and the inequality

$$(3.49) \quad S_m(t) \geq \|w_m'(t)\|^2 + \|\bar{w}_m'(t)\|^2 + \mu_*(\|w_{mx}(t)\|^2 + \|\bar{w}_m(t)\|_a^2)$$

with $\mu_* = \min\{\mu_{1*}, \mu_{2*}\}$, we obtain

$$(3.50) \quad \begin{aligned} J_1 &= \int_0^t (\mu'_{1,m+1}(s) \|w_{mx}(s)\|^2 + \mu'_{2,m+1}(s) \|\bar{w}_m(s)\|_a^2) ds \\ &\leq \frac{1}{\mu_*} \int_0^t (\Psi_{m+1}(M, s) + \tilde{K}_M(1 + 4M^2)) S_m(s) ds. \end{aligned}$$

Integral J_2 . By (H₃) it is clear that

$$\begin{aligned} &\|F_{i,m+1}(t) - F_{im}(t)\| \\ &\leq K_M(2\|\nabla w_{m-1}(t)\| + 2\|\nabla \bar{w}_{m-1}(t)\| + \|w'_{m-1}(t)\| + \|\bar{w}'_{m-1}(t)\|) \\ &\leq 2K_M(\|\nabla w_{m-1}(t)\| + \|\nabla \bar{w}_{m-1}(t)\| + \|w'_{m-1}(t)\| + \|\bar{w}'_{m-1}(t)\|) \\ &\leq 2K_M\|(w_{m-1}, \bar{w}_{m-1})\|_{W_1(T)}. \end{aligned}$$

Hence

$$(3.51) \quad \begin{aligned} J_2 &= 2 \int_0^t (\langle F_{1,m+1}(s) - F_{1m}(s), w'_m(s) \rangle \\ &\quad + \langle F_{2,m+1}(s) - F_{2m}(s), \bar{w}'_m(s) \rangle) ds \\ &\leq 2 \int_0^t (\|F_{1,m+1}(s) - F_{1m}(s)\| \|w'_m(s)\| \\ &\quad + \|F_{2,m+1}(s) - F_{2m}(s)\| \|\bar{w}'_m(s)\|) ds \\ &\leq 4\sqrt{2}K_M\|(w_{m-1}, \bar{w}_{m-1})\|_{W_1(T)} \int_0^t \sqrt{S_m(s)} ds \\ &\leq 8TK_M^2\|(w_{m-1}, \bar{w}_{m-1})\|_{W_1(T)}^2 + \int_0^t S_m(s) ds. \end{aligned}$$

Integrals J_3, J_4 . By the inequalities

$$\begin{aligned} |\mu_{1,m+1}(t) - \mu_{1m}(t)| &\leq \tilde{K}_M |\langle \nabla u_m(t), \nabla u'_m(t) \rangle - \langle \nabla u_{m-1}(t), \nabla u'_{m-1}(t) \rangle| \\ &\leq \tilde{K}_M (\|\nabla w_{m-1}(t)\| \|\nabla u'_m(t)\| + \|\nabla u_{m-1}(t)\| \|\nabla w'_{m-1}(t)\|) \\ &\leq M\tilde{K}_M (\|\nabla w_{m-1}(t)\| + \|\nabla w'_{m-1}(t)\|) \\ &\leq M\tilde{K}_M (\|(w_{m-1}, \bar{w}_{m-1})\|_{W_1(T)} + \|\nabla w'_{m-1}(t)\|) \end{aligned}$$

and

$$\begin{aligned} |\mu_{2,m+1}(t) - \mu_{2m}(t)| &\leq \tilde{K}_M (|\|v_m(t)\|^2 - \|v_{m-1}(t)\|^2| + |\|\nabla v_m(t)\|^2 - \|\nabla v_{m-1}(t)\|^2|) \\ &\leq 2M\tilde{K}_M (\|\bar{w}_{m-1}(t)\| + \|\nabla \bar{w}_{m-1}(t)\|) \\ &\leq 4M\tilde{K}_M\|(w_{m-1}, \bar{w}_{m-1})\|_{W_1(T)} \end{aligned}$$

we obtain that

$$\begin{aligned}
 (3.52) \quad J_3 &= 2 \int_0^t (\mu_{1,m+1}(s) - \mu_{1m}(s)) \langle \Delta u_m(s), w'_m(s) \rangle ds \\
 &\leq 2M^2 \tilde{K}_M \int_0^t (\|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)} + \|\nabla w'_{m-1}(s)\|) \|w'_m(s)\| ds \\
 &= 2M^2 \tilde{K}_M \int_0^t \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)} \|w'_m(s)\| ds \\
 &\quad + 2M^2 \tilde{K}_M \int_0^t \|\nabla w'_{m-1}(s)\| \|w'_m(s)\| ds \\
 &\leq TM^4 \tilde{K}_M^2 \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)}^2 + \int_0^t \|w'_m(s)\|^2 ds \\
 &\quad + \delta M^4 \tilde{K}_M^2 \int_0^t \|\nabla w'_{m-1}(s)\|^2 ds + \frac{1}{\delta} \int_0^t \|w'_m(s)\|^2 ds \\
 &\leq (T + \delta) M^4 \tilde{K}_M^2 \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)}^2 + \left(1 + \frac{1}{\delta}\right) \int_0^t S_m(s) ds
 \end{aligned}$$

and

$$\begin{aligned}
 (3.53) \quad J_4 &= 2 \int_0^t (\mu_{2,m+1}(s) - \mu_{2m}(s)) \langle \Delta v_m(s), \bar{w}'_m(s) \rangle ds \\
 &\leq 8M^2 \tilde{K}_M \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)} \int_0^t \|\bar{w}'_m(s)\| ds \\
 &\leq 16TM^4 \tilde{K}_M^2 \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)}^2 + \int_0^t S_m(s) ds.
 \end{aligned}$$

Combining (3.47), (3.50)–(3.53), we obtain

$$(3.54) \quad S_m(t) \leq D_M(\delta, T) \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)}^2 + \int_0^t \theta_m(M, \delta, s) S_m(s) ds,$$

where $\theta_m(M, \delta, t)$, $D_M(\delta, T)$ are as in (3.33) and (3.36).

Using Gronwall's lemma, we deduce from (3.33), (3.34) and (3.54) that

$$\begin{aligned}
 (3.55) \quad S_m(t) &\leq D_M(\delta, T) \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)}^2 \exp\left(\int_0^T \theta_m(M, \delta, s) ds\right) \\
 &\leq D_M(\delta, T) \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)}^2 \exp(\bar{\gamma}_M(\delta, T)).
 \end{aligned}$$

Hence, we deduce from (3.43) and (3.55) that

$$(3.56) \quad \|w_m, \bar{w}_m\|_{W_1(T)} \leq k_T(\delta) \|w_{m-1}, \bar{w}_{m-1}\|_{W_1(T)} \quad \forall m \in \mathbb{N},$$

where the constant $k_T = k_T(\delta) \in [0, 1)$ is defined as in Lemma 3.6, which implies that

$$(3.57) \quad \|(u_{m+p}, v_{m+p}) - (u_m, v_m)\|_{W_1(T)} \leq \frac{M}{1 - k_T} k_T^m \quad \forall m \text{ and } p \in \mathbb{N}.$$

It follows that $\{(u_m, v_m)\}$ is a Cauchy sequence in $W_1(T)$. Then there exists $(u, v) \in W_1(T)$ such that

$$(3.58) \quad (u_m, v_m) \rightarrow (u, v) \text{ strongly in } W_1(T).$$

On the other hand, $(u_m, v_m) \in W(M, T)$, thus there exists a subsequence $\{(u_{m_j}, v_{m_j})\}$ of $\{(u_m, v_m)\}$ such that

$$(3.59) \quad \begin{cases} (u_{m_j}, v_{m_j}) \rightarrow (u, v) & \text{in } L^\infty(0, T; (H_0^1 \cap H^2) \times (V \cap H^2)) \text{ weak}^*, \\ (u'_{m_j}, v'_{m_j}) \rightarrow (u', v') & \text{in } L^\infty(0, T; (H_0^1 \cap H^2) \times V) \text{ weak}^*, \\ (u''_{m_j}, v''_{m_j}) \rightarrow (u'', v'') & \text{in } L^2(0, T; H_0^1 \times L^2) \text{ weak}, \\ (u, v) \in W(M, T). \end{cases}$$

We note that

$$(3.60) \quad \|F_{im} - f_i[u, v]\|_{L^\infty(0, T; L^2)} \leq 2K_M \|(u_{m-1}, v_{m-1}) - (u, v)\|_{W_1(T)}, \quad i = 1, 2.$$

Hence, we deduce from (3.58) and (3.60) that

$$(3.61) \quad F_{im} \rightarrow f_i[u, v] \text{ strongly in } L^\infty(0, T; L^2), \quad i = 1, 2.$$

We also note that

$$|\mu_{1m}(t) - \mu_1[u](t)| \leq M\tilde{K}_M(\|\nabla u_{m-1}(t) - \nabla u(t)\| + \|\nabla u'_{m-1}(t) - \nabla u'(t)\|).$$

Therefore,

$$\begin{aligned} \|\mu_{1m} - \mu_1[u]\|_{L^2(0, T)} &\leq M\tilde{K}_M(\sqrt{T}\|u_{m-1} - u\|_{L^\infty(0, T, H_0^1)} + \|u'_{m-1} - u'\|_{L^2(0, T, H_0^1)}) \\ &\leq (1 + \sqrt{T})M\tilde{K}_M\|(u_{m-1}, v_{m-1}) - (u, v)\|_{W_1(T)} \rightarrow 0, \end{aligned}$$

which implies that

$$(3.62) \quad \mu_{1m} \rightarrow \mu_1[u] \text{ strongly in } L^2(0, T).$$

We also have

$$\|\mu_{2m} - \mu_2[v]\|_{L^\infty(0, T)} \leq 4M\tilde{K}_M\|(u_{m-1}, v_{m-1}) - (u, v)\|_{W_1(T)} \rightarrow 0,$$

hence

$$(3.63) \quad \mu_{2m} \rightarrow \mu_2[v] \text{ strongly in } L^\infty(0, T).$$

Finally, passing to limit in (3.10)–(3.12) as $m = m_j \rightarrow \infty$, it implies from (3.58), (3.59), (3.61), (3.62) and (3.63) that there exists $(u, v) \in W(M, T)$ satisfying the system

$$(3.64) \quad \begin{cases} \langle u''(t), \varphi \rangle + \lambda \langle u'_x(t), \varphi_x \rangle + \mu_1[u](t) \langle u_x(t), \varphi_x \rangle = \langle f_1[u, v](t), \varphi \rangle, \\ \langle v''(t), \tilde{\varphi} \rangle + \mu_2[v](t) a(v(t), \tilde{\varphi}) = \langle f_2[u, v](t), \tilde{\varphi} \rangle \end{cases}$$

for all $(\varphi, \tilde{\varphi}) \in H_0^1 \times V$ and the initial conditions

$$(3.65) \quad \begin{aligned} (u(0), u'(0)) &= (\tilde{u}_0, \tilde{u}_1), \\ (v(0), v'(0)) &= (\tilde{v}_0, \tilde{v}_1). \end{aligned}$$

Furthermore, from the assumptions (H_2) and (H_3) we obtain from (3.59)₄ and (3.64), that

$$(3.66) \quad \begin{cases} u'' = \lambda \Delta u' + \mu_1[u] \Delta u + f_1[u, v] \in L^\infty(0, T; L^2), \\ v'' = \mu_2[v] \Delta v + f_2[u, v] \in L^\infty(0, T; L^2). \end{cases}$$

Thus, we have $(u, v) \in W_1(M, T)$. The existence proof is completed.

(b) Uniqueness of the solution. Let $(u, v), (\tilde{u}, \tilde{v}) \in W_1(M, T)$ be two weak solutions of the problem (1.1)–(1.3). Then $(w, \bar{w}) = (u - \tilde{u}, v - \tilde{v})$ satisfies the variational problem

$$(3.67) \quad \begin{cases} \langle w''(t), \varphi \rangle + \lambda \langle w'_x(t), \varphi_x \rangle + \mu_1(t) \langle w_x(t), \varphi_x \rangle \\ \quad = \langle F_1(t) - \tilde{F}_1(t), \varphi \rangle + [\mu_1(t) - \tilde{\mu}_1(t)] \langle \Delta \tilde{u}(t), \varphi \rangle \quad \forall \varphi \in H_0^1, \\ \langle \bar{w}''(t), \tilde{\varphi} \rangle + \mu_2(t) a(\bar{w}(t), \tilde{\varphi}) \\ \quad = \langle F_2(t) - \tilde{F}_2(t), \tilde{\varphi} \rangle + [\mu_2(t) - \tilde{\mu}_2(t)] \langle \Delta \tilde{v}(t), \tilde{\varphi} \rangle \quad \forall \tilde{\varphi} \in V, \\ w(0) = w'(0) = \bar{w}(0) = \bar{w}'(0) = 0, \end{cases}$$

where

$$(3.68) \quad \begin{cases} \mu_1(t) = \mu_1[u](t), & \tilde{\mu}_1(t) = \mu_1[\tilde{u}](t), \\ \mu_2(t) = \mu_2[v](t), & \tilde{\mu}_2(t) = \mu_2[\tilde{v}](t), \\ F_1(t) = f_1[u, v](t), & \tilde{F}_1(t) = f_1[\tilde{u}, \tilde{v}](t), \\ F_2(t) = f_2[u, v](t), & \tilde{F}_2(t) = f_2[\tilde{u}, \tilde{v}](t). \end{cases}$$

We take $(\varphi, \tilde{\varphi}) = (w'(t), \bar{w}'(t))$ in (3.67) and integrate with respect to t to get

$$(3.69) \quad \begin{aligned} Z(t) &= 2 \int_0^t (\langle F_1(s) - \tilde{F}_1(s), w'(s) \rangle + \langle F_2(s) - \tilde{F}_2(s), \bar{w}'(s) \rangle) ds \\ &\quad + \int_0^t (\mu'_1(s) \|w_x(s)\|^2 + \mu'_2(s) \|\bar{w}(s)\|_a^2) ds \\ &\quad + 2 \int_0^t (\mu_1(s) - \tilde{\mu}_1(s)) \langle \Delta \tilde{u}(s), w'(s) \rangle ds \\ &\quad + 2 \int_0^t (\mu_2(s) - \tilde{\mu}_2(s)) \langle \Delta \tilde{v}(s), \bar{w}'(s) \rangle ds, \end{aligned}$$

where

$$(3.70) \quad Z(t) = \|w'(t)\|^2 + \|\bar{w}'(t)\|^2 + \mu_1(t) \|w_x(t)\|^2 + \mu_2(t) \|\bar{w}(t)\|_a^2 + 2\lambda \int_0^t \|w'_x(s)\|^2 ds.$$

Put

$$(3.71) \quad \mathcal{H}(s) = \mathcal{H}_M + \frac{2}{\bar{\mu}_*} M \tilde{K}_M \|u''_x(s)\|,$$

where $\bar{\mu}_* = \min\{1, \mu_{1*}, \mu_{2*}\}$ and \mathcal{H}_M is a constant defined by

$$(3.72) \quad \mathcal{H}_M = \frac{16\sqrt{2}K_M}{\sqrt{\bar{\mu}_*}} + 2 \left(\frac{2 + 4\sqrt{2}}{\sqrt{\bar{\mu}_*}} + \frac{M^2 \tilde{K}_M}{\lambda} \right) M^2 \tilde{K}_M + \frac{2}{\bar{\mu}_*} (2 + 5M^2) \tilde{K}_M.$$

Then it follows from (3.69) that

$$(3.73) \quad Z(t) \leq \int_0^t \mathcal{H}(s) Z(s) ds.$$

We remark that by $u'' \in L^2(0, T; H_0^1)$, we obtain $\mathcal{H} \in L^2(0, T)$.

It follows from (3.73) that

$$(3.74) \quad Z^2(t) \leq \|\mathcal{H}\|_{L^2(0, T)}^2 \int_0^t Z^2(s) ds.$$

By Gronwall's lemma, we deduce $Z(t) \equiv 0$, i.e., $u - \tilde{u} = v - \tilde{v} = 0$. Theorem 3.7 is proved completely. \square

4. EXPONENTIAL DECAY OF SOLUTIONS

This section investigates the decay of the solution of the problem (1.1)–(1.3) corresponding to $\zeta = 0$,

$$\begin{aligned} f_1(x, t, u, v, u_x, v_x, u_t, v_t) &= -\lambda_1 u_t + f_1(u, v) + F_1(x, t), \\ f_2(x, t, u, v, u_x, v_x, u_t, v_t) &= -\lambda_2 v_t + f_2(u, v) + F_2(x, t), \\ \mu_1(t, \langle u_x(t), u_{xt}(t) \rangle) &= \mu_* + \sigma(\langle u_x(t), u_{xt}(t) \rangle), \\ \mu_2(t, \|v(t)\|^2, \|v_x(t)\|^2) &= \mu_2(\|v_x(t)\|^2). \end{aligned}$$

Then, the problem (1.1)–(1.3) becomes the problem

$$(4.1) \quad \begin{cases} u_{tt} - \lambda u_{xxt} - (\mu_* + \sigma(\langle u_x(t), u_{xt}(t) \rangle)) u_{xx} + \lambda_1 u_t \\ \qquad \qquad \qquad = f_1(u, v) + F_1(x, t), & 0 < x < 1, t > 0, \\ v_{tt} - \mu_2(\|v_x(t)\|^2) v_{xx} + \lambda_2 v_t = f_2(u, v) + F_2(x, t), & 0 < x < 1, t > 0, \\ u(0, t) = u(1, t) = v_x(0, t) = v(1, t) = 0, \\ (u(x, 0), u_t(x, 0)) = (\tilde{u}_0(x), \tilde{u}_1(x)), \\ (v(x, 0), v_t(x, 0)) = (\tilde{v}_0(x), \tilde{v}_1(x)), \end{cases}$$

where $\lambda > 0$, $\lambda_i > 0$, $\mu_* > 0$ are given constants and σ , μ_2 , $f_i(u, v)$, $F_i(x, t)$, \tilde{u}_i , \tilde{v}_i , μ_i , f_i ($i = 1, 2$) are given functions.

By the same method as in the proof of Theorem 3.1, equation (4.1) has a weak solution $u(x, t)$ such that

$$(4.2) \quad \begin{aligned} (u, v) &\in C([0, T]; (H_0^1 \cap H^2) \times V) \cap C^1([0, T]; H_0^1 \times L^2) \\ &\quad \cap L^\infty(0, T; (H_0^1 \cap H^2) \times (V \cap H^2)), \\ (u', v') &\in C([0, T]; H_0^1 \times L^2) \cap L^\infty(0, T; (H_0^1 \cap H^2) \times V), \\ (u'', v'') &\in L^\infty(0, T; H^2 \cap H_0^1) \times L^\infty(0, T; L^2) \end{aligned}$$

for $T > 0$ small enough.

We make the following assumptions.

(\bar{H}_2) $\sigma \in C^1(\mathbb{R})$ and there exists a positive constant $\sigma_* < \mu_*$ such that

- (i) $\sigma(y) \geq -\sigma_*$ for all $y \in \mathbb{R}$,
- (ii) $y\sigma(y) > 0$ for all $y \in \mathbb{R}$, $y \neq 0$.

(\bar{H}_3) $\mu_2 \in C^1(\mathbb{R}_+)$ and there exist positive constants μ_2^* , χ_* such that

- (i) $\mu_2(y) \geq \mu_2^* > 0$ for all $y \geq 0$,
- (ii) $y\mu_2(y) \geq \chi_* \int_0^y \mu_2(z) dz$ for all $y \geq 0$.

(\bar{H}_4) There exist $\mathcal{F} \in C^2(\mathbb{R}^2; \mathbb{R})$ and the constants α , β , d_1 , $\bar{d}_1 > 0$ with $\alpha > 2$, $\beta > 2$ such that

- (i) $\frac{\partial \mathcal{F}}{\partial u}(u, v) = f_1(u, v)$, $\frac{\partial \mathcal{F}}{\partial v}(u, v) = f_2(u, v)$ for all $(u, v) \in \mathbb{R}^2$,
 - (ii) $uf_1(u, v) + vf_2(u, v) \leq d_1 \mathcal{F}(u, v)$ for all $(u, v) \in \mathbb{R}^2$,
 - (iii) $\mathcal{F}(u, v) \leq \bar{d}_1(|u|^\alpha + |v|^\beta)$ for all $(u, v) \in \mathbb{R}^2$.
- ($\bar{\text{H}}_5$) $F_1, F_2 \in L^\infty(\mathbb{R}_+; L^2) \cap L^1(\mathbb{R}_+; L^2)$ and there exist two constants $C_0, \gamma_0 > 0$ such that $\|F_1(t)\|^2 + \|F_2(t)\|^2 \leq C_0 \exp(-\gamma_0 t)$ for all $t \geq 0$.
- ($\bar{\text{H}}_6$) $p > \max\{2, d_1, d_1/\chi_*\}$ and $\sigma_*/\mu_* \leq 1 - d_1/p$.

Example 4.1. Below we give an example of the functions σ, μ_2 satisfying assumptions ($\bar{\text{H}}_3$), ($\bar{\text{H}}_4$),

$$(4.3) \quad \sigma(y) = \begin{cases} \frac{-\sigma_* y}{y-1} & \text{if } y < 0, \\ (\sigma_* + |y|^{r_1-2})y & \text{if } y \geq 0, \end{cases} \quad \mu_2(y) = \mu_{2*} + y^{r_2-1},$$

where $\sigma_* > 0, \mu_{2*} > 0; r_1, r_2 > 2$ are constants.

First, we construct the Lyapunov functional

$$(4.4) \quad \mathcal{L}(t) = E(t) + \delta\psi(t), \quad t \geq 0,$$

where $\delta > 0$ will be chosen later and

$$(4.5) \quad \begin{aligned} E(t) &= \frac{1}{2}(\|u'(t)\|^2 + \|v'(t)\|^2) + \frac{1}{2}(g * u')(t) + \frac{1}{2}(g * v')(t) + \frac{\mu_*}{2}\|u_x(t)\|^2 \\ &\quad + \frac{1}{2} \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - \int_0^1 \mathcal{F}(u(x, t), v(x, t)) \, dx \\ &= \frac{1}{2}(\|u'(t)\|^2 + \|v'(t)\|^2) \\ &\quad + \left(\frac{1}{2} - \frac{1}{p}\right) \left((g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 \right. \\ &\quad \left. + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \right) + \frac{1}{p} I(t), \end{aligned}$$

$$(4.6) \quad \psi(t) = \langle u'(t), u(t) \rangle + \langle v'(t), v(t) \rangle + \frac{\lambda}{2} \|u_x(t)\|^2 + \frac{\lambda_1}{2} \|u(t)\|^2 + \frac{\lambda_2}{2} \|v(t)\|^2,$$

where

$$\begin{aligned} (g * u')(t) &= \int_0^t g(t-s) \|u'(s)\|^2 \, ds, \\ (g * v')(t) &= \int_0^t g(t-s) \|v'(s)\|^2 \, ds \end{aligned}$$

with $g(t) = 2\bar{\lambda}e^{-2\bar{k}t}$, $\bar{k}, \bar{\lambda}$ are constants with $\bar{k} > 0, 0 < \bar{\lambda} < \lambda_* = \min\{\lambda_1, \lambda_2\}$, and

$$(4.7) \quad \begin{aligned} I(t) &= (g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 \\ &\quad + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - p \int_0^1 \mathcal{F}(u(x, t), v(x, t)) \, dx, \quad t \geq 0. \end{aligned}$$

Then we have the following theorem.

Theorem 4.2. *Assume that (\bar{H}_2) – (\bar{H}_6) hold. Let $(\tilde{u}_0, \tilde{v}_0) \in (H_0^1 \cap H^2) \times (V \cap H^2)$, $(\tilde{u}_1, \tilde{v}_1) \in (H_0^1 \cap H^2) \times V$ such that $I(0) > 0$ and the initial energy $E(0)$ satisfy*

$$(4.8) \quad \eta^* = \tilde{\mu}_* - p\bar{d}_1(R_*^{\alpha-2} + R_*^{\beta-2}) > 0,$$

where $R_*^2 = 2pE_*/((p-2)\tilde{\mu}_*)$, $E_* = (E(0) + \frac{1}{2}\varrho)\exp(\varrho)$, $\varrho = \int_0^\infty (\|F_1(t)\| + \|F_1(t)\|) dt$, $\tilde{\mu}_* = \min\{\mu_*, \mu_{2*}\}$. Then, any global weak solution of the problem (4.1) is exponentially decaying, i.e., there exist positive constants \bar{C} , $\bar{\gamma}$ such that

$$(4.9) \quad \|u'(t)\|^2 + \|v'(t)\|^2 + \|u_x(t)\|^2 + \|v_x(t)\|^2 \leq \bar{C} \exp(-\bar{\gamma}t) \quad \forall t \geq 0.$$

First, to prove the theorem we need the following lemmas.

Lemma 4.3. *The energy functional $E(t)$ defined by (4.5) satisfies*

- (i) $E'(t) \leq \frac{1}{2}(\|F_1(t)\| + \|F_2(t)\|) + \frac{1}{2}(\|F_1(t)\| + \|F_2(t)\|)(\|u'(t)\|^2 + \|v'(t)\|^2)$,
- (ii) $E'(t) \leq -\lambda\|u'_x(t)\|^2 - (\lambda_* - \bar{\lambda} - \frac{1}{2}\varepsilon_1)(\|u'(t)\|^2 + \|v'(t)\|^2) - \bar{k}((g * u')(t) + (g * v')(t)) + \frac{1}{2}\varepsilon_1^{-1}(\|F_1(t)\|^2 + \|F_2(t)\|^2)$

for all $t \geq 0$, $\varepsilon_1 > 0$, where $\lambda_* = \min\{\lambda_1, \lambda_2\}$.

Proof. Multiplying (4.1)₁ by $u'(x, t)$, (4.2)₂ by $v'(x, t)$ and integrating over $[0, 1]$, we get

$$(4.10) \quad \begin{aligned} E'(t) = & -\lambda\|u'_x(t)\|^2 - (\lambda_1 - \bar{\lambda})\|u'(t)\|^2 - (\lambda_2 - \bar{\lambda})\|v'(t)\|^2 \\ & - \bar{k}((g * u')(t) + (g * v')(t)) - \langle u_x(t), u'_x(t) \rangle \sigma(\langle u_x(t), u'_x(t) \rangle) \\ & + \langle F_1(t), u'(t) \rangle + \langle F_2(t), v'(t) \rangle. \end{aligned}$$

On the other hand,

$$(4.11) \quad \begin{aligned} \langle F_1(t), u'(t) \rangle & \leq \frac{1}{2}\|F_1(t)\| + \frac{1}{2}\|F_1(t)\|\|u'(t)\|^2, \\ \langle F_2(t), v'(t) \rangle & \leq \frac{1}{2}\|F_2(t)\| + \frac{1}{2}\|F_2(t)\|\|v'(t)\|^2. \end{aligned}$$

As $\langle u_x(t), u'_x(t) \rangle \sigma(\langle u_x(t), u'_x(t) \rangle) \geq 0$, it follows from (4.10) and (4.11) that Lemma 4.3 (i) holds. Similarly,

$$(4.12) \quad \begin{aligned} \langle F_1(t), u'(t) \rangle & \leq \frac{1}{2\varepsilon_1}\|F_1(t)\|^2 + \frac{\varepsilon_1}{2}\|u'(t)\|^2, \\ \langle F_2(t), v'(t) \rangle & \leq \frac{1}{2\varepsilon_1}\|F_2(t)\|^2 + \frac{\varepsilon_1}{2}\|v'(t)\|^2 \quad \forall \varepsilon_1 > 0, \end{aligned}$$

and it follows from (4.10) and (4.12) that Lemma 4.3 (ii) holds.

Lemma 4.3 is proved. □

Lemma 4.4. Assume that (\bar{H}_2) – (\bar{H}_6) , $I(0) > 0$ and (4.8) hold. Then $I(t) > 0$ for all $t \geq 0$.

P r o o f. By the continuity of $I(t)$ and $I(0) > 0$, there exists $\tilde{T}_1 > 0$ such that

$$(4.13) \quad I(t) = I(u(t), v(t)) > 0 \quad \forall t \in [0, \tilde{T}_1],$$

which implies

$$(4.14) \quad \begin{aligned} E(t) &\geq \frac{1}{2}(\|u'(t)\|^2 + \|v'(t)\|^2) + \left(\frac{1}{2} - \frac{1}{p}\right) \left(\mu_* \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \right) \\ &\geq \frac{1}{2}(\|u'(t)\|^2 + \|v'(t)\|^2) + \frac{(p-2)\tilde{\mu}_*}{2p} (\|u_x(t)\|^2 + \|v_x(t)\|^2) \quad \forall t \in [0, \tilde{T}_1], \end{aligned}$$

where $\tilde{\mu}_* = \min\{\mu_*, \mu_{2*}\}$.

Combining Lemma 4.3 (i) and (4.14) and using Gronwall's inequality, we get

$$(4.15) \quad \begin{aligned} \|u_x(t)\|^2 + \|v_x(t)\|^2 &\leq \frac{2p}{(p-2)\tilde{\mu}_*} E(t) \leq \frac{2pE_*}{(p-2)\tilde{\mu}_*} \equiv R_*^2 \quad \forall t \in [0, \tilde{T}_1], \\ \|u'(t)\|^2 + \|v'(t)\|^2 &\leq 2E(t) \leq 2E_* \quad \forall t \in [0, \tilde{T}_1], \end{aligned}$$

where E_* is as in (4.8). Hence, it follows from (\bar{H}_4) (iii) and (4.15) that

$$(4.16) \quad \begin{aligned} p \int_0^1 \mathcal{F}(u(x, t), v(x, t)) \, dx &\leq p\bar{d}_1 (\|u(t)\|_{L^\alpha}^\alpha + \|v(t)\|_{L^\beta}^\beta) \\ &\leq p\bar{d}_1 (\|u_x(t)\|^\alpha + \|v_x(t)\|^\beta) \\ &\leq p\bar{d}_1 (\|u_x(t)\|^{\alpha-2} + \|v_x(t)\|^{\beta-2}) \\ &\quad \times (\|u_x(t)\|^2 + \|v_x(t)\|^2) \\ &\leq p\bar{d}_1 (R_*^{\alpha-2} + R_*^{\beta-2}) (\|u_x(t)\|^2 + \|v_x(t)\|^2). \end{aligned}$$

Thus,

$$(4.17) \quad \begin{aligned} I(t) &= (g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 \\ &\quad + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - p \int_0^1 \mathcal{F}(u(x, t), v(x, t)) \, dx \\ &\geq (g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 + \mu_{2*} \|v_x(t)\|^2 \\ &\quad - p\bar{d}_1 (R_*^{\alpha-2} + R_*^{\beta-2}) (\|u_x(t)\|^2 + \|v_x(t)\|^2) \\ &\geq (g * u')(t) + (g * v')(t) + \eta^* (\|u_x(t)\|^2 + \|v_x(t)\|^2) \geq 0 \quad \forall t \in [0, \tilde{T}_1], \end{aligned}$$

where $\eta^* = \tilde{\mu}_* - p\bar{d}_1 (R_*^{\alpha-2} + R_*^{\beta-2})$ and $\tilde{\mu}_* = \min\{\mu_*, \mu_{2*}\}$ as in (4.8).

Now, we prove that $I(t) > 0$ for all $t \geq 0$. We put $T_\infty = \sup\{T > 0: I(t) > 0 \text{ for all } t \in [0, T]\}$. If $T_\infty < \infty$ then, by the continuity of $I(t)$, we have $I(T_\infty) \geq 0$.

In case of $I(T_\infty) > 0$, by the same arguments as above, we can deduce that there exists $\tilde{T}_\infty > T_\infty$ such that $I(t) > 0$ for all $t \in [0, \tilde{T}_\infty]$. We obtain a contradiction to the definition of T_∞ .

In case of $I(T_\infty) = 0$, it follows from (4.17) that

$$0 = I(T_\infty) \geq (g * u')(T_\infty) + (g * v')(T_\infty) + \eta^*(\|u_x(T_\infty)\|^2 + \|v_x(T_\infty)\|^2) \geq 0.$$

Therefore

$$u(T_\infty) = v(T_\infty) = 0, \quad (g * u')(T_\infty) = (g * v')(T_\infty) = 0.$$

By the fact that the function $s \mapsto g(T_\infty - s)\|u'(s)\|^2$ is continuous on $[0, T_\infty]$ and $g(T_\infty - s) > 0$ for all $s \in [0, T_\infty]$, we have

$$(g * u')(T_\infty) = \int_0^{T_\infty} g(T_\infty - s)\|u'(s)\|^2 ds = 0,$$

which implies that $\|u'(s)\| = 0$ for all $s \in [0, T_\infty]$. It means that u is a constant function on $[0, T_\infty]$. Then $u(0) = u(T_\infty) = 0$.

Similarly, $v(0) = v(T_\infty) = 0$. It leads to $I(0) = 0$. We get a contradiction with $I(0) > 0$. Consequently, $T_\infty = \infty$, i.e. $I(t) > 0$ for all $t \geq 0$.

Lemma 4.4 is proved completely. □

Lemma 4.5. *Assume that (\overline{H}_2) – (\overline{H}_6) hold. Let $I(0) > 0$ and (4.8) hold. Put*

$$(4.18) \quad E_1(t) = (g * u')(t) + (g * v')(t) + \|u'(t)\|^2 + \|v'(t)\|^2 + \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) dz + I(t).$$

Then there exist positive constants β_1, β_2 such that

$$(4.19) \quad \beta_1 E_1(t) \leq \mathcal{L}(t) \leq \beta_2 E_1(t) \quad \forall t \geq 0$$

for δ is small enough.

Proof. It is easy to see that

$$(4.20) \quad \begin{aligned} \mathcal{L}(t) &= \frac{1}{2}(\|u'(t)\|^2 + \|v'(t)\|^2) \\ &+ \left(\frac{1}{2} - \frac{1}{p}\right) \left((g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) dz \right) \\ &+ \frac{1}{p} I(t) + \delta \langle u'(t), u(t) \rangle + \delta \langle v'(t), v(t) \rangle \\ &+ \frac{\delta}{2} (\lambda \|u_x(t)\|^2 + \lambda_1 \|u(t)\|^2 + \lambda_2 \|v(t)\|^2). \end{aligned}$$

From the inequalities

$$\begin{aligned} |\langle u'(t), u(t) \rangle| &\leq \frac{1}{2} \|u'(t)\|^2 + \frac{1}{2} \|u_x(t)\|^2, \\ |\langle v'(t), v(t) \rangle| &\leq \frac{1}{2} \|v'(t)\|^2 + \frac{1}{2} \|v_x(t)\|^2, \\ -\mu_{2*} \|v_x(t)\|^2 &\geq - \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \end{aligned}$$

we deduce that

$$\begin{aligned} (4.21) \quad \mathcal{L}(t) &\geq \frac{1}{2} (\|u'(t)\|^2 + \|v'(t)\|^2) \\ &\quad + \left(\frac{1}{2} - \frac{1}{p}\right) \left((g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \right) \\ &\quad + \frac{1}{p} I(t) - \frac{\delta}{2} (\|u'(t)\|^2 + \|u_x(t)\|^2) - \frac{\delta}{2} (\|v'(t)\|^2 + \|v_x(t)\|^2) \\ &\geq \frac{1-\delta}{2} (\|u'(t)\|^2 + \|v'(t)\|^2) \\ &\quad + \left(\frac{1}{2} - \frac{1}{p}\right) \left((g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \right) \\ &\quad + \frac{1}{p} I(t) - \frac{\delta}{2} \|u_x(t)\|^2 - \frac{\delta}{2\mu_{2*}} \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \\ &= \frac{1-\delta}{2} (\|u'(t)\|^2 + \|v'(t)\|^2) + \left(\frac{1}{2} - \frac{1}{p}\right) ((g * u')(t) + (g * v')(t)) \\ &\quad + \left(\left(\frac{1}{2} - \frac{1}{p}\right) \mu_* - \frac{\delta}{2} \right) \|u_x(t)\|^2 + \left(\frac{1}{2} - \frac{1}{p} - \frac{\delta}{2\mu_{2*}}\right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz + \frac{1}{p} I(t), \end{aligned}$$

where we choose $\beta_1 = \min\{\frac{1}{2}(1-\delta), (\frac{1}{2} - p^{-1}), (\frac{1}{2} - p^{-1})\mu_* - \frac{1}{2}\delta, (\frac{1}{2} - p^{-1} - \delta/(2\mu_{2*})), p^{-1}\}$ with δ is small enough, $0 < \delta < \min\{1; (1 - 2p^{-1})\mu_*; (1 - 2p^{-1})\mu_{2*}\}$.

Similarly, we can prove that

$$\begin{aligned} (4.22) \quad \mathcal{L}(t) &\leq \frac{1+\delta}{2} (\|u'(t)\|^2 + \|v'(t)\|^2) + \left(\frac{1}{2} - \frac{1}{p}\right) ((g * u')(t) + (g * v')(t)) \\ &\quad + \left(\left(\frac{1}{2} - \frac{1}{p}\right) \mu_* + \frac{\delta(1+\lambda+\lambda_1)}{2} \right) \|u_x(t)\|^2 \\ &\quad + \left(\frac{1}{2} - \frac{1}{p} + \frac{\delta(1+\lambda_2)}{2\mu_{2*}}\right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz + \frac{1}{p} I(t) \\ &\leq \beta_2 E_1(t), \end{aligned}$$

where $\beta_2 = \max\{\frac{1}{2}(1+\delta), (\frac{1}{2} - p^{-1})\mu_* + \frac{1}{2}\delta(1+\lambda+\lambda_1), \frac{1}{2} - p^{-1} + \delta(1+\lambda_2)/(2\mu_{2*})\}$.

Lemma 4.5 is proved completely. \square

Lemma 4.6. Assume that (\overline{H}_2) – (\overline{H}_6) hold. Let $I(0) > 0$ and (4.8) hold. Then, the functional $\psi(t)$ defined by (4.6) satisfies

$$(4.23) \quad \begin{aligned} \psi'(t) \leq & \|u'(t)\|^2 + \|v'(t)\|^2 + \frac{d_1}{p}((g * u')(t) + (g * v')(t)) \\ & - \left(\mu_* - \sigma_* - \frac{d_1 \mu_*}{p} + \frac{d_1(1 - \delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2} \right) \|u_x(t)\|^2 \\ & - \left(\chi_* - \frac{d_1}{p} + \frac{1}{\mu_{2*}} \left(\frac{d_1(1 - \delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2} \right) \right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \\ & - \frac{d_1 \delta_1}{p} I(t) + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2) \end{aligned}$$

for all $\varepsilon_2 > 0$, $\delta_1 \in (0, 1)$.

Proof. Multiplying (4.1)₁ by $u(x, t)$, (4.1)₂ by $v(x, t)$ and integrating over $[0, 1]$, we obtain

$$(4.24) \quad \begin{aligned} \psi'(t) = & \|u'(t)\|^2 + \|v'(t)\|^2 - \mu_* \|u_x(t)\|^2 \\ & - \|u_x(t)\|^2 \sigma(\langle u_x(t), u'_x(t) \rangle) - \|v_x(t)\|^2 \mu_2(\|v_x(t)\|^2) \\ & + \langle F_1(t), u(t) \rangle + \langle F_2(t), v(t) \rangle \\ & + \langle f_1(u(t), v(t)), u(t) \rangle + \langle f_2(u(t), v(t)), v(t) \rangle. \end{aligned}$$

By the inequalities

$$(4.25) \quad \begin{aligned} -\|u_x(t)\|^2 \sigma(\langle u_x(t), u'_x(t) \rangle) & \leq \sigma_* \|u_x(t)\|^2, \\ -\|v_x(t)\|^2 \mu_2(\|v_x(t)\|^2) & \leq -\chi_* \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz, \\ \langle f_1(u(t), v(t)), u(t) \rangle + \langle f_2(u(t), v(t)), v(t) \rangle & \\ & \leq d_1 \int_0^1 \mathcal{F}(u(x, t), v(x, t)) \, dx \\ & = \frac{d_1}{p} \left((g * u')(t) + (g * v')(t) \right. \\ & \quad \left. + \mu_* \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - I(t) \right), \\ I(t) & \geq \eta^* (\|u_x(t)\|^2 + \|v_x(t)\|^2), \\ \langle F_1(t), u(t) \rangle + \langle F_2(t), v(t) \rangle & \leq \frac{\varepsilon_2}{2} (\|u_x(t)\|^2 + \|v_x(t)\|^2) \\ & \quad + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2) \end{aligned}$$

for all $\varepsilon_2 > 0$, we deduce that

(4.26)

$$\begin{aligned}
\psi'(t) &\leq \|u'(t)\|^2 + \|v'(t)\|^2 - \mu_* \|u_x(t)\|^2 + \sigma_* \|u_x(t)\|^2 - \chi_* \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \\
&\quad + \frac{d_1}{p} \left((g * u')(t) + (g * v')(t) + \mu_* \|u_x(t)\|^2 + \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - I(t) \right) \\
&\quad + \frac{\varepsilon_2}{2} (\|u_x(t)\|^2 + \|v_x(t)\|^2) + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2) \\
&= \|u'(t)\|^2 + \|v'(t)\|^2 \\
&\quad + \frac{d_1}{p} ((g * u')(t) + (g * v')(t)) - \frac{d_1 \delta_1}{p} I(t) - \frac{d_1(1 - \delta_1)}{p} I(t) \\
&\quad - \left(\mu_* - \sigma_* - \frac{d_1 \mu_*}{p} \right) \|u_x(t)\|^2 - \left(\chi_* - \frac{d_1}{p} \right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \\
&\quad + \frac{\varepsilon_2}{2} (\|u_x(t)\|^2 + \|v_x(t)\|^2) + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2) \\
&\leq \|u'(t)\|^2 + \|v'(t)\|^2 + \frac{d_1}{p} ((g * u')(t) + (g * v')(t)) \\
&\quad - \left(\mu_* - \sigma_* - \frac{d_1 \mu_*}{p} \right) \|u_x(t)\|^2 - \left(\chi_* - \frac{d_1}{p} \right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - \frac{d_1 \delta_1}{p} I(t) \\
&\quad - \left(\frac{d_1(1 - \delta_1) \eta^*}{p} - \frac{\varepsilon_2}{2} \right) (\|u_x(t)\|^2 + \|v_x(t)\|^2) + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2) \\
&\leq \|u'(t)\|^2 + \|v'(t)\|^2 + \frac{d_1}{p} ((g * u')(t) + (g * v')(t)) \\
&\quad - \left(\mu_* - \sigma_* - \frac{d_1 \mu_*}{p} + \frac{d_1(1 - \delta_1) \eta^*}{p} - \frac{\varepsilon_2}{2} \right) \|u_x(t)\|^2 \\
&\quad - \left(\chi_* - \frac{d_1}{p} \right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz - \frac{d_1 \delta_1}{p} I(t) \\
&\quad - \left(\frac{d_1(1 - \delta_1) \eta^*}{p} - \frac{\varepsilon_2}{2} \right) \frac{1}{\mu_{2*}} \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2) \\
&= \|u'(t)\|^2 + \|v'(t)\|^2 + \frac{d_1}{p} ((g * u')(t) + (g * v')(t)) \\
&\quad - \left(\mu_* - \sigma_* - \frac{d_1 \mu_*}{p} + \frac{d_1(1 - \delta_1) \eta^*}{p} - \frac{\varepsilon_2}{2} \right) \|u_x(t)\|^2 \\
&\quad - \left(\chi_* - \frac{d_1}{p} + \frac{1}{\mu_{2*}} \left(\frac{d_1(1 - \delta_1) \eta^*}{p} - \frac{\varepsilon_2}{2} \right) \right) \int_0^{\|v_x(t)\|^2} \mu_2(z) \, dz \\
&\quad - \frac{d_1 \delta_1}{p} I(t) + \frac{1}{2\varepsilon_2} (\|F_1(t)\|^2 + \|F_2(t)\|^2).
\end{aligned}$$

Hence, Lemma 4.6 is proved by using some simple estimates. \square

Proof of Theorem 4.2. Now we prove Theorem 4.2. Applying the above lemmas, we have

$$\begin{aligned}
 (4.27) \quad \mathcal{L}'(t) &\leq -\left(\lambda_* - \bar{\lambda} - \frac{\varepsilon_1}{2} - \delta\right)(\|u'(t)\|^2 + \|v'(t)\|^2) \\
 &\quad - \left(\bar{k} - \frac{\delta d_1}{p}\right)((g * u')(t) + (g * v')(t)) \\
 &\quad - \delta\left(\mu_* - \sigma_* - \frac{d_1\mu_*}{p} + \frac{d_1(1-\delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2}\right)\|u_x(t)\|^2 \\
 &\quad - \delta\left(\chi_* - \frac{d_1}{p} + \frac{1}{\mu_{2*}}\left(\frac{d_1(1-\delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2}\right)\right)\int_0^{\|v_x(t)\|^2}\mu_2(z) dz \\
 &\quad - \frac{\delta d_1\delta_1}{p}I(t) + \frac{1}{2}\left(\frac{1}{\varepsilon_1} + \frac{\delta}{\varepsilon_2}\right)(\|F_1(t)\|^2 + \|F_2(t)\|^2)
 \end{aligned}$$

for all $\varepsilon_1, \varepsilon_2 > 0$, $\delta_1 \in (0, 1)$, and with δ is small enough such that $0 < \delta < \min\{1, (1 - 2/p)\mu_*, (1 - 2/p)\mu_{2*}\}$.

Because of $p > \max\{d_1, d_1/\chi_*\}$ and $\sigma_*/\mu_* \leq 1 - d_1/p$, we have

$$\begin{aligned}
 (4.28) \quad \lim_{\delta_1 \rightarrow 0_+, \varepsilon_2 \rightarrow 0_+} &\left(\mu_* - \sigma_* - \frac{d_1\mu_*}{p} + \frac{d_1(1-\delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2}\right) \\
 &= \mu_* - \sigma_* - \frac{d_1\mu_*}{p} + \frac{d_1\eta^*}{p} \\
 &= \frac{d_1\eta^*}{p} + \mu_*\left(\left(1 - \frac{d_1}{p}\right) - \frac{\sigma_*}{\mu_*}\right) > 0
 \end{aligned}$$

and

$$(4.29) \quad \lim_{\delta_1 \rightarrow 0_+, \varepsilon_2 \rightarrow 0_+} \left(\chi_* - \frac{d_1}{p} + \frac{1}{\mu_{2*}}\left(\frac{d_1(1-\delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2}\right)\right) = \chi_* - \frac{d_1}{p} + \frac{d_1\eta^*}{p\mu_{2*}} > 0.$$

Therefore, we can choose $\delta_1 \in (0, 1)$ and $\varepsilon_2 > 0$ such that

$$\begin{aligned}
 (4.30) \quad \theta_1 = \theta_1(\delta_1, \varepsilon_2) &= \mu_* - \sigma_* - \frac{d_1\mu_*}{p} + \frac{d_1(1-\delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2} > 0, \\
 \theta_2 = \theta_2(\delta_1, \varepsilon_2) &= \chi_* - \frac{d_1}{p} + \frac{1}{\mu_{2*}}\left(\frac{d_1(1-\delta_1)\eta^*}{p} - \frac{\varepsilon_2}{2}\right) > 0.
 \end{aligned}$$

Moreover, we can choose $\varepsilon_1 > 0$, $\delta > 0$ small enough so that

$$\begin{aligned}
 (4.31) \quad \theta_3 &= \lambda_* - \bar{\lambda} - \frac{\varepsilon_1}{2} - \delta > 0, \quad \theta_4 = \bar{k} - \frac{\delta d_1}{p} > 0, \\
 0 < \delta < \min\left\{1, \left(1 - \frac{2}{p}\right)\mu_*, \left(1 - \frac{2}{p}\right)\mu_{2*}\right\}.
 \end{aligned}$$

By (4.27), (4.30), (4.31), we get

$$(4.32) \quad \mathcal{L}'(t) \leq -\beta_3 E_1(t) + \bar{C}_1 e^{-\gamma_0 t} \leq -\frac{\beta_3}{\beta_2} \mathcal{L}(t) + \bar{C}_1 e^{-\gamma_0 t} \leq -\bar{\gamma} \mathcal{L}(t) + \bar{C}_1 e^{-\gamma_0 t},$$

where $\beta_3 = \min\{\theta_3, \theta_4, \delta\theta_1, \delta\theta_2, \delta d_1 \delta_1/p\}$, $0 < \bar{\gamma} < \min\{\beta_3/\beta_2, \gamma_0\}$, $\bar{C}_1 = \frac{1}{2}(1/\varepsilon_1 + \delta/\varepsilon_2)C_0$.

Therefore, we have

$$(4.33) \quad \mathcal{L}(t) \leq \left(\mathcal{L}(0) + \frac{\bar{C}_1}{\gamma_0 - \bar{\gamma}} \right) e^{-\bar{\gamma}t} = \bar{C}_2 e^{-\bar{\gamma}t}.$$

On the other hand, we have

$$(4.34) \quad E_1(t) \geq \min\{1, \mu_{2*}\} (\|u'(t)\|^2 + \|v'(t)\|^2 + \|u_x(t)\|^2 + \|v_x(t)\|^2).$$

Combining (4.33) and (4.34) we get (4.9). Theorem 4.2 is proved completely. \square

Acknowledgements. The authors wish to express their sincere thanks to the editor and the referees for their valuable comments and remarks that improved the paper.

References

- [1] *A. V. Balakrishnan, L. W. Taylor*: Distributed parameter nonlinear damping models for flight structures. Proceedings Damping 89. Report Number: WRDC-TR-89-3116 Volume II p. FDC-1. Flight Dynamics Laboratory, Chicago, 1989, 9 pages.
- [2] *R. W. Bass, D. Zes*: Spillover, nonlinearity and flexible structures. 4th NASA Workshop on Computational Control of Flexible Aerospace Systems. NASA Conference Publication 10065. NASA. Langley Research Center, Hampton, 1991, pp. 1–14.
- [3] *S. Boulaaras*: Polynomial decay rate for a new class of viscoelastic Kirchhoff equation related with Balakrishnan-Taylor dissipation and logarithmic source terms. *Alexandria Eng. J.* 59 (2020), 1059–1071. [doi](#)
- [4] *S. Boulaaras, A. Draïfa, K. Zennir*: General decay of nonlinear viscoelastic Kirchhoff equation with Balakrishnan-Taylor damping and logarithmic nonlinearity. *Math. Methods Appl. Sci.* 42 (2019), 4795–4814. [zbl](#) [MR](#) [doi](#)
- [5] *S. Boulaaras, D. Ouchenane*: General decay for a coupled Lamé system of nonlinear viscoelastic equations. *Math. Methods Appl. Sci.* 43 (2020), 1717–1735. [zbl](#) [MR](#) [doi](#)
- [6] *N. Boumaza, S. Boulaaras*: General decay for Kirchhoff type in viscoelasticity with not necessarily decreasing kernel. *Math. Methods Appl. Sci.* 41 (2018), 6050–6069. [zbl](#) [MR](#) [doi](#)
- [7] *M. M. Cavalcanti, V. N. Domingos Cavalcanti, J. S. Prates Filho, J. A. Soriano*: Existence and exponential decay for a Kirchhoff-Carrier model with viscosity. *J. Math. Anal. Appl.* 226 (1998), 40–60. [zbl](#) [MR](#) [doi](#)
- [8] *E. Emmrich, M. Thalhammer*: A class of integro-differential equations incorporating nonlinear and nonlocal damping with applications in nonlinear elastodynamics: Existence via time discretization. *Nonlinearity* 24 (2011), 2523–2546. [zbl](#) [MR](#) [doi](#)

- [9] *B. Feng, Y. H. Kang*: Decay rates for a viscoelastic wave equation with Balakrishnan-Taylor and frictional dampings. *Topol. Methods Nonlinear Anal.* *54* (2019), 321–343. [zbl](#) [MR](#) [doi](#)
- [10] *M. M. Freitas*: Pullback attractors for non-autonomous porous elastic system with nonlinear damping and sources terms. *Math. Methods Appl. Sci.* *43* (2020), 658–681. [zbl](#) [MR](#) [doi](#)
- [11] *M. M. Freitas, M. L. Santos, J. A. Langa*: Porous elastic system with nonlinear damping and sources terms. *J. Differ. Equations* *264* (2018), 2970–3051. [zbl](#) [MR](#) [doi](#)
- [12] *E. H. Gomes Tavares, M. A. Jorge Silva, V. Narciso*: Long-time dynamics of Balakrishnan-Taylor extensible beams. *J. Dyn. Differ. Equations* *32* (2020), 1157–1175. [zbl](#) [MR](#) [doi](#)
- [13] *T. G. Ha*: General decay rate estimates for viscoelastic wave equation with Balakrishnan-Taylor damping. *Z. Angew. Math. Phys.* *67* (2016), Article ID 32, 17 pages. [zbl](#) [MR](#) [doi](#)
- [14] *T. G. Ha*: Stabilization for the wave equation with variable coefficients and Balakrishnan-Taylor damping. *Taiwanese J. Math.* *21* (2017), 807–817. [zbl](#) [MR](#) [doi](#)
- [15] *T. G. Ha*: On the viscoelastic equation with Balakrishnan-Taylor damping and acoustic boundary conditions. *Evol. Equ. Control Theory* *7* (2018), 281–291. [zbl](#) [MR](#) [doi](#)
- [16] *J. Hao, Y. Hou*: Stabilization for wave equation of variable coefficients with Balakrishnan-Taylor damping and source term. *Comput. Math. Appl.* *76* (2018), 2235–2245. [zbl](#) [MR](#) [doi](#)
- [17] *J. Hao, F. Wang*: General decay rate for weak viscoelastic wave equation with Balakrishnan-Taylor damping and time-varying delay. *Comput. Math. Appl.* *78* (2019), 2632–2640. [zbl](#) [MR](#) [doi](#)
- [18] *S. Hyder Ali Muttaqi Shah*: Some helical flows of a Burgers fluid with fractional derivative. *Meccanica* *45* (2010), 143–151. [zbl](#) [MR](#) [doi](#)
- [19] *M. Jamil, C. Fetecau*: Helical flows of Maxwell fluid between coaxial cylinders with given shear stresses on the boundary. *Nonlinear Anal., Real World Appl.* *11* (2010), 4302–4311. [zbl](#) [MR](#) [doi](#)
- [20] *J.-R. Kang, M. J. Lee, S. H. Park*: Asymptotic stability for a viscoelastic problem with Balakrishnan-Taylor damping and time-varying delay. *Comput. Math. Appl.* *74* (2017), 1506–1515. [zbl](#) [MR](#) [doi](#)
- [21] *M. J. Lee, D. Kim, J. Y. Park*: General decay of solutions for Kirchhoff type containing Balakrishnan-Taylor damping with a delay and acoustic boundary conditions. *Bound. Value Probl.* *2016* (2016), Article ID 173, 21 pages. [zbl](#) [MR](#) [doi](#)
- [22] *M. J. Lee, J. Y. Park, Y. H. Kang*: Asymptotic stability of a problem with Balakrishnan-Taylor damping and a time delay. *Comput. Math. Appl.* *70* (2015), 478–487. [zbl](#) [MR](#) [doi](#)
- [23] *J. L. Lions*: Quelques méthodes de résolution des problèmes aux limites non linéaires. *Etudes mathématiques*. Dunod, Gauthier-Villars, Paris, 1969. (In French.) [zbl](#) [MR](#)
- [24] *N. T. Long, H. H. Ha, L. T. P. Ngoc, N. A. Triet*: Existence, blow-up and exponential decay estimates for a system of nonlinear viscoelastic wave equations with nonlinear boundary conditions. *Commun. Pure Appl. Anal.* *19* (2020), 455–492. [zbl](#) [MR](#) [doi](#)
- [25] *L. A. Medeiros*: On some nonlinear perturbation of Kirchhoff-Carrier operator. *Comput. Appl. Math.* *13* (1994), 225–233. [zbl](#) [MR](#)
- [26] *C. Mu, J. Ma*: On a system of nonlinear wave equations with Balakrishnan-Taylor damping. *Z. Angew. Math. Phys.* *65* (2014), 91–113. [zbl](#) [MR](#) [doi](#)
- [27] *L. T. P. Ngoc, N. H. Nhan, B. D. Nam, N. T. Long*: Existence and exponential decay of the Dirichlet problem for a nonlinear wave equation with the Balakrishnan-Taylor term. *Lith. Math. J.* *60* (2020), 225–247. [zbl](#) [MR](#) [doi](#)
- [28] *H. Qi, H. Jin*: Unsteady helical flows of a generalized Oldroyd-B fluid with fractional derivative. *Nonlinear Anal., Real World Appl.* *10* (2009), 2700–2708. [zbl](#) [MR](#) [doi](#)
- [29] *M. L. Santos, D. S. Almeida Júnior*: On porous-elastic system with localized damping. *Z. Angew. Math. Phys.* *67* (2016), Article ID 63, 18 pages. [zbl](#) [MR](#) [doi](#)

- [30] *R. E. Showalter*: Hilbert space methods for partial differential equations. Electronic Journal of Differential Equations. Monograph 1. Southwest Texas State University, San Marcos, 1994. [zbl](#) [MR](#)
- [31] *N.-e. Tatar, A. Zarái*: Exponential stability and blow up for a problem with Balakrishnan-Taylor damping. Demonstr. Math. *44* (2011), 67–90. [zbl](#) [MR](#) [doi](#)
- [32] *N.-e. Tatar, A. Zarái*: On a Kirchhoff equation with Balakrishnan-Taylor damping and source term. Dyn. Contin. Discrete Impuls. Syst., Ser. A, Math. Anal. *18* (2011), 615–627. [zbl](#) [MR](#)
- [33] *D. Tong, X. Zhang, X. Zhang*: Unsteady helical flows of a generalized Oldroyd-B fluid. J. Non-Newton. Fluid Mech. *156* (2009), 75–83. [zbl](#) [MR](#) [doi](#)
- [34] *N. A. Triet, L. T. P. Ngoc, N. T. Long*: On a nonlinear Kirchhoff-Carrier wave equation associated with Robin conditions. Nonlinear Anal., Real World Appl. *11* (2010), 3363–3388. [zbl](#) [MR](#) [doi](#)
- [35] *A. Zarái, N.-e. Tatar*: Global existence and polynomial decay for a problem with Balakrishnan-Taylor damping. Arch. Math., Brno *46* (2010), 157–176. [zbl](#) [MR](#)

Authors' addresses: *Bui Duc Nam*, Department of Mathematics and Computer Science, University of Science, 227 Nguyen Van Cu St., Dist. 5, Ho Chi Minh City, Vietnam; Vietnam National University, Ho Chi Minh City, Vietnam, and Ho Chi Minh City University of Food Industry, 140 Le Trong Tan St., Tay Thanh Ward, Tan Phu Dist., Ho Chi Minh City, Vietnam, e-mail: nambd@hufi.edu.vn; *Nguyen Huu Nhan*, Nguyen Tat Thanh University, 300A Nguyen Tat Thanh St., Dist. 4, Ho Chi Minh City, Vietnam; Department of Mathematics and Computer Science, University of Science, 227 Nguyen Van Cu St., Dist. 5, Ho Chi Minh City, Vietnam, and Vietnam National University, Ho Chi Minh City, Vietnam, e-mail: huunhandn@gmail.com; *Le Thi Phuong Ngoc*, University of Khanh Hoa, 01 Nguyen Chanh St., Nha Trang City, Vietnam; Department of Mathematics and Computer Science, University of Science, 227 Nguyen Van Cu St., Dist. 5, Ho Chi Minh City, Vietnam, and Vietnam National University, Ho Chi Minh City, Vietnam, e-mail: ngoc1966@gmail.com; *Nguyen Thanh Long* (corresponding author), Department of Mathematics and Computer Science, University of Science, 227 Nguyen Van Cu St., Dist. 5, Ho Chi Minh City, Vietnam, and Vietnam National University, Ho Chi Minh City, Vietnam, e-mail: longnt2@gmail.com.