

Josef Šlapal

Representative properties of the quasi-ordered set $F(\alpha, M)$

Czechoslovak Mathematical Journal, Vol. 34 (1984), No. 3, 390–395

Persistent URL: <http://dml.cz/dmlcz/101964>

Terms of use:

© Institute of Mathematics AS CR, 1984

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

REPRESENTATIVE PROPERTIES
OF THE QUASI-ORDERED SET $F(\alpha, M)$

JOSEF ŠLAPAL, Ostrava

(Received October 15, 1982)

In [5] V. Novák improved a result of M. Novotný in [4] proving that a set of type $F(\omega_\nu, 2, \aleph_\nu)$ is an \aleph_ν -universal quasi-ordered set. Moreover, he used the quasi-ordered set $F(\alpha, M)$ for the representation of ordered sets and showed that a set of type $F(\omega_\nu, \aleph_\nu)$ is an \aleph_ν -universal quasi-ordered set for every regular cardinal number \aleph_ν . Finally, L. Mišík [6] proved that a set of type $F(\omega_\nu, \aleph_\nu)$ is an \aleph_ν -universal quasi-ordered set for every number \aleph_ν . In this paper the above-mentioned results are improved and supplemented.

A *quasi-ordered* set is a non-empty set G together with a reflexive and transitive binary relation \leq (see for instance [1]). If, moreover, the relation \leq is antisymmetric, the set G is said to be *ordered*. A *chain* is defined as an ordered set such that we have either $x \leq y$ or $y \leq x$ for each pair of its elements x, y . By an *antichain* we understand an ordered set for which the implication $x \leq y \Rightarrow x = y$ holds for each pair of its elements x, y . Two quasi-ordered set G, G' are called *isomorphic* if there exists such a one-one mapping f of the set G onto G' that $x, y \in G, x \leq y \Leftrightarrow f(x) \leq f(y)$. A set H with a binary relation is called an *m-universal set for quasi-ordered sets* (where $m > 0$ is a cardinality) if for every quasi-ordered set G with $\text{card } G \leq m$ there exists a subset $H' \subseteq H$ isomorphic with G . An *m-universal set for ordered sets*, an *m-universal set for chains* and an *m-universal set for antichains* are defined in an analogous way. If an *m-universal set for quasi-ordered sets* is quasi-ordered, then we call it an *m-universal quasi-ordered set*.

Let us recall one important property of every quasi-ordered set. If G is a quasi-ordered set, $x, y \in G$, then put $x \equiv y$ if and only if $x \leq y, y \leq x$. Then the relation \equiv is an equivalence relation, i.e. a reflexive, symmetric and transitive binary relation, which defines a decomposition \bar{G} of G . Let $X, Y \in \bar{G}$ and put $X \leq Y$ if and only if $x \leq y$ for any $x \in X, y \in Y$. Then the set \bar{G} is an ordered set (see [1]).

Let M be a non-empty set and $\alpha > 0$ an ordinal number. Denote by $F(\alpha, M)$ the set of all sequences of type α consisting of elements of the set M together with the relation \leq defined as follows: $\{a_\lambda \mid \lambda < \alpha\} \leq \{b_\lambda \mid \lambda < \alpha\}$ if and only if there exists a strictly increasing sequence $\{\beta_\lambda \mid \lambda < \alpha\}$ of type α of ordinal numbers less than α

such that $a_\lambda = b_{\rho_\lambda}$ for every $\lambda < \alpha$. It is easy to prove that the relation \leq is reflexive and transitive so that $F(\alpha, M)$ is a quasi-ordered set. This relation, however, is in general not antisymmetric as is shown in [4]. Therefore $F(\alpha, M)$ is generally not an ordered set. If N is a set with $\text{card } N = \text{card } M$, then clearly $F(\alpha, N)$ is isomorphic with $F(\alpha, M)$ so that the type of the set $F(\alpha, M)$ depends only on the cardinality m of the set M . We denote this type by $F(\alpha, m)$. Clearly, for $\alpha < \omega_0$ the set of type $F(\alpha, m)$ is an antichain of power $m^{\text{card } \alpha}$.

If α is an ordinal number, then we denote the set of all ordinal numbers less than α ordered according to their magnitude by $W(\alpha)$. It is known that $W(\alpha)$ is a chain of type α (see [2]). Let $\{a_\lambda \mid \lambda < \alpha\}$ be a sequence of type α . Let $G = \{x \mid \text{there exists an ordinal number } \lambda < \alpha \text{ such that } a_\lambda = x\}$. For every $x \in G$ put $m_x(\{a_\lambda \mid \lambda < \alpha\}) = \text{card } \{\lambda \mid \lambda \in W(\alpha), a_\lambda = x\}$. We shall need the following two lemmas proved in [5]:

Lemma 1. *Let G be a non-empty set such that $\text{card } G \leq \aleph_\nu$. Then the elements of the set G can be written in the form of a sequence of type ω_ν , $\{a_\lambda \mid \lambda < \omega_\nu\}$, such that $m_x(\{a_\lambda \mid \lambda < \omega_\nu\}) = \aleph_\nu$ for every $x \in G$.*

Lemma 2. *Let G be a set with $\text{card } G = m$ where $2 \leq m \leq \aleph_\nu$. Let \mathcal{S} be the set of all sequences of type ω_ν , consisting of elements of the set G and such that $m_x(\{a_\lambda \mid \lambda < \omega_\nu\}) = \aleph_\nu$ for any sequence $\{a_\lambda \mid \lambda < \omega_\nu\} \in \mathcal{S}$ and any element $x \in G$. Then $\text{card } \mathcal{S} = 2^{\aleph_\nu}$.*

Let α denote a given ordinal number. If α_1 and α_2 are ordinal numbers such that $\alpha = \alpha_1 + \alpha_2$, then the number α_2 is called the *remainder of number α corresponding to the segment α_1* (see [3]). Now we shall prove the following important theorem:

Theorem 1. *Let α, β be ordinal numbers, $0 < \alpha \leq \beta$, and let m, n be cardinal numbers, $0 < m \leq n$. Let at least one of the following three assumptions hold:*

- (I) $m < n$,
- (II) $m \geq \aleph_0$,
- (III) $\alpha_2 + (\beta - \alpha) > \beta - \alpha$ for every remainder $\alpha_2 > 0$ of number α .

Then for every quasi-ordered set $F(\alpha, M)$ of type $F(\alpha, m)$ there exists a subset of a quasi-ordered set of type $F(\beta, n)$ isomorphic with $F(\alpha, M)$.

Proof. Let $F(\alpha, M), F(\beta, N)$ be quasi-ordered sets of types $F(\alpha, m), F(\beta, n)$ where $0 < \alpha \leq \beta, 0 < m \leq n$, i.e. $\text{card } M = m, \text{card } N = n$. We can suppose $M \subseteq N$ without loss of generality.

Let the assumption (I) hold. Then the set $N - M$ is non-empty. Let $x \in N - M$ be an element and for every sequence $a = \{a_\lambda \mid \lambda < \alpha\} \in F(\alpha, M)$ put $\varphi(a) = b = \{b_\lambda \mid \lambda < \beta\}$ where $\{b_\lambda \mid \lambda < \beta\}$ is a sequence defined in the following way:

$$b_\lambda = \begin{cases} a_\lambda & \text{for } \lambda < \alpha, \\ x & \text{for } \alpha \leq \lambda < \beta. \end{cases}$$

Then clearly $b \in F(\beta, N)$ and φ is a one-one mapping of $F(\alpha, M)$ onto $\Sigma = \{\varphi(a) \mid a \in F(\alpha, M)\} \subseteq F(\beta, N)$. We shall show that φ is an isomorphism of $F(\alpha, M)$ onto Σ .

Let $a = \{a_\lambda \mid \lambda < \alpha\}$, $a' = \{a'_\lambda \mid \lambda < \alpha\} \in F(\alpha, M)$, $a \leq a'$ and $\varphi(a) = b = \{b_\lambda \mid \lambda < \beta\}$, $\varphi(a') = b' = \{b'_\lambda \mid \lambda < \beta\}$. Then there exists a strictly increasing sequence $\{\gamma_\lambda \mid \lambda < \alpha\}$ of type α of ordinal numbers less than α such that $a_\lambda = a'_{\gamma_\lambda}$ for every $\lambda < \alpha$. Let us define a sequence $\{\delta_\lambda \mid \lambda < \beta\}$ of type β of ordinal numbers less than β in the following way:

$$\delta_\lambda = \begin{cases} \gamma_\lambda & \text{for } \lambda < \alpha, \\ \lambda & \text{for } \alpha \leq \lambda < \beta. \end{cases}$$

The sequence $\{\delta_\lambda \mid \lambda < \beta\}$ is strictly increasing and $b_\lambda = a_\lambda = a'_{\gamma_\lambda} = a'_{\delta_\lambda} = b'_{\delta_\lambda}$ for every $\lambda < \alpha$ and $b_\lambda = x = b'_\lambda = b = b'_{\delta_\lambda}$ for every $\alpha \leq \lambda < \beta$. Therefore $b_\lambda = b'_{\delta_\lambda}$ for every $\lambda < \beta$, i.e. $b \leq b'$. Suppose, on the contrary, that $b = \varphi(a) = \{b_\lambda \mid \lambda < \beta\}$, $b' = \varphi(a') = \{b'_\lambda \mid \lambda < \beta\} \in \Sigma$, $b \not\leq b'$. Then there exists a strictly increasing sequence $\{\delta_\lambda \mid \lambda < \beta\}$ of type β of ordinal numbers less than β such that $b_\lambda = b'_{\delta_\lambda}$ for every $\lambda < \beta$. If $\lambda < \alpha$, then $\delta_\lambda < \alpha$, for, if $\delta_{\lambda_0} \geq \alpha$ for some $\lambda_0 < \alpha$, then $b_{\lambda_0} = b'_{\delta_{\lambda_0}} = x$ which contradicts $b_{\lambda_0} = a_{\lambda_0} \in M$. Let us define the sequence $\{\gamma_\lambda \mid \lambda < \alpha\}$ such that $\gamma_\lambda = \delta_\lambda$ for every $\lambda < \alpha$. Then $\{\gamma_\lambda \mid \lambda < \alpha\}$ is a strictly increasing sequence of type α of ordinal numbers less than α and such that $a_\lambda = b_\lambda = b'_{\delta_\lambda} = b'_{\gamma_\lambda} = a'_{\gamma_\lambda}$ for every $\lambda < \alpha$, i.e. $a \leq a'$. Thus φ is an isomorphism.

Let the assumption (II) hold. Then we can suppose that the set $N - M$ is non-empty and the proof coincides with the previous one.

Let the assumption (III) hold. Let $x \in N$ be an element and let us define the mapping φ of $F(\alpha, M)$ into $F(\beta, N)$ in the same way as in the first part of the proof. Then φ is a one-one mapping of $F(\alpha, M)$ onto $\Sigma = \{\varphi(a) \mid a \in F(\alpha, M)\} \subseteq F(\beta, N)$ and we shall show that φ is an isomorphism of $F(\alpha, M)$ onto Σ . Let $a, a' \in F(\alpha, M)$, $a \leq a'$, $b = \varphi(a)$, $b' = \varphi(a')$. We are able to prove that $b \leq b'$ in the same way as in the first part of the proof. Suppose, on the contrary, that $b = \varphi(a) = \{b_\lambda \mid \lambda < \beta\}$, $b' = \varphi(a') = \{b'_\lambda \mid \lambda < \beta\} \in \Sigma$, $b \not\leq b'$. Then there exists a strictly increasing sequence $\{\delta_\lambda \mid \lambda < \beta\}$ of type β of ordinal numbers less than β such that $b_\lambda = b'_{\delta_\lambda}$ for every $\lambda < \beta$. We shall prove that $\delta_\lambda < \alpha$ for every $\lambda < \alpha$. Suppose that there exists $\lambda_0 < \alpha$ such that $\delta_{\lambda_0} \geq \alpha$. Then $\delta_{\lambda_0} \leq \delta_\lambda < \beta$ for every $\lambda_0 \leq \lambda < \beta$, i.e. the sequence $\{b_\lambda \mid \lambda_0 \leq \lambda < \beta\}$ results by omitting a set (empty or non-empty) of members of the sequence $\{b'_\lambda \mid \delta_{\lambda_0} \leq \lambda < \beta\}$. Let α_2 denote the remainder of the number α corresponding to the segment λ_0 , i.e. $\alpha = \lambda_0 + \alpha_2$. As the type of the sequence $\{b_\lambda \mid \lambda_0 \leq \lambda < \beta\}$ is $\alpha_2 + (\beta - \alpha)$ and the type of the sequence $\{b'_\lambda \mid \delta_{\lambda_0} \leq \lambda < \beta\}$ is $\leq \beta - \alpha$ we have $\alpha_2 + (\beta - \alpha) \leq \beta - \alpha$, which is a contradiction. Therefore $\delta_\lambda < \alpha$ for every $\lambda < \alpha$ and this implies, similarly as in the first part of the proof, that $a \leq a'$. Thus φ is an isomorphism and the theorem is proved.

Now we shall investigate the set $F(\alpha, M)$ as an m -universal set.

Theorem 2. *Let m be a cardinal number such that $0 < m \leq \aleph_\nu$. Then a quasi-ordered set of type $F(\omega_\nu, m)$ is an m -universal set for ordered sets.*

Proof. Let the assumptions of Theorem be true and let G be an ordered set such

that $\text{card } G \leq m$. Then there exists a one-one mapping f of G into M where M is a set with $\text{card } M = m$. Denote by \mathcal{S} the set of all subsets of M , i.e. $\mathcal{S} = \{N \mid N \subseteq M\}$, ordered by the set inclusion. If we assign to every element $x \in G$ a subset $\psi(x) = \{f(t) \mid t \leq x\} \subseteq M$, then clearly ψ is an isomorphism of G onto a certain subset $\mathcal{S}' \subseteq \mathcal{S}$ and $\text{card } N' \geq 1$ for every $N' \in \mathcal{S}'$. Since $\text{card } M = m \leq \aleph_\nu$, according to Lemma 1 it is possible to write the elements of the set M in the form of a sequence $\{b_\lambda \mid \lambda < \omega_\nu\}$ of type ω_ν such that $m_x(\{b_\lambda \mid \lambda < \omega_\nu\}) = \aleph_\nu$ for every $x \in M$. Now let us define a mapping φ of \mathcal{S}' into $F(\omega_\nu, M)$ in the same way as in the proof of Theorem 1 of [5], i.e., let us assign to every set $N' \in \mathcal{S}'$ a sequence $\varphi(N') = \{a_\lambda \mid \lambda < \omega_\nu\}$ of type ω_ν in the following way: $a_0 = b_{\mu_0}$ where μ_0 is the smallest ordinal number such that $b_{\mu_0} \in N'$; suppose that we have defined a_λ for every $\lambda < \lambda_0$ ($\lambda_0 < \omega_\nu$) and put $a_{\lambda_0} = b_{\mu_{\lambda_0}}$ where μ_{λ_0} is the smallest ordinal number with the following properties: $\mu_{\lambda_0} > \mu_\lambda$ for every $\lambda < \lambda_0$, $\mu_{\lambda_0} < \omega_\nu$, $b_{\mu_{\lambda_0}} \in N'$. In the above mentioned proof [5] it is shown that such an ordinal number always exists and that φ is an isomorphism of \mathcal{S}' onto $\Sigma = \{\varphi(N') \mid N' \in \mathcal{S}'\} \subseteq F(\omega_\nu, M)$. Hence it follows that the composite mapping $\varphi\psi$ is an isomorphism of G onto $\Sigma \subseteq F(\omega_\nu, M)$. Because the type of the set $F(\omega_\nu, M)$ is $F(\omega_\nu, m)$, the theorem is proved.

Theorem 3. *Let \aleph_ν be a regular cardinal number and let m be a cardinal number such that $0 < m \leq \aleph_\nu$. Then a quasi-ordered set of type $F(\omega_\nu, m + 1)$ is an m -universal quasi-ordered set.*

Proof. Let the assumptions of Theorem 3 be fulfilled and let G be a quasi-ordered set such that $\text{card } G \leq m$. Then $\text{card } \bar{G} \leq m$ and similarly as in the proof of Theorem 2 there exists an isomorphism ψ of the ordered set \bar{G} onto a certain subset $\mathcal{S}' \subseteq \mathcal{S}$ where \mathcal{S} is the set of all subsets of a set M with $\text{card } M = m$ ordered by the set inclusion. The definition of the mapping ψ yields that $\text{card } N' \geq 1$ for every $N' \in \mathcal{S}'$. Let $a \in M$ be an element and for every $N' \in \mathcal{S}'$ put $N'' = N' \cup \{a\}$. Then the system $\mathcal{S}'' = \{N'' \mid N' \in \mathcal{S}'\}$ is a system of sets such that $2 \leq \text{card } N'' \leq \aleph_\nu$ for every $N'' \in \mathcal{S}''$ which — ordered by the set inclusion — is isomorphic with \bar{G} . Denote by χ an isomorphism of \bar{G} onto \mathcal{S}'' . Let $\Sigma(N'')$ be the set of all sequences $\{a_\lambda \mid \lambda < \omega_\nu\}$ of type ω_ν consisting of elements of the set N'' and such that $m_x(\{a_\lambda \mid \lambda < \omega_\nu\}) = \aleph_\nu$ for every $x \in N''$. According to Lemma 2 we have $\text{card } \Sigma(N'') = 2^{\aleph_\nu}$ for every $N'' \in \mathcal{S}''$. As $\text{card } X \leq \aleph_\nu$ for every $X \in \bar{G}$ it is possible to define a one-one mapping φ_X of the set X into $\Sigma[\chi(X)]$. Finally, let us define a mapping φ of G into $F(\omega_\nu, M \cup \{a\})$ in the same way as in the proof of Theorem 3 of [5], i.e. let $\varphi(x) = \varphi_X(x)$ where $x \in X \in \bar{G}$. In [5] it is shown that φ is an isomorphism of G onto a certain subset of $F(\omega_\nu, M \cup \{a\})$. Because the type of the set $F(\omega_\nu, M \cup \{a\})$ is $F(\omega_\nu, m + 1)$, the theorem is proved.

Theorem 4. *Let m be a cardinal number such that $0 < m \leq \aleph_\nu$. Then a quasi-ordered set of type $F(\omega_\nu, m + 2)$ is an m -universal quasi-ordered set.*

Proof. Let the assumptions of Theorem 4 be fulfilled. If $m \leq \aleph_0$, then the

statement follows from Theorem 3 and Theorem 1. If $m > \aleph_0$, then we obtain the statement in the following way:

Let G be a quasi-ordered set such that $\text{card } G \leq m$. Then $\text{card } \bar{G} \leq m$ and according to Theorem 2 the ordered set \bar{G} is isomorphic with a certain subset $H \subseteq F(\omega_\nu, M)$, where $\nu > 0$ and M is a set with $\text{card } M = m$. Denote by ψ an isomorphism of \bar{G} onto H . Let $a \in M, b \in M, a \neq b$, be two elements. Let us construct the class $\Sigma[\psi(X)]$ for every element $\psi(X) = \{a_\lambda \mid \lambda < \omega_\nu\} \in H (X \in \bar{G})$ where $\Sigma[\psi(X)]$ is the set of all sequences which we obtain by inserting the sequence $\{a, b, a, b, \dots\}$ or $\{b, a, b, a, \dots\}$ of type ω_0 after every element $a_\lambda, \lambda < \omega_\nu$. Every element $\xi \in \Sigma[\psi(X)]$ belongs to the set $F(\omega_\nu, M \cup \{a, b\})$ for every $X \in \bar{G}$. For $\psi(X) \leq \psi(Y)$ and $\xi \in \Sigma[\psi(X)], \eta \in \Sigma[\psi(Y)]$ $\xi \leq \eta$ holds. Because $\text{card } \Sigma[\psi(X)] = 2^{\aleph_\nu}$ for every $X \in \bar{G}$ there exists a one-one mapping φ_X of X into $\Sigma[\psi(X)]$ for every $X \in \bar{G}$. If we define a mapping φ of G into $F(\omega_\nu, M \cup \{a, b\})$ in the same way as in the proof of Theorem of [6], i.e. $\varphi(x) = \varphi_X(x)$ for $x \in X \in \bar{G}$, then φ is an isomorphism of G onto a certain subset of $F(\omega_\nu, M \cup \{a, b\})$. Because the type of the set $F(\omega_\nu, M \cup \{a, b\})$ is $F(\omega_\nu, m + 2)$, the theorem is proved.

Now we shall deal with representations of finite chains and finite antichains by the set $F(\alpha, M)$.

Theorem 5. *If B is a chain of type ω_ν , then a quasi-ordered set of type $F(\omega_\nu, 2)$ contains a subset isomorphic with B .*

Proof. If B is a chain of type ω_ν , then we can suppose $B = W(\omega_\nu)$ without loss of generality. Let $F(\omega_\nu, M)$ be a quasi-ordered set of type $F(\omega_\nu, 2)$, where $M = \{a, b\}$. To every ordinal number $\mu \in W(\omega_\nu)$ let us assign a sequence $f(\mu) = \{c_\lambda^\mu \mid \lambda < \omega_\nu\}$ defined in the following way:

$$c_\lambda^\mu = \begin{cases} a & \text{for } \lambda < \mu, \\ b & \text{for } \mu \leq \lambda < \omega_\nu. \end{cases}$$

It is clear that $f(\mu) \in F(\omega_\nu, M)$ for every $\mu \in W(\omega_\nu)$ and that f is a one-one mapping of the chain $W(\omega_\nu)$ onto a certain subset $K \subseteq F(\omega_\nu, M)$. We shall show that f is an isomorphism of $W(\omega_\nu)$ onto K . Hence let $\mu_1, \mu_2 \in W(\omega_\nu), \mu_1 \leq \mu_2$. Then $f(\mu_1) = \{c_\lambda^{\mu_1} \mid \lambda < \omega_\nu\}, f(\mu_2) = \{c_\lambda^{\mu_2} \mid \lambda < \omega_\nu\}$ and put

$$\gamma_\lambda = \begin{cases} \lambda & \text{for } \lambda < \mu_1, \\ \mu_2 + (\lambda - \mu_1) & \text{for } \mu_1 \leq \lambda < \omega_\nu. \end{cases}$$

Then $\{\gamma_\lambda \mid \lambda < \omega_\nu\}$ is a strictly increasing sequence of ordinal numbers of type ω_ν and because $\mu_2 + (\lambda - \mu_1) < \mu_2 + (\omega_\nu - \mu_1) = \omega_\nu$ for $\mu_1 \leq \lambda < \omega_\nu$ we have $\gamma_\lambda < \omega_\nu$ for every $\lambda < \omega_\nu$. Now if $c_\lambda^{\mu_1} = a$, then $\lambda < \mu_1$ and therefore $\gamma_\lambda = \lambda < \mu_1 \leq \mu_2$. This implies $c_{\gamma_\lambda}^{\mu_2} = a$. If $c_\lambda^{\mu_1} = b$, then $\mu_1 \leq \lambda < \omega_\nu$ and therefore $\gamma_\lambda = \mu_2 + (\lambda - \mu_1) \geq \mu_2$. This implies $c_{\gamma_\lambda}^{\mu_2} = b$. Thus, $c_\lambda^{\mu_1} = c_{\gamma_\lambda}^{\mu_2}$ for every $\lambda < \omega_\nu$, i.e. $f(\mu_1) \leq f(\mu_2)$. Suppose, on the contrary, that $f(\mu_1) = \{c_\lambda^{\mu_1} \mid \lambda < \omega_\nu\} \not\leq \{c_\lambda^{\mu_2} \mid \lambda < \omega_\nu\} = f(\mu_2)$. Then there exists a strictly increasing sequence

$\{\gamma_\lambda \mid \lambda < \omega_\nu\}$ of type ω_ν of ordinal numbers less than ω_ν such that $c_\lambda^{\mu_1} = c_{\gamma_\lambda}^{\mu_2}$ for every $\lambda < \omega_\nu$. If $\lambda < \mu_1$, then $c_\lambda^{\mu_1} = a$ and therefore $c_{\gamma_\lambda}^{\mu_2} = a$ which implies $\gamma_\lambda < \mu_2$. Because $\lambda \leq \gamma_\lambda$ for every $\lambda < \omega_\nu$ we obtain $\lambda < \mu_1 \Rightarrow \lambda < \mu_2$. This implies $\mu_1 \leq \mu_2$ and the proof is complete.

Corollary. *Let m be a cardinal number such that $0 < m < \aleph_0$. Then a quasi-ordered set of type $F(\omega_0, 2)$ is an m -universal set for chains.*

Proof. Every finite chain is isomorphic with a certain subset of a chain of type ω_0 . Now the statement follows from Theorem 5 for $\nu = 0$.

Theorem 6. *Let m be a cardinal number such that $0 < m < \aleph_0$. Then a quasi-ordered set of type $F(\omega_0, 3)$ is an m -universal set for antichains.*

Proof. Let the assumptions of Theorem 6 be fulfilled. Let P be an antichain such that $\text{card } P \leq m$. Let α be an ordinal number with $\text{card } \alpha = m$. Then a set of type $F(\alpha, 2)$ is an antichain of power 2^m and thus it contains a certain subset isomorphic with P . According to Theorem 1 every set of type $F(\alpha, 2)$ is isomorphic with a certain subset of a set of type $F(\omega_0, 3)$. Thus Theorem 6 is proved.

References

- [1] *G. Birkhoff*: Lattice Theory, Third Edition, Providence, Rhode Island, 1967.
- [2] *F. Hausdorff*: Grundzüge der Mengenlehre, Leipzig, 1914.
- [3] *W. Sierpiński*: Cardinal and Ordinal Numbers, Warszawa, 1958.
- [4] *M. Novotný*: Über quasi-geordnete Mengen, Czech. Math. Journ. 9 (84) (1959), 327–333.
- [5] *V. Novák*: On universal quasi-ordered sets, Czech. Math. Journ. 15 (90) (1965), 589–595.
- [6] *L. Mišík*: About one theorem of V. Novák, Czech. Math. Journ. 15 (90) (1965), 596.

Author's address: 708 33 Ostrava-Poruba, tř. Vítězného února, ČSSR (katedra matematiky VŠB).