

Michal Greguš; Michal Greguš, Jr.

On certain singular third order eigenvalue problem

Archivum Mathematicum, Vol. 39 (2003), No. 3, 173--177

Persistent URL: <http://dml.cz/dmlcz/107863>

Terms of use:

© Masaryk University, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON CERTAIN SINGULAR THIRD ORDER EIGENVALUE PROBLEM

MICHAL GREGUŠ AND MICHAL GREGUŠ, JR.

Dedicated to Professor's Šeda 70th birthday

ABSTRACT. In this paper a singular third order eigenvalue problem is studied. The results of the paper complete the results given in the papers [3], [5].

1. We consider the third order linear differential equation in the normal form

$$(a) \quad y''' + 2A(t)y' + [A'(t) + \lambda b(t)]y = 0,$$

where $A(t) \geq 0$, $A'(t)$ and $b(t) > 0$ are continuous functions of $t \in [a, \infty)$, $a > -\infty$ and λ is a positive parameter. It is assumed that (a) is strongly nonoscillatory on $[a, \infty)$, that is (a) is nonoscillatory there for each real positive λ . By nonoscillation of (a) we mean that all of its nontrivial solutions are nonoscillatory on $[a, \infty)$.

A nontrivial solution of (a) is called oscillatory on $[a, \infty)$ if ∞ is a limit point of zeros of that solution. In the contrary case the solution is called nonoscillatory on $[a, \infty)$.

The equation (a) is said to be oscillatory on $[a, \infty)$ if it has at least one oscillatory solution on $[a, \infty)$.

The problem studied in Section 2, is to find a nontrivial (nonoscillatory) solution $y(t, \lambda)$ of (a) which satisfies either of the boundary conditions at finite points

$$(1) \quad y(a, \lambda) = y'(a, \lambda) = y(b, \lambda) = 0, \quad a < b,$$

$$(2) \quad y(a, \lambda) = y(b, \lambda) = y(c, \lambda) = 0, \quad a < b < c$$

a , b and c being any given constants, as well as the boundary condition at infinity

$$(3) \quad y(t, \lambda) = o(t[k_1 u_1(t)u_2(t) + k_2 u_2^2(t)]) \quad \text{for } t \rightarrow \infty,$$

2000 *Mathematics Subject Classification*: 34B05, 34B24, 34C10.

Key words and phrases: singular eigenvalue problem, normal form of third order differential equation, zeros of nonoscillatory solutions.

Received June 7, 2001.

together with the requirement that

$$y(t, \lambda) \neq 0$$

in a certain neighbourhood of infinity (t_0, ∞) , where $b \leq t_0 < \infty$ (or $c \leq t_0 < \infty$ in the conditions (2)), and u_1, u_2 form a fundamental set of solutions of the second order differential equation

$$(4) \quad u'' + \frac{1}{2}A(t)u = 0$$

with initial conditions $u_1(t_0) = 1, u_1'(t_0) = 0, u_2(t_0) = 0, u_2'(t_0) = 1, k_1, k_2$ are suitable constants. The motivation for this paper was given by the paper [1] of A. Elbert, T. Kusano and M. Naito for linear second order nonoscillatory differential equations.

In the paper [3] the case $A(t) < 0$ on $[a, \infty)$ was studied.

2. At the beginning of this section we introduce certain auxiliary statements on the linear third order differential equation, given in monograph [4].

Consider equation (a) and the third order differential equation

$$(a_1) \quad y''' + 2A(t)y' + [A'(t) + b(t)] = 0,$$

Lemma 1. [4, Theorem 2.41] *Let the differential equation*

$$(5) \quad y'' + 2A(t)y = 0$$

be disconjugate in (a, ∞) and let the functions $A(t), A'(t) + b(t), b(t) - A'(t)$ be positive in $[a, \infty)$. If

$$(6) \quad \int_a^\infty t^2 [b(t) - A'(t)] dt < \infty,$$

then the differential equation (a_1) is non-oscillatory in $[a, \infty)$.

Remark 1. If $A(t) \equiv 0$ on $[a, \infty)$ and (6) holds then [2, Theorem 4] equation (a_1) is non-oscillatory in $[a, \infty)$.

By Lemma 1 and Remark 1 the following lemma can be proved.

Lemma 2. *Let the differential equation (5) be disconjugated in $[a, \infty)$ and let $A(t) \geq 0, A'(t) \leq 0, b(t) > 0$ and $A'(t) + b(t) > 0$ in $[a, \infty)$. Let further $\bar{\lambda}$ be any fixed positive value of the parameter λ . If (6) holds then*

$$(7) \quad \int_a^\infty t^2 [\bar{\lambda}b(t) - A'(t)] dt < \infty,$$

and the differential equation (a) is non-oscillatory for $\lambda = \bar{\lambda}$ in $[a, \infty)$.

Lemma 3. [4, Theorem 2.31] *Let $b(t) > 0$, $A'(t)$ be continuous in $[a, \infty)$. The equation (a_1) is oscillatory in $[a, \infty)$ if and only if its adjoint equation*

$$(b_1) \quad z''' + 2A(t) + [A'(t) - b(t)]z = 0,$$

is oscillatory in $[a, \infty)$.

If we apply Lemma 3 we can formulate Theorem 2.51 [4] as follows.

Lemma 4. *Let $A(t) \geq 0$, $A'(t) - b(t) \leq 0$, $b(t) > 0$ in $[a, \infty)$ and let $\int_a^\infty b(t) dt = \infty$, then the differential equation (a_1) is oscillatory in $[a, \infty)$.*

Corollary 1. *Let the suppositions of Lemma 4 be fulfilled and $A'(t) \leq 0$ in $[a, \infty)$. Then the differential equation (a) for $\lambda = \bar{\lambda} > 0$ is oscillatory in $[a, \infty)$.*

Lemma 5. *Let $A(t) \geq 0$ in $[a, \infty)$ and let the differential equation (4) be disconjugated in $[a, \infty)$. Let u_1, u_2 be independent solutions of (4) and let $u_1(t_0) = 1, u_1'(t_0) = 0, u_2(t_0) = 1, u_2'(t_0) = 0, a < t_0 < \infty$. Then there is $u_2(t) > 0$ for $t > t_0$. And $u_1(t)$ has at most one zero to the right of t_0 .*

Remark 2 [6, Lemma 2.23]. *Let the suppositions of Lemma 5 be fulfilled. If u is a solution of (4) and $u(t) \neq 0$ for $t \geq t_1$, then*

$$0 < (t + d)v(t) \leq 1, \quad t \geq t_1$$

where $v(t) = \frac{u'(t)}{u(t)}$, $d = -t_1 + 1/v(t_1)$.

Lemma 6. *Let the suppositions of Lemma 5 be fulfilled and let $b(t) > 0$ for $t \in [a, \infty)$ and $\lambda > 0$. Let further y be a solution of (a) and let for $\lambda = \bar{\lambda}$ be $y(t_0, \bar{\lambda}) = 0, y'(t_0, \bar{\lambda}) \neq 0, y''(t_0, \bar{\lambda}) \neq 0$ for $a \leq t_0 < \infty$. Let, moreover $y(t, \bar{\lambda}) \neq 0$ for $t > t_0$. Then*

$$(8) \quad y(t, \bar{\lambda}) = u_2(t) \left[\frac{y''(t_0, \bar{\lambda})}{2} u_2(t) + y'(t_0, \bar{\lambda}) u_1(t) \right] - \frac{1}{2} \bar{\lambda} \int_{t_0}^t b(\tau) \begin{vmatrix} u_1(t) & u_2(t) \\ u_1(\tau) & u_2(\tau) \end{vmatrix}^2 y(\tau, \bar{\lambda}) d\tau,$$

where u_1, u_2 form a fundamental set of solutions of (4) with the properties given in Lemma 5.

The proof of Lemma 6 is given in [4] at the beginning of Section 3, Chap. I, §3.

Corollary 2. *If $y(t, \bar{\lambda}) > 0$ [$y(t, \bar{\lambda}) < 0$] for $t > t_0$ in (8) then $y'(t_0, \bar{\lambda}) > 0$ [$y'(t_0, \bar{\lambda}) < 0$], $u_2(t) > 0$ and $u(t) = y'(t_0, \bar{\lambda}) u_1(t) + \frac{y''(t_0, \bar{\lambda})}{2} u_2(t) > 0$ [$u(t) < 0$] for $t > t_0$.*

Corollary 3. *Let the suppositions of Lemma 6 be fulfilled then there exist constants $k_1 = y'(t_0, \bar{\lambda})$, $k_2 = \frac{y''(t_0, \bar{\lambda})}{2}$ such that $|y(t, \bar{\lambda})| \leq u_2(t) |k_1 u_1(t) + k_2 u_2(t)|$ for $t > t_0$, or $y(t, \bar{\lambda}) = o(tu_2(t) [k_1 u_1(t) + k_2 u_2(t)])$ for $t \rightarrow \infty$.*

Adaptation of Oscillation theorem [4, Theorem 4.5] to (a) in our case yields the following lemma.

Lemma 7. *Suppose that $A(t) \geq 0$, $A'(t) \leq 0$ and $b(t) \geq k > 0$ for $t \in [a, \infty)$. Let $\lambda \in (0, \infty)$ and let $y(t, \lambda)$ be a nontrivial solution of (a) with $y(a, \lambda) = 0$. Then for any fixed $b > a$, the number of zeros of y on $[a, b]$ increases to infinity as $\lambda \rightarrow \infty$ and the distance between any consecutive zeros of y converges to zero.*

The continuous dependence of zeros of solutions of (a) upon the parameter λ is given in following lemma.

Lemma 8. [4, Lemma 4.2] *Let $A'(t)$, $b(t) > 0$ be continuous functions in $[a, \infty)$. Let y be a nontrivial solution of (a) on $[a, \infty)$ such that $y(\alpha, \lambda) = 0$, $a \leq \alpha < \infty$, for all $\lambda \in (0, \infty)$. Then the zeros of y on (α, ∞) (if they exist) are continuous functions of the parameter $\lambda \in (0, \infty)$.*

With the help of results given in the preceding lemmas, remarks and corollaries one can prove the following theorem regarding the singular eigenvalue problem (a), (1), (3) or (a), (2), (3).

Theorem 1. *Let $A(t) \geq 0$, $A'(t) \leq 0$, $b(t) > 0$ be continuous functions in $[a, \infty)$ and let $A'(t) + b(t) > 0$ for $t \in [a, \infty)$. Let $\int_a^\infty t^2[b(t) - A'(t)]dt < \infty$ and let the second order differential equation (5) be disconjugate in $[a, \infty)$. Let further $a \leq b < c$ be fixed arbitrarily. Then there exists a natural number ν and a sequence of parameters $\lambda \left\{ \lambda_{\nu+p} \right\}_{p=0}^\infty$ (eigenvalues) such that $\lambda_{\nu+p} < \lambda_{\nu+p+1}$, $p = 0, 1, 2, \dots$ and $\lim_{p \rightarrow \infty} \lambda_{\nu+p} = \infty$ and a corresponding sequence of functions $\left\{ y_{\nu+p} \right\}_{p=0}^\infty$ (eigenfunctions) such that $y_{\nu+p} = y(t, \lambda_{\nu+p})$ is a solution of (a) for $\lambda = \lambda_{\nu+p}$, has a finite number of zeros on (a, ∞) with the last zero at $t_0^{\nu+p}$. This solution $y_{\nu+p}$ fulfills the boundary conditions (1), (3) or (2), (3) and has exactly $\nu + p$ zeros in (b, c) .*

Proof. We prove the case $a < b < c$. In the case $a = b$, i.e. (1), (3) the proof is similar.

Let $a < b < c < \infty$. Let $y = y(t, \lambda), \lambda > 0$, be a nontrivial solution of (a) such that $y(a, \lambda) = y(b, \lambda) = 0$ for all $\lambda > 0$. By Lemma 2 solution y is nonoscillatory for each $\lambda = \bar{\lambda} > 0$. Now, construct the differential equation

$$(A) \quad Y''' + 2A(t)Y' + [A'(t) + \lambda B(t)]Y = 0 \quad \text{on } [a, \infty)$$

where

$$B(t) = \begin{cases} b(t) & \text{for } t \in [b, c] \\ b(c) & \text{for } t \geq c. \end{cases}$$

Let $Y = Y(t, \lambda)$ be a solution of (A) on $[a, \infty)$ such that $Y(a, \lambda) = Y(b, \lambda) = 0$ and $Y(t, \lambda) = y(t, \lambda)$ for $t \in [a, c]$ and $\lambda \in (0, \infty)$.

By Lemma 4 and Corollary 1 the differential equation (A) is oscillatory on $[a, \infty)$ for each $\bar{\lambda} \in (0, \infty)$ and therefore the solution Y is oscillatory on $[a, \infty)$ for each $\bar{\lambda} > 0$.

Let $\lambda = \bar{\lambda}$ be fixed. Let $Y(t, \bar{\lambda})$ have exactly ν zeros in (b, c) . Let $t_\nu(\lambda)$ be ν -th zero of $Y(t, \lambda)$. Then there is $t_\nu(\bar{\lambda}) < c \leq t_{\nu+1}(\bar{\lambda})$. By Lemma 7 there exist λ^* such that $t_{\nu+1}(\lambda^*) < c$ and by Lemma 8 (continuous dependence of zeros) there exists λ_ν , $\bar{\lambda} \leq \lambda_\nu < \lambda^*$ such that $t_{\nu+1}(\lambda_\nu) = c$ and $y(t, \lambda_\nu)$ has exactly ν zeros in (b, c) . But, we know that $Y(t, \lambda_\nu) = y(t, \lambda_\nu)$ on $[a, c]$. By Lemma 2 applied to λ_ν there exists $t'_0 \geq c$ such that $y(t, \lambda_\nu)$ has finite number of zeros to the right of c and t'_0 is its last zero on $[c, \infty)$. Then by Corollary 3, when $t_0 = t'_0$, the inequality (3) holds.

Continuing in the same manner we can find a sequence of values

$$\lambda_\nu, \lambda_{\nu+1}, \dots, \lambda_{\nu+p}, \dots$$

and the corresponding sequence of functions $\left\{ y_{\nu+p} \right\}_{p=0}^{\infty}$ (eigenfunctions) with the prescribed properties and the theorem is proved. \square

REFERENCES

- [1] Elbert, A., Takaši, K. and Naito, M., *Singular eigenvalue problems for second order linear ordinary differential equations*, Arch. Math. (Brno), T. **34** (1998), 59–72.
- [2] Gera, M., *Bedingungen der Nicht-oszillationsfähigkeit für die lineare Differentialgleichung dritter Ordnung*, Mat. Čas. Slov. akad. vied, **21** (1971), 65–80.
- [3] Greguš, M., *On certain third order eigenvalue problem*, CDDE Brno 2000, Masaryk University, Arch. Math. (Brno), T. **36** (2000), 461–464.
- [4] Greguš, M., *Third Order Differential Equations*, D. Riedel Publishing Company, Dordrecht, Boston, Lancaster, Tokyo (1987).
- [5] Greguš, M., Jr., *On certain third order boundary value problems on infinite interval*, CDDE Brno 2000, Masaryk University, Arch. Math. (Brno), T. **36** (2000), 465–468.
- [6] Swanson, C. A., *Comparison and oscillation theory of linear differential equations*, Academic Press, New York and London 1968.

DEPARTMENT OF INFORMATION SYSTEMS
 FACULTY OF MANAGEMENT, COMENIUS UNIVERSITY
 ODBOJAROV 10, P.O. BOX 95
 820 05 BRATISLAVA 25, SLOVAKIA
 E-mail: michal.gregus@fm.uniba.sk