

Johan Fastenakels

Ideal tubular hypersurfaces in real space forms

Archivum Mathematicum, Vol. 42 (2006), No. 3, 295--305

Persistent URL: <http://dml.cz/dmlcz/108009>

Terms of use:

© Masaryk University, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

IDEAL TUBULAR HYPERSURFACES IN REAL SPACE FORMS

JOHAN FASTENAKELS

ABSTRACT. In this article we give a classification of tubular hypersurfaces in real space forms which are $\delta(2, 2, \dots, 2)$ -ideal.

1. IDEAL IMMERSIONS

Let M be a Riemannian n -manifold. Denote by $K(\pi)$ the sectional curvature of M associated with a plane section $\pi \subset T_p M$, $p \in M$. For any orthonormal basis e_1, \dots, e_n of the tangent space $T_p M$, the scalar curvature τ at p is defined to be

$$(1) \quad \tau(p) = \sum_{i < j} K(e_i \wedge e_j).$$

When L is a 1-dimensional subspace of $T_p M$, we put $\tau(L) = 0$. If L is a subspace of $T_p M$ of dimension $r \geq 2$, we define the scalar curvature $\tau(L)$ of L by

$$(2) \quad \tau(L) = \sum_{\alpha < \beta} K(e_\alpha \wedge e_\beta), \quad 1 \leq \alpha, \beta \leq r,$$

where $\{e_1, \dots, e_r\}$ is an orthonormal basis of L .

For an integer $k \geq 0$, denote by $\mathcal{S}(n, k)$ the finite set consisting of unordered k -tuples (n_1, \dots, n_k) of integers ≥ 2 satisfying $n_1 < n$ and $n_1 + \dots + n_k \leq n$. Let $\mathcal{S}(n)$ be the union $\cup_{k \geq 0} \mathcal{S}(n, k)$. If $n = 2$, we have $k = 0$ and $\mathcal{S}(2) = \{\emptyset\}$.

For each $(n_1, \dots, n_k) \in \mathcal{S}(n)$, the invariant $\delta(n_1, \dots, n_k)$ is defined in [3] by:

$$(3) \quad \delta(n_1, \dots, n_k)(p) = \tau(p) - S(n_1, \dots, n_k)(p),$$

where

$$S(n_1, \dots, n_k)(p) = \inf \{ \tau(L_1) + \dots + \tau(L_k) \}$$

and L_1, \dots, L_k run over all k mutually orthogonal subspaces of $T_p M$ such that $\dim L_j = n_j$, $j = 1, \dots, k$. Clearly, the invariant $\delta(\emptyset)$ is nothing but the scalar curvature τ of M .

2000 *Mathematics Subject Classification*: 53B25.

Key words and phrases: tubular hypersurfaces, ideal immersion, real space form.

The author is Research assistant of the Research Foundation - Flanders (FWO).

Received April 28, 2006.

For a given partition $(n_1, \dots, n_k) \in \mathcal{S}(n)$, we put

$$(4) \quad b(n_1, \dots, n_k) = \frac{1}{2} \left(n(n-1) - \sum_{j=1}^k n_j(n_j-1) \right),$$

$$(5) \quad c(n_1, \dots, n_k) = \frac{n^2(n+k-1 - \sum n_j)}{2(n+k - \sum n_j)}.$$

For each real number c and each $(n_1, \dots, n_k) \in \mathcal{S}(n)$, the associated normalized invariant $\Delta_c(n_1, \dots, n_k)$ is defined by

$$(6) \quad \Delta_c(n_1, \dots, n_k) = \frac{\delta(n_1, \dots, n_k) - b(n_1, \dots, n_k)c}{c(n_1, \dots, n_k)}.$$

We recall the following general result from [3].

Theorem 1. *Let M be an n -dimensional submanifold of a real space form $R^m(c)$ of constant sectional curvature c . Then for each $(n_1, \dots, n_k) \in \mathcal{S}(n)$ we have*

$$(7) \quad H^2 \geq \Delta_c(n_1, \dots, n_k),$$

where H^2 is the squared norm of the mean curvature vector.

The equality case of inequality (7) holds at a point $p \in M$ if and only if, with respect to a suitable orthonormal basis $e_1, \dots, e_n, e_{n+1}, \dots, e_m$ at p , the shape operators $A_r = A_{e_r}$, $r = n+1, \dots, m$ of M in $R^m(c)$ at p take the following forms:

$$(8) \quad A_{n+1} = \begin{pmatrix} a_1 & 0 & 0 & \cdots & 0 \\ 0 & a_2 & 0 & \cdots & 0 \\ 0 & 0 & a_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_n \end{pmatrix},$$

$$(9) \quad A_r = \begin{pmatrix} A_1^r & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & A_k^r & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad r = n+2, \dots, m,$$

where a_1, \dots, a_n satisfy

$$(10) \quad a_1 + \cdots + a_{n_1} = \cdots = a_{n_1+\dots+n_{k-1}+1} + \cdots + a_{n_1+\dots+n_k} \\ = a_{n_1+\dots+n_k+1} = \cdots = a_n$$

and each A_j^r is an $n_j \times n_j$ submatrix such that

$$(11) \quad \text{trace}(A_j^r) = 0, \quad (A_j^r)^t = A_j^r, \quad r = n+2, \dots, m; \quad j = 1, \dots, k.$$

For an isometric immersion $x : M \rightarrow R^m(c)$ of a Riemannian n -manifold into $R^m(c)$, this theorem implies that

$$(12) \quad H^2(p) \geq \hat{\Delta}_c(p),$$

where $\hat{\Delta}_c$ denotes the invariant on M defined by

$$(13) \quad \hat{\Delta}_c = \max \{ \Delta_c(n_1, \dots, n_k) \mid (n_1, \dots, n_k) \in \mathcal{S}(n) \}.$$

In general, there do not exist direct relations between these new invariants.

Applying inequality (12) B. Y. Chen introduced in [4] the notion of ideal immersions as follows.

Definition 1. *An isometric immersion $x : M \rightarrow R^m(c)$ is called an ideal immersion if the equality case of (12) holds at every point $p \in M$. An isometric immersion is called (n_1, \dots, n_k) -ideal if it satisfies $H^2 = \Delta_c(n_1, \dots, n_k)$ identically for $(n_1, \dots, n_k) \in \mathcal{S}(n)$.*

Physical Interpretation of Ideal Immersions. An isometric immersion $x : M \rightarrow R^m(c)$ is ideal means that M receives the least possible amount of tension (given by $\hat{\Delta}_c(p)$) at each point $p \in M$ from the ambient space. This is due to (12) and the well-known fact that the mean curvature vector field is exactly the tension field for isometric immersions. Therefore, the squared mean curvature $H^2(p)$ at a point $p \in M$ simply measures the amount of tension M is receiving from the ambient space $R^m(c)$ at that point.

2. TUBULAR HYPERSURFACES

Recall the definition of the exponential mapping \exp of a Riemannian manifold M . Denote by $\gamma_v, v \in T_pM$, the geodesic of M through p such that $\gamma'(p) = v$. Then we have that

$$\exp : TM \rightarrow M : (p, v) \mapsto \exp_p(v) = \gamma_v(1)$$

for every $v \in T_pM$ for which γ_v is defined on $[0, 1]$.

Let B^ℓ be a topologically imbedded ℓ -dimensional ($\ell < n$) submanifold in an $n + 1$ -dimensional real space form $R^{n+1}(c)$. Denote by $\nu_1(B^\ell)$ the unit normal subbundle of the normal bundle $T^\perp(B^\ell)$ of B^ℓ in $R^{n+1}(c)$. Then, for a sufficiently small $r > 0$, the mapping

$$\psi : \nu_1(B^\ell) \rightarrow R^{n+1}(c) : (p, e) \mapsto \exp_p(re)$$

is an immersion which is called the *tubular hypersurface* with radius r about B^ℓ . We denote it by $T_r(B^\ell)$.

In this article, we consider $r > 0$ such that the map is an immersion only. Thus, the shape operator of the tubular hypersurface $T_r(B^\ell)$ is a well defined self-adjoint linear operator at each point.

Now take an arbitrary point p in B^ℓ and a vector u in $\nu_1(B^\ell)$. Denote with $\kappa_1(u), \dots, \kappa_\ell(u)$ the eigenvalues of the shape operator of B^ℓ in $R^{n+1}(c)$ with respect to u at the point p . Then we can give an expression for the principal curvatures $\bar{\kappa}_1, \dots, \bar{\kappa}_m$ of the tubular hypersurface in the point $\exp(p, u)$. We consider three cases.

(i) $c = 0$. In the Euclidean case, we find

$$(14) \quad \bar{\kappa}_i = \frac{\kappa_i(u)}{1 - r\kappa_i(u)}, \quad i = 1, \dots, \ell,$$

$$(15) \quad \bar{\kappa}_\alpha(r) = -\frac{1}{r}, \quad \alpha = \ell + 1, \dots, n.$$

(ii) $c = 1$. For the unit sphere, we can simplify the expressions by denoting $\kappa_1(u) = \tan(\theta_1), \dots, \kappa_\ell(u) = \tan(\theta_\ell)$ with $-\frac{\pi}{2} < \theta_i < \frac{\pi}{2}$. Then we have

$$(16) \quad \bar{\kappa}_i = \tan(\theta_i + r), \quad i = 1, \dots, \ell,$$

$$(17) \quad \bar{\kappa}_\alpha(r) = -\cot(r), \quad \alpha = \ell + 1, \dots, n.$$

(iii) $c = -1$. In the hyperbolic space we have

$$(18) \quad \bar{\kappa}_i = \frac{\kappa_i(u) \coth(r) - 1}{\coth(r) - \kappa_i(u)}, \quad i = 1, \dots, \ell,$$

$$(19) \quad \bar{\kappa}_\alpha(r) = -\coth(r), \quad \alpha = \ell + 1, \dots, n.$$

More details can be found in [2].

3. $\delta_{(2,2,\dots,2)}$ -IDEAL TUBULAR HYPERSURFACES

In this section we will give a complete classification of tubular hypersurfaces in real space forms for which the immersion defined in the previous section is a $\delta_{(2,2,\dots,2)}$ -ideal immersion. We again consider three cases.

In the Euclidean space \mathbb{E}^{n+1} .

Theorem 2. *A tubular hypersurface $T_r(B^\ell)$ in \mathbb{E}^{n+1} ($n > 2$) satisfies equality in (7) for k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$ if and only if one of the following three cases occurs:*

- (1) $\ell = 0$ and the tubular hypersurface is a hypersphere.
- (2) $\ell = k \in \{1, \dots, [\frac{n}{2}]\}$ and the tubular hypersurface is an open part of a spherical hypercylinder: $\mathbb{E}^\ell \times S^{n-\ell}(r)$.
- (3) n is even, $\ell = k = \frac{n}{2}$ and B^ℓ is totally umbilical.

Proof. Let $\kappa_1(u), \dots, \kappa_\ell(u)$ be the eigenvalues of the shape operator of B^ℓ in \mathbb{E}^{n+1} with respect to a unit normal vector u at p . Then we find, according to the previous section, that the principal curvatures of the tubular hypersurface $T_r(B)$

at $p + ru$ are given by

$$(20) \quad \bar{\kappa}_i = \frac{\kappa_i(u)}{1 - r\kappa_i(u)}, \quad i = 1, \dots, \ell,$$

$$(21) \quad \bar{\kappa}_\alpha(r) = -\frac{1}{r}, \quad \alpha = \ell + 1, \dots, n.$$

Suppose now that $T_r(B)$ satisfies equality in (7) for a k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$.

If $\ell = 0$, the tubular hypersurface is an open part of an hypersphere. This gives us the first case in the theorem.

If $\ell = 1$, the multiplicity of $-\frac{1}{r}$ is $n - 1$. From (8) and (10) we find the following three cases:

- $\bar{\kappa}_1 + (-\frac{1}{r}) = -\frac{1}{r}$, which implies that $\bar{\kappa}_1 = 0$.
- $\frac{\kappa_1}{1-r\kappa_1} = -\frac{1}{r} - \frac{1}{r} = -\frac{2}{r}$, so we have that $r\kappa_1 = 2$. This gives a contradiction with the fact that $\kappa_1(-u) = -\kappa_1(u)$.
- $\frac{\kappa_1}{1-r\kappa_1} + (-\frac{1}{r}) = -\frac{2}{r}$, from which we also get a contradiction.

So we see that $\bar{\kappa}_1 = 0$ and that $k = 1$. Thus B^1 is an open part of a line segment and the tubular hypersurface is an open part of $\mathbb{E}^1 \times S^{n-1}(r)$. This gives a special case of case (2) of the theorem.

Suppose now that $\ell \geq 2$, then (8) and (10) imply that we have one of the following five cases:

- (a) for all unit normal vectors u of B^ℓ , we have

$$(22) \quad \kappa_1(u) = \dots = \kappa_\ell(u) = 0$$
 and $\ell = k \leq \frac{n}{2}$;
- (b) for all unit normal vectors u of B^ℓ , we have

$$(23) \quad \bar{\kappa}_1(u) = \dots = \bar{\kappa}_\ell(u) \neq 0,$$
 n is even and $k = \ell = \frac{n}{2}$;
- (c) for all $i \in \{1, \dots, \ell\}$ there exists a $j \in \{1, \dots, \ell\}$ such that $i \neq j$ and such that:

$$(24) \quad \frac{\kappa_i(u)}{1 - r\kappa_i(u)} + \frac{\kappa_j(u)}{1 - r\kappa_j(u)} = -\frac{1}{r};$$
- (d) for all $i \in \{1, \dots, \ell\}$ there exists a $j \in \{1, \dots, \ell\}$ such that $i \neq j$ and such that:

$$(25) \quad \frac{\kappa_i(u)}{1 - r\kappa_i(u)} - \frac{1}{r} = \frac{\kappa_j(u)}{1 - r\kappa_j(u)};$$
- (e) $\ell = k = 2, n = 4$ and

$$(26) \quad \frac{\kappa_1(u)}{1 - r\kappa_1(u)} + \frac{\kappa_2(u)}{1 - r\kappa_2(u)} = -\frac{2}{r}.$$

Case (a) implies that B^ℓ is totally geodesic. Thus the tubular hypersurface is an open part of a spherical hypercylinder $\mathbb{E}^\ell \times S^{n-\ell}(r)$, which gives case (2) of the theorem.

Case (b) gives us case (3) of the theorem because $\bar{\kappa}_i = \bar{\kappa}_j$ if and only if $\kappa_i = \kappa_j$. Next we want to proof that cases (c), (d) and (e) cannot occur.

From (24), we find that

$$(27) \quad 1 = r^2 \kappa_i(u) \kappa_j(u)$$

for every u . This is impossible since the codimension of B^ℓ in \mathbb{E}^{n+1} is at least 2. We can see this in the following way. Because the codimension is at least 2, we can take a plane in the normal space which contains u . If $\kappa_i(u) = 0$, then we have a contradiction at once. Otherwise $\kappa_i(u)$ is strict positive or strict negative. Then we have that $\kappa_i(-u)$ is strict negative or strict positive respectively. Now we rotate u in the chosen plane to $-u$. Because the principal curvature is a continuous function, there exists a normal vector ξ for which $\kappa_i(\xi) = 0$. Putting ξ in equation (27) gives a contradiction.

From (25) we find analogously that

$$(28) \quad 1 - 2r\kappa_i(u) - r^2\kappa_i(u)\kappa_j(u) = 0.$$

Because $\kappa_i(-u) = -\kappa_i(u)$ we have also that

$$(29) \quad 1 + 2r\kappa_i(u) - r^2\kappa_i(u)\kappa_j(u) = 0.$$

Combining (28) and (29) then gives

$$4r\kappa_i(u) = 0,$$

which gives a contradiction unless all the principal curvatures of B^ℓ are zero. But then we are again in case (a).

Similarly case (e) gives a contradiction since we find from (26) that $\kappa_1 + \kappa_2 = \frac{2}{r}$. The converse is trivial. □

In the sphere $S^{n+1}(1)$. First we recall the definition of an austere submanifold in the sense of Harvey and Lawson [5].

Definition 2. We call a submanifold M of a Riemannian manifold \widetilde{M} austere if for every normal $\xi \in T^\perp M$ the set of all eigenvalues of the shape operator counted with multiplicities is invariant under multiplication with -1 .

Theorem 3. A tubular hypersurface $T_r(B^\ell)$ in $S^{n+1}(1)$ ($n > 2$) satisfies equality in (7) for a k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$ if and only if one of the following four cases occur:

- (1) $\ell = 0$ and the tubular hypersurface is a geodesic sphere with radius $r \in]0, \pi[$.
- (2) $n > \ell \geq \frac{n}{2}$, $k = n - \ell$, $r = \frac{\pi}{2}$ and B^ℓ is a totally umbilical submanifold in $S^{n+1}(1)$.

- (3) $\ell = 2k < n$, $r = \frac{\pi}{2}$ and B^ℓ is an austere submanifold in $S^{n+1}(1)$.
- (4) n is even, $\ell = k = \frac{n}{2}$ and B^ℓ is totally umbilical.

Proof. Let B^ℓ be an ℓ -dimensional submanifold inbedded in $S^{n+1}(1)$. For every unit normal vector u of B^ℓ at a point p we denote by $\kappa_1(u), \dots, \kappa_\ell(u)$ the eigenvalues of the shape operator of B^ℓ in $S^{n+1}(1)$ with respect to u . Suppose now that

$$(30) \quad \kappa_i(u) = \tan(\theta_i), \quad -\frac{\pi}{2} < \theta_i < \frac{\pi}{2}, \quad 1 \leq i \leq \ell.$$

Then we know from the previous section that the principal curvatures of the tubular hypersurface $T_r(B^\ell)$ in $S^{n+1}(1)$ at $\cos(r)p + \sin(r)u$ are given by

$$(31) \quad \begin{aligned} \bar{\kappa}_i &= \tan(\theta_i + r), & i &= 1, \dots, \ell, \\ \bar{\kappa}_\alpha(r) &= -\cot(r), & \alpha &= \ell + 1, \dots, n. \end{aligned}$$

Suppose that $T_r(B^\ell)$ satisfies (7) for a k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$.

If $\ell = 0$, the tubular hypersurface is totally umbilical in $S^{n+1}(1)$. Then theorem 1 implies that $T_r(B^\ell)$ with radius $r \in]0, \pi[$ satisfies (7) for a k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$ if and only if $k = 0$ or $k = \frac{n}{2}$. So we find that $T_r(B^\ell)$ is a geodesic sphere. This gives us case (1).

If $\ell = 1$, then (8) and (10) imply that we are in one of the following cases:

- $\frac{\kappa_1 + \tan r}{1 - \kappa_1 \tan r} + (-\cot(r)) = -\cot(r)$, which implies that $\kappa_1(u) = -\tan(r)$ for every unit normal vector u of B^1 in $S^{n+1}(1)$. This gives a contradiction with the fact that $\kappa_1(-u) = -\kappa_1(u)$.
- $\frac{\kappa_1 + \tan r}{1 - \kappa_1 \tan r} = -2 \cot r$, so we find $\kappa_1 \tan r = 2 + \tan^2 r$. Because $\kappa_1(-u) = -\kappa_1(u)$ we have $2 + \tan^2 r = 0$ which also gives a contradiction.
- $\frac{\kappa_1 + \tan r}{1 - \kappa_1 \tan r} + (-\cot r) = -2 \cot r$, which becomes $\tan^2 r = -1$. This clearly also gives a contradiction.

In each case we get a contradiction, so $\ell = 1$ cannot occur.

Suppose now that $\ell \geq 2$, then theorem 1 implies that we are in one of the following cases:

- (a) for all unit normal vectors u of B^ℓ we have that

$$(32) \quad \tan(\theta_j + r) = 0, \quad j = 1, \dots, \ell$$

and $\ell = k \leq \frac{n}{2}$;

- (b) for any unit normal vector u of B^ℓ we have that

$$(33) \quad \tan(\theta_1 + r) = \dots = \tan(\theta_\ell + r) \neq 0,$$

n is even and $k = \ell = \frac{n}{2}$;

- (c) for all $i \in \{1, \dots, \ell\}$ there exists a $j \in \{1, \dots, \ell\}$ such that $i \neq j$ and such that:

$$(34) \quad \tan(\theta_i + r) - \cot(r) = \tan(\theta_j + r);$$

(d) for all $i \in \{1, \dots, \ell\}$ there exists a $j \in \{1, \dots, \ell\}$ such that $i \neq j$ and such that:

$$(35) \quad \tan(\theta_i + r) + \tan(\theta_j + r) = -\cot(r);$$

(e) $\ell = k = 2, n = 4$ and

$$(36) \quad \tan(\theta_1 + r) + \tan(\theta_2 + r) = -2\cot(r).$$

Suppose now that we are in case (a) and thus (32) holds. Then we see that $\kappa_j(u)\cot(r) + 1 = 0$ for any unit normal vector u of B^ℓ in $S^{n+1}(1)$. This is impossible since $\kappa_j(-u) = -\kappa_j(u)$.

If case (b) holds, then we get case (4) of the theorem, since

$$\frac{\kappa_i + \tan r}{1 - \kappa_i \tan r} = \frac{\kappa_j + \tan r}{1 - \kappa_j \tan r}$$

implies that

$$(\kappa_i - \kappa_j)(1 + \tan^2 r) = 0.$$

Suppose now that we are in case (c). Then we have from (34) that:

$$(37) \quad \cot^3(r) - 2\kappa_i \cot^2(r) + \kappa_i \kappa_j \cot(r) + (\kappa_j - \kappa_i) = 0.$$

We use again the fact that $\kappa_i(-u) = -\kappa_i(u)$ and therefore we find

$$(38) \quad \cot(r)(\cot^2(r) + \kappa_i(u)\kappa_j(u)) = 0$$

and

$$(39) \quad 2\kappa_i(u)\cot^2(r) + \kappa_i(u) - \kappa_j(u) = 0.$$

If $\cot(r) \neq 0$, then (38) implies that $\cot^2(r) = -\kappa_i(u)\kappa_j(u)$. Because $\ell < n$ we get a contradiction with the same argument as in the preceding proof.

Thus we have $\cot(r) = 0$, and thus $r = \frac{\pi}{2}$. From (39) we also see that $\kappa_i(u) = \kappa_j(u)$. Without loss of generality, we may assume

$$a_1 = \mu, a_2 = 0, a_3 = \mu, a_4 = 0, \dots, a_{2k-1} = \mu, a_{2k} = 0, a_{2k+1} = \mu, \dots, a_n = \mu$$

where $\mu = -\frac{1}{\kappa_1}$ and a_1, \dots, a_n are given by theorem (1).

Furthermore we see that $\tan(\theta_i + r) \neq 0$ since $-\frac{\pi}{2} < \theta_i < \frac{\pi}{2}$ and from (31) we find that $\cot(r)$ has multiplicity $n - \ell$. So theorem (1) implies that $\ell \geq \frac{n}{2}$ and $\tan(\theta_1 + r) = \dots = \tan(\theta_\ell + r)$. This implies also that $\tan(\theta_1) = \dots = \tan(\theta_\ell)$ and thus that B^ℓ is totally umbilical. Moreover we see that theorem (1) implies that $k = n - \ell$. This gives rise to case (2).

Suppose now that we are in case (d) and thus that (35) holds. Then we have

$$(40) \quad \cot^3(r) + 2\cot(r) - \kappa_i \kappa_j \cot(r) - (\kappa_i + \kappa_j) = 0.$$

If we use that $\kappa_i(-u) = -\kappa_i(u)$ we find

$$(41) \quad \cot(r)(\cot^2(r) + 2 - \kappa_i \kappa_j) = 0$$

and

$$(42) \quad \kappa_i + \kappa_j = 0.$$

Like in case (c) we get a contradiction if $\cot(r) \neq 0$. So we find $\cot(r) = 0$ and thus $r = \frac{\pi}{2}$. Moreover we have $\kappa_i = -\kappa_j$. Without loss of generality, we may assume

$$a_1 = \tan(\theta_1 + r) = -\frac{1}{\kappa_1}, \quad a_2 = \tan(\theta_2 + r) = -\frac{1}{\kappa_2}, \dots, \quad a_n = -\cot(r) = 0$$

We also know that $\tan(\theta_j + r) \neq 0$ (since $-\frac{\pi}{2} < \theta_j < \frac{\pi}{2}$). Thus (31) and theorem 1 imply that B^ℓ is an austere submanifold in $S^{n+1}(1)$; in particular ℓ is even. This gives case (3).

A similar computation as in case (d) shows that case (e) gives a contradiction. The converse can be verified easily. □

In the hyperbolic space $H^{n+1}(-1)$.

Theorem 4. *A tubular hypersurface $T_r(B^\ell)$ in $H^{n+1}(-1)$ ($n > 2$) satisfies equality in (7) for a k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$ if and only if we are in one of the following three cases:*

- (1) $\ell = 0$ and the tubular hypersurface is a geodesic sphere with radius $r > 0$.
- (2) $\ell = 2k$, B^ℓ is totally geodesic and $r = \coth^{-1}(\sqrt{2})$.
- (3) n is even, $\ell = k = \frac{n}{2}$ and B^ℓ is totally umbilical.

Proof. Let B^ℓ be an ℓ -dimensional submanifold in the hyperbolic space $H^{n+1}(-1)$ and $T_r(B^\ell)$ be the tubular hypersurface of B^ℓ in $H^{n+1}(-1)$. Suppose that $T_r(B^\ell)$ satisfies (7) for a k -tuple $(n_1, \dots, n_k) = (2, \dots, 2)$. For any unit normal vector u of B^ℓ at a point p of B^ℓ denote with $\kappa_1(u), \dots, \kappa_\ell(u)$ the principal curvatures of B^ℓ in $H^{n+1}(-1)$ at p with respect to u . Then it follows from section 2 that the principal curvatures $\bar{\kappa}_1, \dots, \bar{\kappa}_n$ of the shape operator of $T_r(B^\ell)$ are given by:

$$(43) \quad \bar{\kappa}_i = \frac{\kappa_i(u) \coth(r) - 1}{\coth(r) - \kappa_i(u)}, \quad i = 1, \dots, \ell,$$

$$(44) \quad \bar{\kappa}_\alpha(r) = -\coth(r), \quad \alpha = \ell + 1, \dots, n.$$

If $\ell = 0$, then the tubular hypersurface is totally umbilical. So we find from theorem (1) that $k = 0$ or $k = \frac{n}{2}$ and $T_r(B^\ell)$ is a geodesic sphere. Thus we are in case (1).

If $\ell = 1$, then from theorem 1 and (43) it follows that we are in one of the following cases:

- $\bar{\kappa}_1 - \cot r = -\cot r$, which implies immediately that $\bar{\kappa}_1(u) = 0$ for any unit normal vector u of B^1 in $S^{n+1}(1)$. Then (43) would imply that $\kappa_1(u) = -\tanh(r)$ which gives a contradiction with the fact that $\kappa_1(-u) = -\kappa_1(u)$ since $r \in \mathbb{R}_0^+$.
- $\frac{\kappa_1 \coth r - 1}{\coth r - \kappa_1} = -2 \coth r$, so we find $\kappa_1 \coth r = 2 \coth^2 r - 1$. Because $\kappa_1(-u) = -\kappa_1(u)$ this implies that $\coth^2 r = \frac{1}{2}$ which gives a contradiction since $\coth^2 r$ is always greater than 1.
- $\frac{\kappa_1 \coth r - 1}{\coth r - \kappa_1} + (-\cot r) = -2 \cot r$, this implies $\coth^2 r = 1$ which gives a contradiction as above.

Thus we see that the case $\ell = 1$ cannot occur.

Suppose now that $\ell \geq 2$, then theorem (1) implies that one of the following cases occur:

(a) for all unit normal vectors u of B^ℓ we have

$$(45) \quad \kappa_i(u) \coth(r) = 1, \quad \text{for all } i \in \{1, \dots, \ell\}$$

and $\ell = k \leq \frac{n}{2}$;

(b) for alle unit normal vectors u of B^ℓ we have

$$(46) \quad \bar{\kappa}_1(u) = \dots = \bar{\kappa}_\ell(u) \neq 0,$$

n is even and $k = \ell = \frac{n}{2}$;

(c) for all $i \in \{1, \dots, \ell\}$ there exists a $j \in \{1, \dots, \ell\}$ such that $i \neq j$ and such that:

$$(47) \quad \frac{\kappa_i(u) \coth(r) - 1}{\coth(r) - \kappa_i(u)} - \coth(r) = \frac{\kappa_j(u) \coth(r) - 1}{\coth(r) - \kappa_j(u)};$$

(d) for all $i \in \{1, \dots, \ell\}$ there exists a $j \in \{1, \dots, \ell\}$ such that $i \neq j$ and such that:

$$(48) \quad \frac{\kappa_i(u) \coth(r) - 1}{\coth(r) - \kappa_i(u)} + \frac{\kappa_j(u) \coth(r) - 1}{\coth(r) - \kappa_j(u)} = -\coth(r);$$

(e) $\ell = k = 2$, $n = 4$ and

$$(49) \quad \frac{\kappa_1(u) \coth(r) - 1}{\coth(r) - \kappa_1(u)} + \frac{\kappa_2(u) \coth(r) - 1}{\coth(r) - \kappa_2(u)} = -2\coth(r).$$

We see at once that (45) and thus case (a) cannot occur since $\kappa_i(-u) = -\kappa_i(u)$. Suppose now that we are in case (b). The condition $\bar{\kappa}_i = \bar{\kappa}_j$ gives us

$$(\kappa_i - \kappa_j)(\coth^2 r - 1) = 0.$$

Because $\coth^2 r > 1$ this implies $\bar{\kappa}_i = \bar{\kappa}_j$ if and only if $\kappa_i = \kappa_j$. This is case (3) of the theorem.

Suppose that we are in case (c). Then from (47), we find

$$(50) \quad \coth^3(r) - 2\kappa_i \coth^2(r) + \kappa_i \kappa_j \coth(r) + \kappa_i - \kappa_j = 0.$$

Because $\kappa_i(-u) = -\kappa_i(u)$ we have

$$(51) \quad \coth^3(r) + \kappa_i \kappa_j \coth(r) = 0,$$

$$(52) \quad -2\kappa_i \coth^2(r) + \kappa_i - \kappa_j = 0.$$

From (51), it follows that $\kappa_i(u)\kappa_j(u) = -\coth^2(r)$ since $\coth(r) \neq 0$. But this gives a contradiction with the same argument as in the Euclidean case because the codimension is at least 2.

Analogously from (48) we find:

$$(53) \quad (\kappa_i + \kappa_j) \tanh^3(r) - (2 + \kappa_i \kappa_j) \tanh^2(r) + 1 = 0.$$

By switching to $-u$ we get:

$$(54) \quad -(\kappa_i + \kappa_j) \tanh^3(r) - (2 + \kappa_i \kappa_j) \tanh^2(r) + 1 = 0.$$

This implies that $\kappa_i(u) + \kappa_j(u) = 0$. Substituting this in (53) gives $\kappa_i(u)^2 = 2 - \coth^2(r)$. We can also substitute the other way round, then we find $\kappa_j(u)^2 = 2 - \coth^2(r)$. Thus κ_i must be zero for every $i \in \{1, \dots, \ell\}$. We see that B^ℓ is totally geodesic. We see also that in this case $r = \coth^{-1}(\sqrt{2})$. Thus we get as principal curvatures for $T_r(B^\ell)$ $\bar{\kappa}_i = -\frac{1}{\sqrt{2}} = -\frac{\sqrt{2}}{2}$, $i = 1, \dots, \ell$ and $\bar{\kappa}_\alpha = -\sqrt{2}$, $\alpha = \ell + 1, \dots, n$. From theorem (1) it follows that $\ell = 2k$. So we get case (2).

Case (e) cannot occur since similar computations as in case (d) give a contradiction.

The converse can be verified easily. □

REFERENCES

- [1] B. Y. Chen, *Some pinching and classification theorems for minimal submanifolds*, Arch. Math. **60** (1993), 568-578.
- [2] B. Y. Chen, *Tubular hypersurfaces satisfying a basic equality.*, Soochow Journal of Mathematics **20** No. 4 (1994), 569-586.
- [3] B. Y. Chen, *Some new obstructions to minimal and Lagrangian isometric immersions*, Japan J. Math. **26** (2000), 105-127.
- [4] B. Y. Chen, *Strings of Riemannian invariants, inequalities, ideal immersions and their applications*, in Third Pacific Rim Geom. Conf., (Intern. Press, Cambridge, MA), (1998), 7-60.
- [5] R. Harvey and H. B. Lawson, Jr., *Calibrated geometries*, Acta Math. **148** (1982), 47-157.

KATHOLIEKE UNIVERSITEIT LEUVEN, DEPARTEMENT WISKUNDE
 CELESTIJNENLAAN 200 B, B-3001 LEUVEN, BELGIUM
E-mail: johan.fastenakels@wis.kuleuven.be