

Bernhard Banaschewski
Bourbaki's Fixpoint Lemma reconsidered

Commentationes Mathematicae Universitatis Carolinae, Vol. 33 (1992), No. 2, 303--309

Persistent URL: <http://dml.cz/dmlcz/118498>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1992

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Bourbaki's Fixpoint Lemma reconsidered

B. BANASCHEWSKI

Abstract. A constructively valid counterpart to Bourbaki's Fixpoint Lemma for chain-complete partially ordered sets is presented to obtain a condition for one closure system in a complete lattice L to be stable under another closure operator of L . This is then used to deal with coproducts and other aspects of frames.

Keywords: complete lattice, closure operator, fixpoint, frame coproduct, compact frame

Classification: 06A15, 54A99, 54B99

A *preclosure operator* on a complete lattice L is a map $k_0 : L \rightarrow L$ which preserves the partial order and is upward, that is, $x \leq k_0(x)$ for all $x \in L$. For such k_0 , $\text{Fix}(k_0) = \{x \in L \mid k_0(x) = x\}$ is readily seen to be a closure system in L , that is, closed under arbitrary meets in L , and we let k be the associated closure operator. In various contexts, one would like to be able to conclude, for certain subsets $S \subseteq L$, the following

Stability Lemma. *S is k -stable whenever it is k_0 -stable.*

Now, one way of describing k is as the stable transfinite iterate of k_0 : if one defines, for any $x \in L$, any ordinal α and any limit ordinal λ ,

$$k_0^0(x) = x, \quad k_0^{\alpha+1}(x) = k_0(k_0^\alpha(x)), \quad k_0^\lambda(x) = \bigvee \{k_0^\alpha(x) \mid \alpha < \lambda\},$$

then $k = k_0^\gamma$ for the first γ such that $k_0^{\gamma+1} = k_0^\gamma$. Here, one sees by induction that any $\{k_0^\alpha(x) \mid \alpha < \beta\}$ is a chain, and hence the desired result follows for any $S \subseteq L$ closed under taking joins, in L , of (non-void) chains.

The same conclusion can also be obtained, without the use of ordinals, as an application of

Bourbaki's Fixpoint Lemma. *Any upward map of a chain-complete partially ordered set into itself has a fixpoint.*

For any S as above and $a \in S$,

$$P = \{x \in S \mid a \leq x \leq k(a)\}$$

It is a pleasure to acknowledge the hospitality of the Department of Mathematics & Computer Science, University of Lethbridge, during the summer of 1991 while this note was written. Also, the support of Natural Sciences and Engineering Research Council of Canada in the form of an operating grant is gratefully recorded

is chain-complete and is mapped into itself by k_0 . For the resulting fixpoint $c = k_0(c)$ in P , $a \leq c \leq k(a)$ implies $c = k(a)$ and thus $k(a) \in S$.

It is an open problem precisely what rules of logic are needed to establish this lemma, and specifically, whether it is constructively valid. The known proofs (for instance, Witt [6]) use, for the partially ordered set in question, that $x \leq y$ implies $x < y$ or $x = y$ for all elements x and y , and that, for certain subsets U, V, W , if $U \subseteq V \cup W$ then $U \subseteq V$ or there exist $x \in U$ such that $x \in W$. These steps are not constructively valid but they do hold in any Boolean topos (Johnstone [2]), and hence so does the Stability Lemma, for any S closed under taking joins of chains.

The purpose of this note is to establish a constructively valid counterpart of Bourbaki's Fixpoint Lemma, to derive a form of the Stability Lemma from this, and to apply the latter to certain considerations concerning the coproducts of frames.

I am much indebted to Japie Vermeulen for a stimulating correspondence on this subject. For a slightly different treatment related to the Stability Lemma, see [4].

Consider, then, any preclosure operator k_0 on a complete lattice L , with associated closure operator k . For any $a \in L$, let W be the smallest downset (= containing all $y \leq x$ with any x) in $\uparrow a = \{x \in L \mid x \geq a\}$ such that

- (1) $a \in W$,
- (2) W is k_0 -stable, and
- (3) $\bigvee D \in W$ for any updirected $D \subseteq W$.

Then we have

Lemma 1. $W = \{x \in L \mid a \leq x \leq k(a)\}$.

PROOF: Let $V = \{x \in W \mid x \vee y \in W \text{ for all } y \in W\}$. This is a downset since W is. Also, $a \in V$ because $a \vee y = y$ for all $y \in W$. Further, for $x \in V$ and $y \in W$, $k_0(x) \vee y \leq k_0(x \vee y)$, and since $k_0(x \vee y) \in W$ by (2) and the definition of V it follows that $k_0(x) \vee y \in W$, showing that $k_0(x) \in V$. Finally, if $D \subseteq V$ is updirected and $y \in W$ then $E = \{t \vee y \mid t \in D\}$ is an updirected subset of W , hence $\bigvee E \in W$ by (3), but $\bigvee E = (\bigvee D) \vee y$ and therefore $\bigvee D \in V$. It follows now that $V = W$, thus $x \vee y \in W$, for any $x, y \in W$, making W itself updirected so that $s = \bigvee W$ belongs to W . Consequently, by (2), $k_0(s) \leq s$ and hence $s = k_0(s)$. Now, $W \subseteq \{x \in L \mid a \leq x \leq k(a)\}$ since its intersection with the latter still satisfies the conditions (1)–(3), and therefore $a \leq s \leq k(a)$. This implies $s = k(a)$ which proves the lemma. \square

We now apply Lemma 1 to obtain a form of the Stability Lemma. For this, a closure system S in a complete lattice L will be called *finitary* if it is closed under taking joins, in L , of arbitrary updirected subsets. Note that, for the closure operator ℓ associated with S , this condition means that ℓ preserves joins of updirected subsets of L .

Lemma 2. Any finitary closure system in L which is k_0 -stable is also k -stable.

PROOF: Let S be the finitary closure system, with associated closure operator ℓ . Then, for all $x \in L$, $k_0(\ell(x)) \in S$, hence $\ell k_0 \ell(x) = k_0 \ell(x)$, and consequently

$\ell k_0(x) \leq k_0 \ell(x)$. Now for any $a \in S$, let

$$U = \{x \in L \mid a \leq x, \ell(x) \leq k(a)\}.$$

Then U is a downset in $\uparrow a$. Also, $a \in U$ since $a = \ell(a)$. Further, for any $x \in U$, $\ell(x) \leq k(a)$ implies $k_0 \ell(x) \leq k_0(k(a)) = k(a)$ and hence $\ell(k_0(x)) \leq k(a)$, showing that $k_0(x) \in U$. Finally, for any updirected $D \subseteq U$, $\ell[D] \subseteq \downarrow k(a)$, hence $t = \bigvee \ell[D] \leq k(a)$; further, $t \in S$ since S is finitary, and from $\bigvee D \leq t$ it then follows that $\ell(\bigvee D) \leq t \leq k(a)$. This shows $\bigvee D \in U$. As a result, U satisfies the conditions (1)–(3) stated above, hence $W \subseteq U$ and therefore $k(a) \in U$ by Lemma 1. This means that $\ell(k(a)) \leq k(a)$, showing that $k(a) \in S$. \square

As an application of Lemma 2, we now give an improved version of the description of frame coproducts presented in Banaschewski [1]. For general facts concerning frames we refer to Johnstone [3].

Recall that, on a frame L , a *nucleus* is a closure operator such that $k(x \wedge y) = k(x) \wedge k(y)$, and a *prenucleus* is a preclosure operator k_0 for which $k_0(x) \wedge y \leq k_0(x \wedge y)$. The significance of these notions lies in the fact that, for any nucleus k on L , $\text{Fix}(k)$ is a frame such that the map $L \rightarrow \text{Fix}(k)$ given by k is a frame homomorphism, and for any prenucleus on L , the associated closure operator is a nucleus.

Now, for any family $(L_i)_{i \in I}$ of frames, the coproduct may be obtained by suitable constructs originating from the weak product A of the $(L_i)_{i \in I}$ as meet-semilattices. The first stage in this is the lattice \mathcal{D} of all downsets of A ; being closed under arbitrary unions and intersections, \mathcal{D} is certainly a topology and hence a frame. Now, A is not only a meet-semilattice but also has joins, taken componentwise, for arbitrary *updirected* subsets. This suggests the consideration of the Scott-closed subsets of A , that is, the downsets closed under taking joins of arbitrary updirected subsets. These form a closure system \mathcal{S} in \mathcal{D} , obviously determined by the preclosure operator σ_0 such that, for any $U \in \mathcal{D}$,

$$\sigma_0(U) = \{\bigvee D \mid D \subseteq U, \text{updirected}\}.$$

Moreover, σ_0 is a prenucleus, and hence \mathcal{S} is a frame, with frame homomorphism $\mathcal{D} \rightarrow \mathcal{S}$ induced by the associated nucleus σ .

For each $i \in I$ we have a map $k_i : L_i \rightarrow A$ such that $k_i(x)$ has component x for the index i and the unit of L_j for each index $j \neq i$. Then, the map $L_i \rightarrow \mathcal{S}$ taking x to $\downarrow k_i(x)$ preserves all finite meets and updirected joins.

Now, consider a further operator $\tau_0 : \mathcal{D} \rightarrow \mathcal{D}$ such that, for each $U \in \mathcal{D}$, $\tau_0(U)$ consists of all $a \wedge k_i(\bigvee Z)$ for any $a \in A$, $i \in I$ and finite $Z \subseteq L_i$ for which all $a \wedge k_i(t) \in U$, $t \in Z$. This is obviously a preclosure operator, but also easily checked to be a prenucleus. Let $\mathcal{T} = \text{Fix}(\tau_0)$ and τ be the associated nucleus. Note that the maps $L_i \rightarrow \mathcal{T}$ taking $x \in L_i$ to $\tau(\downarrow k_i(x))$ preserve all finite meets and joins.

We are interested in the relationship between the two nuclei σ and τ . Since the definition of τ_0 makes it obvious that \mathcal{T} is a finitary closure system in \mathcal{D} , we can conclude by Lemma 2 that \mathcal{T} is σ -stable provided we show that it is σ_0 -stable. For

this, we first note that the closure condition defining \mathcal{J} can be checked by just taking the cases $Z = \emptyset$ and $Z = \{s, t\}$ for the finite set Z involved — the general case then resulting by obvious induction. Here, the condition for $Z = \emptyset$ requires that all $a \in A$ for which some component is zero belong to the $U \in \mathcal{J}$, and since $U \subseteq \sigma_0(U)$ this also holds for $\sigma_0(U)$. Hence, in order to see that $\sigma_0(U) \in \mathcal{J}$ for any $U \in \mathcal{J}$ it remains to deal with the case $Z = \{s, t\}$. Let, then, $a \wedge k_i(s)$ and $a \wedge k_i(t)$ belong to $\sigma_0(U)$ for some $a \in A$, $i \in I$, and $s, t \in L_i$, and take, accordingly, updirected $D, E \subseteq U$ such that $a \wedge k_i(s) = \bigvee D$ and $a \wedge k_i(t) = \bigvee E$. Now, for any $x = (x_i)_{i \in I}$ in A , define $\bar{x} = \bigwedge \{k_j(x_j) \mid j \neq i\}$ and note that $x = \bar{x} \wedge k_i(x_i)$. Then, for each $x \in D$ and $y \in E$,

$$\bar{x} \wedge \bar{y} \wedge k_i(x_i) \quad \text{and} \quad \bar{x} \wedge \bar{y} \wedge k_i(y_i)$$

belong to U so that

$$\bar{x} \wedge \bar{y} \wedge k_i(x_i \vee y_i) \in U$$

since $U \in \mathcal{J}$. Now, the set of these elements is again updirected and hence

$$b = \bigvee \{\bar{x} \wedge \bar{y} \wedge k_i(x_i \vee y_i) \mid x \in D, y \in E\} \in \sigma_0(U).$$

Finally, since directed joins in A are taken componentwise,

$$\begin{aligned} b &= \bigvee \{\bar{x} \wedge \bar{y} \mid x \in D, y \in E\} \wedge k_i(\bigvee \{x_i \vee y_i \mid x \in D, y \in E\}) \\ &= \bar{a} \wedge k_i(a_i \wedge (s \vee t)) = a \wedge k_i(s \vee t), \end{aligned}$$

showing that the latter elements also belongs to $\sigma_0(U)$, as desired.

The result thus obtained shows that $\tau\sigma\tau = \sigma\tau$, which in turn implies that $\sigma\tau$ is idempotent and therefore a nucleus on \mathcal{D} . Now, Banaschewski [1] describes the coproduct of a family $(L_i)_{i \in I}$ of frames as the closure system \mathcal{L} in \mathcal{S} given by the condition that corresponds to the definition of τ_0 . It follows that $\mathcal{L} = \mathcal{S} \cap \mathcal{J}$, and in all this proves:

Proposition. $\sigma\tau$ is a nucleus on \mathcal{D} such that $\text{Fix}(\sigma\tau)$ is the coproduct of the family $(L_i)_{i \in I}$ of frames, with coproduct maps $L_i \rightarrow \text{Fix}(\sigma\tau)$ taking x to $\sigma\tau(\downarrow k_i(x))$, for each $x \in L_i$ and $i \in I$.

Remark. A crucial stage in the proof in [1] of the localic Tychonoff Theorem that the coproduct of compact frames is compact was the result that, for any family of frames, the nucleus on \mathcal{S} determining $\mathcal{L} = \mathcal{S} \cap \mathcal{J}$ is finitary. Here, this follows from the trivial fact that the nucleus τ on \mathcal{D} is finitary, given that, by the proposition, the nucleus in question is the restriction of $\sigma\tau$. We note that it was at this stage that Bourbaki’s Fixpoint Lemma was used in [1]. The argument here replaces this by Lemma 1 and hence is constructively valid. This amendment makes the results of [1] concerning frame coproducts valid in any topos, provided the family $(L_i)_{i \in I}$ has decidable index set I . The latter restriction enters because the arguments involved here do make use of the condition that $i \neq j$ or $i = j$ for any $i, j \in I$.

As a further application of Lemma 1 we derive an important lemma due to Vermeulen [5].

For any frames L and M , let \mathcal{D} be the frame of all downsets of $L \times M$, σ and τ the nuclei considered earlier, $\varrho = \sigma\tau$, and $\mathcal{K} = \text{Fix}(\varrho)$. Thus \mathcal{K} is the coproduct of L and M , with coproduct maps $L \rightarrow \mathcal{K}$ and $M \rightarrow \mathcal{K}$ given, respectively, by

$$x \rightsquigarrow \varrho(\downarrow(x, e)) \quad \text{and} \quad y \rightsquigarrow \varrho(\downarrow(e, y)).$$

We put

$$x \oplus y = \varrho(\downarrow(x, e)) \cap \varrho(\downarrow(e, y))$$

and note that, for any $U \in \mathcal{D}$,

$$\bigvee \{x \oplus y \mid (x, y) \in U\} = \varrho(U).$$

Further, $U \in \mathcal{D}$ will be called closed under *first* (or *second*) *slice joins* whenever $X \times \{b\} \subseteq U$ implies $(\bigvee X, b) \in U$, for any $X \subseteq L$ and $b \in M$ (or $\{a\} \times Y \subseteq U$ implies $(a, \bigvee Y) \in U$, for any $a \in L$ and $Y \subseteq M$). If X or Y in this condition are restricted to *finite* sets, we refer to *finitary* slice joins.

The result to be proved now is

Vermeulen's Lemma. *For any compact frame L and arbitrary frame M , if $S \in \mathcal{D}$ is closed under finitary first and arbitrary second slice joins then $e \oplus a \leq \bigvee \{x \oplus y \mid (x, y) \in S\}$ implies $(e, a) \in S$.*

PROOF: Consider the set \mathcal{M} of all $U \in \mathcal{D}$ such that $S \subseteq U$ and $(e, z) \in U$ implies $(e, z) \in S$, for all $z \in M$. Clearly, \mathcal{M} is a downset in $\uparrow S$ and $S \in \mathcal{M}$. Further, for any $U \in \mathcal{M}$, let $(e, z) \in \sigma_0(U)$. Then, $(e, z) = \bigvee D$ for some updirected $D \subseteq U$, hence by compactness there exists $(e, t_0) \in D$, and then $z = \bigvee \{t \in M \mid (e, t_0) \in D\}$. Here all $(e, t) \in U$ but since $U \in \mathcal{M}$ also $(e, t) \in S$, and therefore $(e, z) \in S$ by hypothesis on S . This shows that $\sigma_0(U) \in \mathcal{M}$ for all $U \in \mathcal{M}$. Finally, $\bigcup \mathcal{A} \in \mathcal{M}$ for any updirected $\mathcal{A} \subseteq \mathcal{M}$, immediately from the definition of \mathcal{M} . It now follows by Lemma 1 that $\sigma(S) \in \mathcal{M}$. Moreover, since S is closed under finitary first and second slice joins, $\tau(S) = S$ and hence $\varrho(S) = \sigma(S)$ so that, in fact, $\varrho(S) \in \mathcal{M}$. Now $e \oplus a \leq \bigvee \{x \oplus y \mid (x, y) \in S\}$ means $(e, a) \in \varrho(S)$, and we conclude $(e, a) \in S$, as desired. \square

Remark. It might be worth noting that the above proof does not use the full force of the hypothesis on S . It is actually sufficient to have that $S = \tau(S)$, that is, S is closed under all finitary slice joins, and that $\{e\} \times Y \subseteq S$ implies $(e, \bigvee Y) \in S$.

We conclude with a presentation, in slightly different language, of two applications Vermeulen [5] makes of his lemma.

For this, recall that the frame version of the Hausdorff separation axiom for topological spaces is the condition that the codiagonal map $\nabla : L \oplus L \rightarrow L$, given by $\nabla(x \oplus y) = x \wedge y$ be closed, that is, induce an isomorphism $\uparrow s \rightarrow L$ where

$$s = \bigvee \{U \in L \oplus L \mid \nabla(U) = 0\} = \bigvee \{x \oplus y \mid x \wedge y = 0\}.$$

We shall call a frame L *separated* if it satisfies this (although elsewhere such L are also called strongly Hausdorff). It is easy to see that a frame L is separated iff $(e \oplus a) \vee s = (a \oplus e) \vee s$ for all $a \in L$.

Now, the results in question are as follows, with emphasis on the fact that their proofs are constructively valid [5]:

- (R) *Every compact separated frame is regular.*
- (I) *Any dense homomorphism from a separated frame onto a compact frame is an isomorphism.*

PROOF OF (R): Since $(e \oplus a) \vee s = (a \oplus e) \vee s$ one has, for any $a \in L$,

$$\begin{aligned} e \oplus a &\leq \bigvee \{x \oplus y \mid x \leq a \text{ or } x \wedge y = 0\} \\ &\leq \bigvee \{x \oplus y \mid x \leq a \vee y^*\} \leq \bigvee \{x \oplus y \mid (x, y) \in S\} \end{aligned}$$

where $()^*$ stands for pseudocomplement and

$$S = \{(x, y) \mid y \leq \bigvee \{t \mid x \leq a \vee t^*\}\}.$$

Now, S is clearly a downset, closed under arbitrary second slice joins. Moreover it obviously contains $(0, e)$, and if $(x, y), (z, y) \in S$ then

$$y \leq \bigvee \{u \wedge v \mid x \leq a \vee u^*\} \leq \bigvee \{t \mid x \vee z \leq a \vee t^*\}$$

since $u^* \vee v^* \leq (u \wedge v)^*$, and hence $(x \vee z, y) \in S$. This shows S is also closed under finitary first slice joins, and Vermeulen's Lemma then implies that $(e, a) \in S$, meaning

$$a = \bigvee \{t \mid e = a \vee t^*\},$$

which just expresses the regularity of L . □

PROOF OF (I): For separated L and compact M , let $h : L \rightarrow M$ be dense onto. Further, let $k : L \oplus L \rightarrow L \oplus M$ be the homomorphism determined by id_L and h , and $s = \bigvee \{x \oplus y \mid x \wedge y = 0\}$ in $L \oplus L$. Then, for any $a, b \in L$,

$$a \oplus e \leq (e \oplus a) \vee s \text{ and } (e \oplus b) \leq (b \oplus e) \vee s$$

in $L \oplus L$ since L is separated. Now, let $h(a) = h(b)$. Acting k on these two inequalities, one obtains

$$a \oplus e \leq (e \oplus h(a)) \vee k(s) = (e \oplus h(b)) \vee k(s) \leq (b \oplus e) \vee k(s)$$

in $L \oplus M$, and therefore

$$a \oplus e \leq \bigvee \{x \oplus h(y) \mid x \leq b \text{ or } x \wedge y = 0\} \leq \bigvee \{x \oplus h(y) \mid x \leq b \vee y^*\},$$

where y^* is the pseudocomplement of y . Here, $S = \{(x, h(y)) \mid x \leq b \vee y^*\}$ is a downset in $L \times M$, closed under arbitrary first and finitary second slice joins, the latter since (e, o) clearly belongs to S , and if $x \leq b \vee y^*$ and $x \leq b \vee z^*$ then

$$x \leq (b \vee y^*) \wedge (b \vee z^*) = b \vee (y \vee z)^*.$$

Hence Vermeulen's Lemma implies that $(a, e) \in S$, meaning there exist $y \in L$ such that $h(y) = e$ and $a \leq b \vee y^*$. Now

$$0 = h(y \wedge y^*) = h(y) \wedge h(y^*) = h(y^*)$$

shows $y^* = 0$ since h is dense, hence $a \leq b$, and thus $a = b$ by symmetry, as desired. \square

REFERENCES

- [1] Banaschewski B., *Another look at the localic Tychonoff Theorem*, Comment. Math. Univ. Carolinae **29** (1988), 647–656.
- [2] Johnstone P.T., *Topos Theory*, Academic Press, London-New York-San Francisco, 1977.
- [3] ———, *Stone Spaces*, Cambridge University Press, 1982.
- [4] Vermeulen J.J.C., *A note on iterative arguments in a topos*, preprint, 1990.
- [5] ———, *Some constructive results related to compactness and the (strong) Hausdorff property for locales*, preprint, 1991.
- [6] Witt E., *Beweisstudien zum Satz von M. Zorn*, Math. Nachr. **4** (1951), 434–438.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON,
ONT. L8S 4K1, CANADA

(Received December 2, 1991)