

Salvador García-Ferreira; Angel Tamariz-Mascarúa
p-sequential like properties in function spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 35 (1994), No. 4, 753--771

Persistent URL: <http://dml.cz/dmlcz/118717>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

p-sequential like properties in function spaces

SALVADOR GARCÍA-FERREIRA, ANGEL TAMARIZ-MASCARÚA

Abstract. We introduce the properties of a space to be strictly $WFU(M)$ or strictly $SFU(M)$, where $\emptyset \neq M \subset \omega^*$, and we analyze them and other generalizations of p -sequentiality ($p \in \omega^*$) in Function Spaces, such as Kombarov's weakly and strongly M -sequentiality, and Kocinac's $WFU(M)$ and $SFU(M)$ -properties. We characterize these in $C_\pi(X)$ in terms of cover-properties in X ; and we prove that weak M -sequentiality is equivalent to $WFU(L(M))$ -property, where $L(M) = \{\lambda p : \lambda < \omega_1 \text{ and } p \in M\}$, in the class of spaces which are p -compact for every $p \in M \subset \omega^*$; and that $C_\pi(X)$ is a $WFU(L(M))$ -space iff X satisfies the M -version δ_M of Gerlitz and Nagy's property δ . We also prove that if $C_\pi(X)$ is a strictly $WFU(M)$ -space (resp., $WFU(M)$ -space and every RK-predecessor of $p \in M$ is rapid), then X satisfies C'' (resp., X is zero-dimensional), and, if in addition, $X \subset \mathbb{R}$, then X has strong measure zero (resp., X has measure zero), and we conclude that $C_\pi(\mathbb{R})$ is not p -sequential if $p \in \omega^*$ is selective. Furthermore, we show: (a) if $p \in \omega^*$ is selective, then $C_\pi(X)$ is an $FU(p)$ -space iff $C_\pi(X)$ is a strictly $WFU(T(p))$ -space, where $T(p)$ is the set of RK-equivalent ultrafilters of p ; and (b) $p \in \omega^*$ is semiselective iff the subspace $\omega \cup \{p\}$ of $\beta\omega$ is a strictly $WFU(T(p))$ -space. Finally, we study these properties in $C_\pi(Z)$ when Z is a topological product of spaces.

Keywords: selective, semiselective and rapid ultrafilter; Rudin-Keisler order; weakly M -sequential, strongly M -sequential, $WFU(M)$ -space, $SFU(M)$ -space, strictly $WFU(M)$ -space, strictly $SFU(M)$ -space; countable strong fan tightness, Id-fan tightness, property C'' , measure zero

Classification: 04A20, 54C40, 54D55

0. Introduction and preliminaries

In this paper, space will mean a Tychonoff topological space, and $C_\pi(X)$ will denote the space of all the continuous real valued functions defined on X and endowed with the topology of pointwise convergence. For $f \in C_\pi(X)$ and $\delta > 0$, we put $\text{coz}_\delta f = \{x \in X : |f(x)| < \delta\}$. ω is the set of natural numbers equipped with the discrete topology and $\beta\omega$ is the Stone-Ćech compactification of ω which can be viewed as the set of all ultrafilters on ω , where $\hat{A} = \{q \in \beta\omega : A \in q\}$ is a basic open neighborhood of p for each $A \in p$. The remainder $\omega^* = \beta\omega \setminus \omega$ coincides with the set of free ultrafilters on ω and if $f : \omega \rightarrow \beta\omega$ is a function, we will denote by $\bar{f} : \beta\omega \rightarrow \beta\omega$ the Stone extension of f .

We say that $p \in \omega^*$ is *selective* if for every collection $\{A_n\}_{n < \omega}$ of disjoint infinite subsets of ω , with $A_n \notin p$ for every $n < \omega$, there exists $A \in p$ such that $|A \cap A_n| \leq 1$ for every $n < \omega$; $p \in \omega^*$ is *semiselective* if for every sequence $\{A_n\}_{n < \omega} \subset p$ there is $a_n \in A_n$ for each $n < \omega$ such that $\{a_n : n < \omega\} \in p$; and

$p \in \omega^*$ is *rapid* if for every $f \in {}^\omega\omega$ we can find $A \in p$ such that $|A \cap f(n)| \leq n$ for each $n < \omega$. It is not difficult to prove that every selective ultrafilter is semiselective, and every semiselective is a rapid ultrafilter. The existence of these sorts of ultrafilters is independent from the axioms of ZFC; the reader is referred to [Bo], [Ku], [L] and [M].

The *Rudin-Keisler (pre)-order* in ω^* is defined as follows: for $p, q \in \omega^*$, $p \leq_{\text{RK}} q$ if there is $f : \omega \rightarrow \omega$ such that $\bar{f}(q) = p$. If $p \leq_{\text{RK}} q$ and $q \leq_{\text{RK}} p$, then we say that p and q are RK-equivalent (in symbols, $p \simeq_{\text{RK}} q$). It is not difficult to verify that $p \simeq_{\text{RK}} q$ iff there is a permutation σ of ω such that $\bar{\sigma}(q) = p$. The *type* of $p \in \omega^*$ is the set $T(p)$ of all RK-equivalent ultrafilters of p . Observe that the Rudin-Keisler pre-order in ω^* is an order in $\{T(p) : p \in \omega^*\}$. Kunen showed (see [CN, 9.6]) that selective ultrafilters on ω^* are precisely the RK-minimal points of ω^* .

For $p \in \omega^*$, the *p-sum* of a sequence $\{p_n : n < \omega\}$ of free ultrafilters on ω , studied by Frolík ([F]) and, in a more general context, by Vopěnka ([V]) and Katětov ([K]), is the ultrafilter

$$\Sigma_p p_n = \{A \subset \omega \times \omega : \{n < \omega : \{m < \omega : (n, m) \in A\} \in p_n\} \in p\}.$$

Throughout this paper, $\Sigma_p p_n$ will be viewed either as an ultrafilter on ω via a bijection between $\omega \times \omega$ and ω , or as an ultrafilter on $\omega \times \omega$. If $p, q \in \omega^*$ and $p_n = q$ for every $n < \omega$, then $\Sigma_p p_n$ is the usual tensor product $p \otimes q$. Booth [Bo] showed that the induced product $T(p) \otimes T(q) = T(p \otimes q)$, for $p, q \in \omega^*$, produces a semigroup structure in the set of types in ω^* .

For each $\nu \in \omega_1$, we choose an increasing sequence $(\nu(n))_{n < \omega}$ of ordinals in ω_1 such that

- (a) if $2 \leq \nu < \omega$, $\nu(n) = \nu - 1$;
- (b) $\omega(n) = n$ for every $n < \omega$;
- (c) if ν is a limit ordinal, then $\nu(n) \nearrow \nu$;
- (d) if $\nu = \mu + m$ where μ is a limit ordinal and $m < \omega$, then $\nu(n) = \mu(n) + m$ for each $n < \omega$.

For each $p \in \omega^*$, we can define the right powers and the left powers of $T(p)$ as follows (see [Bo] and [GT 2]):

$$T(p)^2 = T(p) \otimes T(p) = {}^2T(p);$$

If $T(p)^\lambda$ and ${}^\lambda T(p)$ have already been defined for every $\lambda < \nu < \omega_1$, then

$$T(p)^\nu = T(p)^\mu \otimes T(p) \quad \text{and} \quad {}^\mu T(p) = T(p) \otimes {}^\mu T(p)$$

for $\nu = \mu + 1$; and, if ν is a limit ordinal,

$$T(p)^\nu = T(\overline{f}(p)) \quad \text{and} \quad {}^\nu T(p) = T(\overline{g}(p)),$$

where $f, g \in \omega \rightarrow \omega^*$ are embeddings defined in such a way that $f(n) \in T(p)^{\nu(n)}$ and $g(n) \in {}^\nu(n)T(p)$ for $n < \omega$.

Observe that, by the associativity of \otimes , ${}^n T(p) = T(p)^n$ for every $n < \omega$, and therefore ${}^\omega T(p) = T(p)^\omega$. On the other hand, it is proved in [Bo, Corollary 2.23] that $T(p)^{\omega+1} <_{\text{RK}} {}^{\omega+1} T(p)$.

If $0 < \nu < \omega_1$ and $p \in \omega^*$, then p^ν and ${}^\nu p$ stand for arbitrary points in $T(p)^\nu$ and ${}^\nu T(p)$, respectively.

The basic properties of these products are the following:

0.1. (1) If $p \in \omega^*$ and $o < \mu < \nu < \omega_1$, then $p^\mu <_{\text{RK}} p^\nu$ ([Bo]) and ${}^\mu p <_{\text{RK}} {}^\nu p$ ([GT 2]).

(2) For each $0 < \mu < \omega_1$ there are $\theta, \tau < \omega_1$ such that $p^\mu \leq_{\text{RK}} \theta_p$ and ${}^\mu p \leq_{\text{RK}} p^\tau$ ([GT 2]).

0.2 Notation. For $\emptyset \neq M \subset \omega$, we set $L(M) = \{\lambda p : \lambda < \omega_1, p \in M\}$ and $R(M) = \{p^\lambda : \lambda < \omega_1, p \in M\}$.

Bernstein introduced in [B] the notion of p -limit of a sequence for $p \in \omega^*$: Let $(x_n)_{n < \omega}$ be a sequence in X . Then $x \in X$ is a p -limit point of $(x_n)_{n < \omega}$ (in symbols, $x = p\text{-lim } x_n$ or $x = p\text{-lim}_{n \rightarrow \infty} x_n$ if it is necessary to emphasize what the indexes of the sequence are being considered) if for each $V \in \mathcal{N}(x)$, $\{n < \omega : x_n \in V\} \in p$. This notion suggests the following definitions which are natural generalizations of the concepts of sequentiality and Fréchet-Urysohn.

0.3 Definitions. Let $\emptyset \neq M \subset \omega^*$ and let X be a space.

(1) (Kombarov [Km]) X is said to be *weakly M -sequential* if for each non-closed subset A of X there are a sequence $(x_n)_{n < \omega}$ in A , $p \in M$ and $x \in X \setminus A$ such that $x = p\text{-lim } x_n$.


(2) (Kombarov [Km]) X is said to be *strongly M -sequential* if for each non-closed subset A of X there are a sequence $(x_n)_{n < \omega}$ in A and $x \in X \setminus A$ such that $x = p\text{-lim } x_n$ for all $p \in M$.

(3) (Kocinac [Ko]) X is a $\text{WFU}(M)$ -space if for every $A \subset X$ and $x \in \text{cl } A$, there are $p \in M$ and a sequence $(x_n)_{n < \omega}$ in A such that $x = p\text{-lim } x_n$.

(4) (Kocinac [Ko]) X is a $\text{SFU}(M)$ -space if for $A \subset X$ and $x \in \text{cl } A$, there is a sequence $(x_n)_{n < \omega}$ in A such that $x = p\text{-lim } x_n$ for every $p \in M$.

Observe that, for $p \in \omega^*$, weakly $\{p\}$ -sequential = strongly $\{p\}$ -sequential and $\text{WFU}(\{p\})$ -space = $\text{SFU}(\{p\})$ -space; in this case, we simply say p -sequential and $\text{FU}(p)$ -space, respectively (the concept of $\text{FU}(p)$ -space was discovered by Comfort and Savchenko independently). We remark that for a space X we have: (a) X is strongly ω^* -sequential if and only if X is sequential; (b) X is weakly ω^* -sequential if and only if X has countable tightness if and only if X is a $\text{WFU}(\omega^*)$ -space; and (c) X is a $\text{SFU}(\omega^*)$ -space if and only if X is Fréchet-Urysohn.

Moreover, if $p \in M \subset \omega^*$, then



Next we give examples to show that the arrows cannot be reversed.

Let $p, q \in \omega^*$ be such that $p <_{\text{RK}} q$, then $\xi(q)$ is a $\text{FU}(q)$ -space and is not p -sequential (see [GF 2]); the sequential space S_2 (see [AF]) is not Fréchet-Urysohn, and its p -version $S_2(p)$ (see [GF 2]) is p -sequential and is not a $\text{FU}(p)$ -space.

If $p \leq_{\text{RK}} q$, then every p -limit point is also a q -limit point, as it is stated in the next lemma.

0.4 Lemma. *Let $(x_n)_{n < \omega}$ be a sequence in a space X such that $p\text{-lim } x_n = x \in X$. If $f : \omega \rightarrow \omega$ is a function such that $\bar{f}(q) = p$, then $x = q - \lim_{n \rightarrow \infty} x_{f(n)}$.*

As a consequence of the previous lemma and 0.1, we have that if $\emptyset \neq M \subset \omega^*$, then

- (a) weak $R(M)$ -sequentiality \Leftrightarrow weak $L(M)$ -sequentiality;
- (b) strong $R(M)$ -sequentiality \Leftrightarrow strong $L(M)$ -sequentiality;
- (c) $\text{WFU}(R(M))$ -space \Leftrightarrow $\text{WFU}(L(M))$ -space;
- (d) $\text{SFU}(R(M))$ -space \Leftrightarrow $\text{SFU}(L(M))$ -space.

We observe next that strong M -sequentiality and weak M -sequentiality imply $\text{SFU}(N)$ -space and $\text{WFU}(N)$ -space for a suitable $N \subset \omega^*$, respectively. The proof of our result resembles the one given in Theorems 3.6 and 3.8 in [GT 2]. We recall that a space X is p -compact, where $p \in \omega^*$, if every sequence in X has a p -limit point in X .

0.5 Theorem. *For $\emptyset \neq M \subset \omega^*$, we have that every weakly M -sequential space is a $\text{WFU}(L(M))$ -space. In addition, if X is p -compact for every $p \in M$, then X is a $\text{WFU}(L(M))$ -space if and only if X is weakly M -sequential.*

If $p \in \omega^*$, then $\xi(p \otimes p)$ is a $\text{WFU}(L(p))$ -space, but it is not p -sequential. Also, as pointed out by the referee, $\xi(p \otimes p)$ is an example of a p^2 -sequential non $\text{SFU}(L(p^2))$ -space.

Theorem 0.5 leads us to ask:

0.6 Problem. For $\emptyset \neq M \subset \omega^*$ and X a space, if $C_\pi(X)$ is a $\text{WFU}(L(M))$ -space (resp. $\text{SFU}(L(M))$ -space) must $C_\pi(X)$ be weakly (resp. strongly) M -sequential?

0.7 Definitions. Let $\emptyset \neq M \subset \omega^*$.

(a) A space Y is a *strictly* $\text{WFU}(M)$ -space if for every sequence $(F_n)_{n < \omega}$ of subsets of Y and every $y \in \bigcap_{n < \omega} \text{cl}_Y F_n$, there exist $y_n \in F_n$ for each $n < \omega$, and $p \in M$ such that $y = p\text{-lim } y_n$.

(b) A space Y is a *strictly* $\text{SFU}(M)$ -space if for every sequence $(F_n)_{n < \omega}$ of subsets of Y and every $y \in \bigcap_{n < \omega} \text{cl}_Y F_n$, there exists $y_n \in F_n$ for each $n < \omega$ such that $y = p\text{-lim } y_n$ for every $p \in M$.

(c) ([S]) A space Y has *countable strong fan tightness* if for every $y \in Y$ and every sequence $(A_n)_{n < \omega}$ of subsets of Y such that $y \in \bigcap_{n < \omega} \text{cl } A_n$, there exists $y_n \in A_n$ for each $n < \omega$, such that $y \in \text{cl}\{y_n : n < \omega\}$.

Note that, for $p \in \omega^*$, a space Y is a strictly $\text{WFU}(\{p\})$ -space $\Leftrightarrow Y$ is a strictly $\text{SFU}(\{p\})$ -space; so, in this case, we say that Y is a *strictly $\text{FU}(p)$ -space*.

Observe also that: (a) Y is a strictly $\text{SFU}(\omega^*)$ -space $\Leftrightarrow Y$ is strictly Fréchet-Urysohn; (b) Y is strictly $\text{WFU}(\omega^*)$ -space $\Leftrightarrow Y$ has countable strong fan tightness; and (c) for $p \in M \subseteq \omega^*$, Y is a strictly $\text{SFU}(M)$ -space $\Rightarrow Y$ is a strictly $\text{FU}(p)$ -space $\Rightarrow Y$ is a strictly $\text{WFU}(M)$ -space $\Rightarrow Y$ has countable strong fan tightness.

A collection \mathcal{G} of subsets of a space X is an ω -cover of X if for every finite subset F of X there exists $G \in \mathcal{G}$ such that $F \subset G$.

The following definition was introduced in [GN 2].

A space X is said to have property γ if for every open ω -cover \mathcal{G} of X , there is a sequence $(G_n)_{n < \omega}$ in \mathcal{G} such that

$$X = \lim G_n = \bigcup_{n < \omega} \bigcap_{n < m} G_m.$$

It is shown in [GN 2] and [G] that the following statements are equivalent:

- (a) $C_\pi(X)$ is a Fréchet-Urysohn space.
- (b) $C_\pi(X)$ is a strictly Fréchet-Urysohn space .
- (c) $C_\pi(X)$ is sequential.
- (d) X satisfies γ .

The equivalence (a) \Leftrightarrow (b) was proved by Pytkeev ([Py]) as well, and Nyikos ([N]) showed that every Fréchet-Urysohn topological group is strictly Fréchet-Urysohn.

In this paper, we principally study the p -sequentiality and the $\text{FU}(p)$ -property on function spaces, and their effects on the base space. This leads us to consider the following generalization of property γ .

0.8 Definition ([GT 1]). X satisfies property γ_p if for each open ω -cover \mathcal{G} of X there is a sequence $(G_n)_{n < \omega}$ in \mathcal{G} such that $X = \lim_p G_n$, where $\lim_p G_n = \bigcup_{A \in p} \bigcap_{n \in A} G_n$.

It is natural to ask whether the p -limit version of the statements (a), (b), (c) and (d), quoted above, are also equivalent for each $p \in \omega^*$. The authors proved in [GT 1] the following.

0.9 Theorem. *Let X be a space and let $p \in \omega^*$. $C_\pi(X)$ is a $\text{FU}(p)$ -space if and only if X satisfies γ_p .*

Unfortunately, the next problem remains unsolved.

0.10 Problem ([GT 1]). *If $C_\pi(X)$ is p -sequential, must X satisfy γ_p ?*

The equivalence between the p -Fréchet-Urysohn property and strictly p -Fréchet-Urysohn property does not hold, in general, on function spaces. In fact, we showed in [GT1] that $C_\pi(\mathbb{R})$ is a $FU(q)$ -space for some $q \in \omega^*$, meanwhile $C_\pi(\mathbb{R})$ cannot be a strictly $FU(p)$ -space for all $p \in \omega^*$ (see Corollary 2.5 below).

Before finishing this section we will give some results concerning limit of sets, sum of ultrafilters and RK-order which will be fundamental for our purposes; their proofs are not difficult, so we have omitted them. If q is an ultrafilter on $\omega \times \omega$ and $(G_{n,m})_{n,m}$ is a bisequence of sets, then the symbol $\lim_q G_{n,m}$ has a clear meaning (see 0.8).

0.11 Lemma. *Let $p \in \omega^*$, $\{q_n\}_{n < \omega} \subset \omega^*$ and let $(G_{n,m})_{n,m < \omega}$ be a bisequence of subsets of a space X . Then*

$$\lim_{\Sigma_p q_n} G_{n,m} = \lim_{n \rightarrow \infty} p \left(\lim_{m \rightarrow \infty} q_n G_{n,m} \right).$$

(Observe that the symbol $\lim_{\Sigma_p q_n} G_{n,m}$ makes sense because $\Sigma_p q_n$ can be considered as an ultrafilter on $\omega \times \omega$.)

0.12 Lemma. *Let $p, q \in \omega^*$ and let $(G_n)_{n < \omega}$ be a sequence of subsets of a space X . If $f : \omega \rightarrow \omega$ is a function satisfying $\bar{f}(q) = p$, then $\lim_p G_n = \lim_{n \rightarrow \infty} q G_{f(n)}$.*

In the next section of this article we give some necessary and sufficient conditions on a space X in order that its function space $C_\pi(X)$ be either $WFU(M)$ -spaces or $SFU(M)$ -spaces, for $\emptyset \neq M \subset \omega^*$. We also prove that $C_\pi(\mathbb{R})$ is not p -sequential if $p \in \omega^*$ is selective. In Section 2 we study the strictly $WFU(M)$ and strictly $SFU(M)$ - function spaces, and we characterize the semiselective ultrafilters in terms of these properties. In the last section we study the product of spaces having one of the cover-properties analyzed in this work.

1. Weakly and strongly Fréchet-Urysohn function spaces and rapid ultrafilters

We are going to characterize the $WFU(M)$ and $SFU(M)$ -properties on a function space $C_\pi(X)$ in terms of cover-properties in X .

1.1 Definitions. Let $\emptyset \neq M \subset \omega^*$.

(1) A space X satisfies property $W\gamma_M$ if for each open ω -cover \mathcal{G} of X , there exist a sequence $(G_n)_{n < \omega}$ in \mathcal{G} and $p \in M$ such that $\lim_p G_n = X$.

(2) A space X satisfies property $S\gamma_M$ if for each open ω -cover \mathcal{G} of X , there is a sequence $(G_n)_{n < \omega}$ such that $\lim_p G_n = X$ for every $p \in M$.

(3) A space X satisfies property WT_M if X has ε and for each sequence $(\mathcal{G}_n)_{n < \omega}$, where $\mathcal{G}_n = \{G_k^n : k < \omega\}$ for $n < \omega$, of open ω -covers of X satisfying $G_k^{n+1} \subset G_k^n$ for each $n, k < \omega$, there exists $p \in M$ and a sequence $(k_m)_{m < \omega}$ of positive integers such that $X = \lim_{m \rightarrow \infty} p G_{k_m}^m$ for each $n < \omega$.

(4) A space X satisfies property ST_M if X has ε and for each sequence $(\mathcal{G}_n)_{n < \omega}$, where $\mathcal{G}_n = \{G_k^n : k < \omega\}$ for $n < \omega$, of open ω -covers of X satisfying $G_k^{n+1} \subset G_k^n$ for each $n, k < \omega$, there is a sequence $(k_m)_{m < \omega}$ of positive integers such that, for every $p \in M$, $X = \lim_{m \rightarrow \infty} {}_p G_{k_m}^n$ for each $n < \omega$.

If $p \in \omega^*$, then we denote by Γ_p the equivalent properties $WT_{\{p\}}$ and $ST_{\{p\}}$.

We recall that a space X satisfies property ε if for each open ω -cover \mathcal{G} of X we can find a countable ω -subcover. It is shown in [GN 2] and [Ar 3] that X has $\varepsilon \Leftrightarrow X^n$ is Lindelöf for every $n < \omega \Leftrightarrow C_\pi(X)$ has countable tightness.

Observe that, if $p \in M \subset \omega^*$, then $S\gamma_M \Rightarrow \gamma_p \Rightarrow W\gamma_M \Rightarrow \varepsilon$ and $S\gamma_M \Rightarrow \Gamma_p \Rightarrow WT_M \Rightarrow \varepsilon$.

In [GT 1] we proved the following four lemmas which will be useful.

1.2 Lemma. Let $\emptyset \neq \Phi \subset C_\pi(X)$ and $f \in \text{cl } \Phi$, and let $\varrho > 0$. Then $\mathcal{G} = \{\text{coz}_\varrho(g - f) : g \in \Phi\}$ is an open ω -cover of X .

1.3 Lemma. Let $f \in C_\pi(X)$, let $\varrho > 0$ and let \mathcal{H} be an open ω -cover of X with $X \notin \mathcal{H}$. If $\Phi = \{g \in C_\pi(X) : \text{coz}_\varrho(g - f) \subset H \text{ for some } H \in \mathcal{H}\}$, then $f \in \text{cl } \Phi \setminus \Phi$.

1.4 Lemma. Let $p \in \omega^*$, let X be a space and let $f, f_0, \dots, f_n, \dots \in C_\pi(X)$. Then $f = p\text{-lim } f_n$ if and only if $\lim_p \text{coz}_\varrho(f_n - f) = X$ for every $\varrho > 0$.

1.5 Lemma. Let $p, X, f, f_0, \dots, f_n, \dots$ as in the previous lemma. Let $(\varrho_n)_{n < \omega}$ be a strictly decreasing sequence of positive real numbers converging to 0. If $X = \lim_p \text{coz}_{\varrho_n}(f_n - f)$, then $f = p\text{-lim } f_n$.

1.6 Theorem. Let X be a space and let $\emptyset \neq M \subset \omega^*$. Then the following statements are equivalent.

- (a) $C_\pi(X)$ is a WFU(M)-space.
- (b) X satisfies $W\gamma_M$.
- (c) X satisfies WT_M .

PROOF: (a) \Rightarrow (b). Let \mathcal{G} be an open ω -cover of X . If $X \in \mathcal{G}$, we take $G_n = X$ for every $n < \omega$ and, obviously, $\lim_p G_n = X$ for every $p \in M$. Now we assume that $X \notin \mathcal{G}$, so $0 \in \text{cl } \Phi \setminus \Phi$ where $\Phi = \{f \in C_\pi(X) : \text{coz}_1 f \subset G \text{ for some } G \in \mathcal{G}\}$ (Lemma 1.3). By hypothesis, there is $p \in M$ and $(f_n)_{n < \omega} \subset \Phi$ such that $\lim_p f_n = 0$. It follows from Lemma 1.4 that $\lim_p \text{coz}_1 f_n = X$. If, for each $n < \omega$, $G_n \in \mathcal{G}$ satisfies that $\text{coz}_1 f_n \subset G_n$, then $\lim_p G_n = X$.

(b) \Rightarrow (c). We know that $W\gamma_M$ implies ε . Let $\mathcal{G}_n = \{G_k^n : k < \omega\}$ be an open ω -cover of X , for each $n < \omega$, such that $G_k^{n+1} \subset G_k^n$ for each $n, k < \omega$. We may assume that X is infinite and hence choose $\{x_n : n < \omega\}$ an infinite subset of X . For each $n < \omega$, define $\mathcal{F}_n = \{G_k^n \setminus \{x_n\} : k < \omega\}$ and put $\mathcal{F} = \bigcup_{n < \omega} \mathcal{F}_n$. It is not hard to prove that \mathcal{F} is an open ω -cover of X as well. By assumption, there is a sequence $(F_j)_{j < \omega}$ in \mathcal{F} and $p \in M$ for which $X = \lim_p F_j$. For each $j < \omega$, we have that $F_j = G_{k_j}^{n_j} \setminus \{x_{n_j}\}$ for some $n_j < \omega$. We claim that $X = \lim_{j \rightarrow \infty} {}_p G_{k_j}^{n_j}$ for each $n < \omega$. In fact, fix $n < \omega$. First, observe that $\{j < \omega : n \leq n_j\} \in p$.

Otherwise, there would be $s < n$ for which $B = \{j < \omega : n_j = s\} \in p$ and so $X = \bigcup_{j \in B} F_j \subset \bigcup \mathcal{F}_s$, a contradiction. If $x \in X$, then $\{j < \omega : x \in F_j\} \in p$ and hence $\{j < \omega : n \leq n_j, x \in G_{k_j}^{n_j}\} \in p$. Since $G_{k_j}^{n_j} \subset G_{k_j}^n$ for $n \leq n_j$, $\{j < \omega : x \in G_{k_j}^n\} \in p$. This proves our claim.

(c) \Rightarrow (a). Let $\Phi \subset C_\pi(X)$ such that $0 \in \text{cl } \Phi \setminus \Phi$. Since X satisfies ε , $C_\pi(X)$ has countable tightness, so we can find $\Psi = \{f_n : n < \omega\} \subset \Phi$ such that $0 \in \text{cl } \Psi$. We take, for each $k < \omega$, $\mathcal{G}_k = \{\text{coz}_{1/2^k} f_n : n < \omega\}$. According to Lemma 1.2, each \mathcal{G}_k is an open ω -cover of X . Besides, $\text{coz}_{1/2^{k+1}} f_n \subset \text{coz}_{1/2^k} f_n$ for every $k, n < \omega$. Thus, by hypothesis, there is $p \in M$ and a sequence $(n_s)_{s < \omega}$ in ω such that $\lim_{s \rightarrow \infty}^p \text{coz}_{1/2^k} f_{n_s} = X$ for every $k < \omega$. Therefore, $0 = p\text{-}\lim_{s \rightarrow \infty} f_{n_s}$ (by Lemma 1.4). □

Analogously, we can prove the strong version of the previous theorem:

1.7 Theorem. *Let X be a space and let $\emptyset \neq M \subset \omega^*$. Then the following statements are equivalent.*

- (a) $C_\pi(X)$ is a SFU(M)-space.
- (b) X satisfies $S\gamma_M$.
- (c) X satisfies $S\Gamma_M$.

A natural cover-property that is closely related to p -sequentiality in function spaces is the p -version δ_p of property δ (δ was introduced and studied in [GN₂]). In order to define δ_p we need some notation:

Let X be a space, $\mathcal{G} \subset \mathcal{P}(X)$ and $p \in \omega^*$. We proceed by induction: Set $S(p, \mathcal{G}, 0) = \mathcal{G}$ and if $S(p, \mathcal{G}, \lambda)$ has been defined for all $\lambda < \mu < \omega_1$, we put $S(p, \mathcal{G}, \mu) = \{\lim_p G_n : (G_n)_{n < \omega}$ is a sequence in $\bigcup_{\lambda < \mu} S(p, \mathcal{G}, \lambda)\}$. Finally, we define $S_p(\mathcal{G}) = \bigcup_{\lambda < \omega_1} S(p, \mathcal{G}, \lambda)$.

1.8 Definition. Let $\emptyset \neq M \subset \omega^*$. A space X satisfies property δ_M if for every open ω -cover \mathcal{G} , X belongs to $S_p(\mathcal{G})$ for some $p \in M$.

We will write δ_p instead of $\delta_{\{p\}}$ for each $p \in \omega^*$.

We will prove that δ_M is the translation in X of WFU($L(M)$)-property in $C_\pi(X)$, where $L(M)$ is the set defined in 0.2. First we will give some lemmas.

- 1.9 Lemma.** (a) *If $\lambda < \mu < \omega_1$, then $S(p, \mathcal{G}, \lambda) \subset S(p, \mathcal{G}, \mu)$;*
 (b) *if $p \leq_{\text{RK}} q$, then $S(p, \mathcal{G}, \lambda) \subset S(q, \mathcal{G}, \lambda)$ for every $\lambda < \omega_1$;*
 (c) *if $\gamma < \mu < \omega_1$, then $S(\gamma p, \mathcal{G}, \lambda) \subset S(\mu p, \mathcal{G}, \lambda)$.*

PROOF: (a) is trivial and (c) is a consequence of (b) and 0.1 (2).

The proof of (b) is by induction. The containment $S(p, \mathcal{G}, 1) \subset S(q, \mathcal{G}, 1)$ follows from Lemma 0.12. Suppose now that $S(p, \mathcal{G}, \lambda) \subset S(q, \mathcal{G}, \lambda)$ for all $\lambda < \mu$. Thus, $S(p, \mathcal{G}, \mu) = \{\lim_p G_n : (G_n)_{n < \omega} \subset \bigcup_{\lambda < \mu} S(p, \mathcal{G}, \lambda)\} \subset \{\lim_p G_n : (G_n)_{n < \omega} \subset \bigcup_{\lambda < \omega} S(q, \mathcal{G}, \lambda)\}$. It follows from Lemma 0.12 that $S(p, \mathcal{G}, \mu) \subset \{\lim_q G_n : (G_n)_{n < \omega} \subset \bigcup_{\lambda < \mu} S(q, \mathcal{G}, \lambda)\} = S(q, \mathcal{G}, \mu)$. □

1.10 Lemma. *Let X be a space, $\mathcal{A} \subset \mathcal{P}(X)$ and $p \in \omega^*$. Then*

- (a) *for every $\nu < \omega_1$, $S(p, \mathcal{A}, \nu) \subset S(\nu^{+1}p, \mathcal{A}, 1)$;*
- (b) *for every $\nu < \omega_1$, $S(\nu p, \mathcal{A}, 1) \subset S(p, \mathcal{A}, \nu)$.*

PROOF: We proceed by transfinite recursion.

(a) The statement is true when $\nu = 1$ because of Lemma 1.9. Suppose that, for every $\nu < \mu$, $S(p, \mathcal{A}, \nu) \subset S(\nu^{+1}p, \mathcal{A}, 1)$. Let $G \in S(p, \mathcal{A}, \mu)$; so $G = \lim_p G_n$ with $(G_n)_{n < \omega}$ a sequence in $\bigcup_{\nu < \mu} S(p, \mathcal{A}, \nu)$. Thus, we obtain that $\{G_n : n < \omega\} \subset S(\mu p, \mathcal{A}, 1)$ (by Lemma 1.9). So, for each $n < \omega$, there is a sequence $(G_{n,m})_{n,m < \omega}$ in \mathcal{A} such that $G_n = \lim_{m \rightarrow \infty} \mu_p(G_{n,m})$. Because of Lemma 0.11, we must have $G = \lim_{n \rightarrow \infty} p(\lim_{m \rightarrow \infty} \mu_p G_{n,m}) = \lim_{\mu+1_p} G_{n,m}$. Thus, $G \in S(\mu^{+1}p, \mathcal{A}, 1)$.

(b) This is evidently true when $\nu = 1$. Suppose that, for every $\nu < \mu$, $S(\nu p, \mathcal{A}, 1) \subset S(p, \mathcal{A}, \nu)$. Let $G \in S(\mu p, \mathcal{A}, 1)$, so $G = \lim_{\mu_p} G_n$ for some sequence $(G_n)_{n < \omega}$ in \mathcal{A} . If $\mu = \lambda + 1$, then there is a bisequence $(G_{n,m})_{n,m < \omega} \subset \mathcal{A}$ such that $G = \lim_{\lambda+1_p} G_n = \lim_{n \rightarrow \infty} p(\lim_{m \rightarrow \infty} \lambda_p G_{n,m})$ (Lemmas 0.11 and 0.12). By induction hypothesis $\lim_{m \rightarrow \infty} \lambda_p G_{n,m} \in S(p, \mathcal{A}, \lambda)$ for each $n < \omega$. Therefore, $G \in S(p, \mathcal{A}, \mu)$.

If μ is a limit ordinal, then $\mu p \simeq_{\text{RK}} \Sigma_p^{\mu(n)} p$ and so, using the results in 0.11 and 0.12 again, we obtain that $G = \lim_{\mu_p} G_n = \lim_{\Sigma_p^{\mu(n)} p} G_n = \lim_{n \rightarrow \infty} p \lim_{m \rightarrow \infty} \mu^{(n)}_p G_{n,m}$ for some bisequence $(G_{n,m})_{n,m < \omega}$ in \mathcal{A} . But $\lim_{m \rightarrow \infty} \mu^{(n)}_p G_{n,m} \in \bigcup_{\nu < \mu} S(p, \mathcal{A}, \nu)$ for every $n < \omega$. Hence, $G \in S(p, \mathcal{A}, \mu)$. □

To prove the following theorem we only need to observe that a space X satisfies $W\gamma_{L(M)}$, for $\emptyset \neq M \subset \omega^*$, if and only if for every open ω -cover \mathcal{G} of X , $X \in \bigcup_{\nu < \omega_1} S(\nu p, 1, \mathcal{G})$ for some $p \in M$, and apply the previous lemma.

1.11 Theorem. *Let $\emptyset \neq M \subset \omega^*$. A space X satisfies δ_M if and only if X satisfies $W\gamma_{L(M)}$ (iff $C_\pi(X)$ is a $\text{WFU}(L(M))$ -space).*

The following lemma follows from Lemma 0.12.

1.12 Lemma. *Let $\emptyset \neq M, N \subset \omega^*$. If for each $p \in M$ there is $q \in N$ such that $p \leq_{\text{RK}} q$, then every space having δ_M has δ_N , and every space having $W\gamma_M$ satisfies $W\gamma_N$.*

From 0.1, 1.11 and 1.12 it follows that:

1.13 Corollary. *Let $p \in \omega^*$ and $\mu, \nu < \omega_1$. All the properties $\delta_p, \delta_{p^\nu}, \delta_{\mu p}, W\gamma_{L(p)}, W\gamma_{L(p^\nu)}, W\gamma_{L(\mu p)}, W\gamma_{R(p)}, W\gamma_{R(p^\nu)}$ and $W\gamma_{R(\mu p)}$ are equivalent.*

Now we are going to study the properties $W\gamma_M$ on \mathbb{R} . In particular we will give some results related to question 0.10.

1.14 Definition. Let Y be a p -sequential space. $\sigma_p(Y)$ will denote the degree of p -sequentiality of Y which is defined to be the least ordinal $\mu \leq \omega_1$ such that for every $A \subset Y$, $\text{cl}_Y A = \bigcup_{\lambda < \mu} A(p, \lambda)$, where

$$A(p, 0) = A, \quad \text{and}$$

$$A(p, \lambda) = \{y \in Y : \exists (y_n)_{n < \omega} \subset \bigcup_{\eta < \lambda} A(p, \eta) \ (y = p\text{-}\lim y_n)\},$$

for $\lambda \leq \omega_1$.

Observe that $\sigma_p(Y)$ is defined iff Y is p -sequential.

As a consequence of Theorem 3.6 in [GT 2] we obtain:

1.15 Theorem. Let $p \in \omega^*$. If $\sigma_p(C_\pi(X)) = \mu < \omega_1$, then $C_\pi(X)$ is a $\text{FU}(\mu+1)p$ -space.

1.16 Corollary. Let $p \in \omega^*$. If $\sigma_p(C_\pi(X)) = \mu < \omega_1$, then X satisfies $\gamma_{\mu+1}p$.

The next result is a generalization of, and its proof is similar to, Lemma 3.15 in [GT 1].

1.17 Lemma. Let $p \in \omega^*$ be selective. Then, for $\lambda < \omega_1$, the RK-predecessors of ${}^\lambda p$ are rapid.

The previous results and Theorem 3.6 in [GT 1] imply, in particular, that $C_\pi(\mathbb{R})$ does not have a degree of p -sequentiality $< \omega_1$ if p is selective. We will obtain a stronger result in 1.20: if p is selective, then $C_\pi(\mathbb{R})$ is not p -sequential.

The following theorem can be proved in a similar way as Theorem 3.6 in [GT 1].

1.18 Theorem. Let $M \subset \omega^*$ such that every RK-predecessor of any element of M is rapid. If $X \subset \mathbb{R}$ satisfies W_{γ_M} , then X has measure zero.

Because of 1.11, 1.17 and 1.18 we obtain

1.19 Theorem. Let $p \in \omega^*$ be selective and $X \subset \mathbb{R}$. If X has δ_p , then X has measure zero.

1.20 Corollary. Let $p \in \omega^*$ be selective and let $X \subset \mathbb{R}$. If $C_\pi(X)$ is p -sequential, then X has measure zero. In particular, $C_\pi(\mathbb{R})$ is not p -sequential.

2. Strictly Fréchet-Urysohn (M)-function spaces and semiselective ultrafilters

2.1 Definitions. Let $\emptyset \neq M \subset \omega^*$.

(a) A space X has strictly W_{γ_M} if for each sequence $(\mathcal{G}_n)_{n < \omega}$ of open ω -covers of X , there are $p \in M$ and, for each $n < \omega$, $G_n \in \mathcal{G}_n$ such that $X = \lim_p G_n$.

(b) A space X has strictly S_{γ_M} if for each sequence $(\mathcal{G}_n)_{n < \omega}$ of open ω -covers of X , there is $G_n \in \mathcal{G}_n$ for each $n < \omega$ such that $X = \lim_p G_n$ for every $p \in M$.

In particular, we say that a space X has *strictly* γ_p if X has strictly $W\gamma_{\{p\}}$ (or, equivalently, if X has strictly $S\gamma_{\{p\}}$).

With a slight modification in the proof of the equivalence (a) \Leftrightarrow (b) of Theorem 1.6, we can show the following result.

2.2 Theorem ([GT 1]). *Let X be a space and $p \in \omega^*$. $C_\pi(X)$ is a strictly $FU(p)$ -space if and only if X has strictly γ_p .*

By applying the same techniques used so far, it is not difficult to prove the following generalization of Theorem 2.2.

2.3 Theorem. *Let X be a space, and let $\emptyset \neq M \subset \omega^*$. Then*

- (a) $C_\pi(X)$ is a strictly $WFU(M)$ -space if and only if X has strictly $W\gamma_M$.
- (b) $C_\pi(X)$ is a strictly $SFU(M)$ -space if and only if X has strictly $S\gamma_M$.

We recall that a space X has property C'' provided that for each sequence $(\mathcal{G}_n)_{n < \omega}$ of open covers of X , there is a sequence $(G_n)_{n < \omega}$, with $G_n \in \mathcal{G}_n$ and $X = \bigcup_{n < \omega} G_n$. A space Y has *Id-fan tightness* if for every $y \in Y$ and every sequence $(A_n)_{n < \omega}$ of subsets of Y such that $y \in \bigcap_{n < \omega} \text{cl } A_n$, there is $F_n \in [A_n]^{\leq n}$ for each $n < \omega$, such that $y \in \text{cl } \bigcup_{n < \omega} F_n$. This concept was introduced in [GT 3] and the authors pointed out that the following result holds (the equivalence between (a) and (c) was shown by Sakai in [S]).

2.4 Theorem. *For a space X , the following are equivalent.*

- (a) $C_\pi(X)$ has countable strong fan tightness.
- (b) $C_\pi(X)$ has Id-fan tightness.
- (c) Each finite product of X has property C'' .

Since every strictly $WFU(M)$ -space, for $\emptyset \neq M \subset \omega^*$, has countable strong fan tightness, then we obtain the following corollary of Theorem 2.4.

2.5 Corollary. *Let $\emptyset \neq M \subset \omega^*$ and let X be a space. If X satisfies strictly $W\gamma_M$, then X^n has C'' for every $n < \omega$. In particular, if X is a subset of \mathbb{R} having strictly $W\gamma_M$, then X has strong measure zero.*

We may also obtain the conclusions in Corollary 2.5 by developing a proof similar to that given in [D, p. 100] of the fact that γ implies Rothberger property C'' (see [GN 2]).

Corollary 2.5 implies that γ_p and strictly $W\gamma_{T(p)}$ are not necessarily equivalent because \mathbb{R} satisfies γ_p for some $p \in \omega^*$ (Theorem 2.15, [GT 1]), but \mathbb{R} does not satisfy strictly $W\gamma_{T(p)}$. Nevertheless, the two concepts coincide when p is selective:

2.6 Theorem. *If $p \in \omega^*$ is selective, then a space X has γ_p if and only if X has strictly $W\gamma_{T(p)}$.*


PROOF: It is not difficult to verify that strictly $W\gamma_{T(p)}$ implies property γ_p for every $p \in \omega^*$. Let $p \in \omega^*$ be selective and assume that X has γ_p . Let $(\mathcal{G}_m)_{m < \omega}$

be a sequence of open ω -covers of X . If X is finite, there is nothing to prove, so we suppose that we can take $\{x_n : n < \omega\} \subset X$ such that $x_i \neq x_j$ if $i \neq j$. For each $m < \omega$, we put $\mathcal{F}_m = \{G - \{x_m\} : G \in \mathcal{G}_m\}$. Then $\mathcal{F} = \bigcup_{m < \omega} \mathcal{F}_m$ is an open ω -cover of X . Since X satisfies γ_p , there is a sequence $(F_n)_{n < \omega}$ in \mathcal{F} such that $X = \lim_p F_n$. The sets $A_k = \{n < \omega : F_n \in \mathcal{F}_k \setminus \bigcup_{i < k} \mathcal{F}_i\}$, for $k < \omega$, constitute a partition of ω . Besides, $A_k \notin q$ for every $k < \omega$ because, on the contrary, if $A_k \in q$ for some $k < \omega$, then $X = \bigcup_{n \in A_k} F_n$ which is in contradiction with the construction of \mathcal{F}_k . Hence, since q is selective, there is $A \in q$ such that $|A \cap A_k| \leq 1$ for every $k < \omega$. By adding points if it is necessary, we may assume that $A \cap A_k = \{n_k\}$ for each $k < \omega$ and hence $A = \{n_k : k < \omega\}$. Thus, the function $f : \omega \rightarrow \omega$ defined by $f(i) = n_i$ is injective and $f[\omega] = A \in p$. Let $q \in \omega^*$ satisfying $\bar{f}(q) = p$. Then, by 0.12, $X = \lim_{k \rightarrow \infty} q F_{n_k}$. Since f is one-to-one, $q \simeq_{\text{RK}} p$ (see [CN] or [C1]). Now, for each $k < \omega$ we choose $G_k \in \mathcal{G}_k$ so that $F_{n_k} \subset G_k$. Then $X = \lim_q G_k$ and $q \in T(p)$. This completes our proof. \square


2.7 Corollary. *Let $p \in \omega^*$ be selective. Then $C_\pi(X)$ is an $\text{FU}(p)$ -space if and only if $C_\pi(X)$ is a strictly $\text{WFU}(T(p))$ -space.*

2.8 Problem. Are γ_p and strictly γ_p the same property whenever p is selective?

2.9. In the following diagrams we summarize some of the results that we have already proved in this article and in [GT 1] concerning generalizations of the FU -property in $C_\pi(X)$ and cover-properties in X . The expression $p \leq_{\text{RK}}^* q$, for $p, q \in \omega^*$, will mean that ${}^\lambda p \leq_{\text{RK}} q$ (equivalently, $p^\lambda \leq_{\text{RK}} q$) for every $\lambda < \omega_1$.



Moreover, if $p \in M \subset \omega^*$, then



2.10 Problems. (1) Under what conditions on $p \in \omega^*$, and on $M \subset \omega^*$ (and on $N \subset \omega^*$), does $W\gamma_M$ imply γ_p (resp., $W\gamma_M$ implies $W\gamma_N$)?

(2) Is p -sequentiality a consequence of δ_p if p is selective?

Now we are going to see that semiselective ultrafilters can be characterized in terms of strictly WFU(M)-properties in $\xi(p)$ -type spaces, where $p \in \omega^*$ and $\xi(p)$ is the subspace $\omega \cup \{p\}$ of $\beta\omega$.

2.11 Lemma. (a) Let $\emptyset \neq M \subset \omega^*$ and $p \in \omega^*$. If $\xi(p)$ is a strictly WFU(M)-space, then p is semiselective.

(b) Let $p, q \in \omega^*$. If $\xi(p)$ is a strictly FU(q)-space, then p is semiselective and $p <_{\text{RK}} q$.

(c) For every $p \in \omega^*$, $\xi(p)$ is not a strictly FU(q)-space for every $q \in T(p)$.

PROOF: (a) Let $\{A_n\}_{n < \omega} \subset p$. For each $n < \omega$ there are $x_n \in A_n$ and $q \in M$ such that $p = q\text{-lim } x_n$, because $p \in \text{cl}_{\xi(p)} A_n$ for all $n < \omega$. Take $A = \{x_n : n < \omega\}$. If $A \notin p$, then $\omega \setminus A \in p$, and so $\emptyset = \{n < \omega : x_n \in \omega \setminus A\} \in q$, a contradiction. Therefore, $A \in p$.

(b) By (a), p is semiselective. Let $A_n \in p$ such that $n \notin A_n$ for $n < \omega$. Then, there is $x_n \in A_n$, for $n < \omega$, such that $p = q\text{-lim } x_n$. If $f(n) = x_n$ for each $n < \omega$, then $\overline{f}(q) = p$. Suppose that f is one-to-one. Let B_n be such that $f(n) \notin B_n$ for each $n < \omega$. By assumption, there is $g : \omega \rightarrow \omega$ such that $g(n) \in B_n$, for all $n < \omega$, and $\overline{g}(q) = \overline{f}(q)$. By [G-F 1, Lemma 2.1], $\{n < \omega : g(n) = f(n)\} \in p$ which is a contradiction. Thus, $p <_{\text{RK}} q$.

(c) is a consequence of (b). □

2.12 Theorem. Let $p \in \omega^*$. p is semiselective if and only if $\xi(p)$ is a strictly WFU($T(p)$)-space.

PROOF: According to Lemma 2.11 (a), we only have to prove the necessity. Assume that $p \in \omega^*$ is semiselective and let $p \in \bigcap_{n < \omega} \text{cl}_{\xi(p)} A_n$, where $A_n \subset \xi(p)$ for every $n < \omega$. Let $B = \{n < \omega : p \in A_n\}$. We have two cases:

First case: $B \in p$.

In this case, for each $n \in B$ we take $x_n = p$, and for every $n \notin B$ we choose some $x_n \in A_n$. If $W \in p$, $\{n < \omega : x_n \in \widehat{W}\} \cap B \in p$; so $p = p\text{-}\lim x_n$.

Second case: $B \notin p$.

Let $\omega \setminus B = \{n_k : k < \omega\}$ such that $n_i < n_j$ if $i < j$. By assumption, for every $k < \omega$, there is $x_{n_k} \in A_{n_k}$ such that $A = \{x_{n_k} : k < \omega\} \in p$. We take $y_0 = x_{n_0}$ and

$$y_{k+1} = \begin{cases} x_{n_{k+1}} & \text{if } x_{n_{k+1}} \notin \{x_{n_s} : 0 \leq s \leq k\} \\ x \in A_{n_{k+1}} \setminus \{x_{n_s} : 0 \leq s \leq k\} & \text{otherwise.} \end{cases}$$

We have that $A \subset \{y_k : k < \omega\} = A'$ and so $A' \in p$. Let $f : \omega \rightarrow A'$ defined by $f(n) = y_k$ if $n = n_k$ and $f(n) = y_0$ if $n \in B$. Let $q \in \omega^*$ such that $\overline{f}(q) = p$. It follows that $p = q\text{-}\lim y_k$. On the other hand, since $f|_{\omega \setminus B}$ is a one-to-one function and $\omega \setminus B \in p$, $p \simeq_{\text{RK}} q$ (see [C1, Lemma 3.2 (c)]). So $\xi(p)$ is a strictly $\text{WFU}(T(p))$ -space. \square

Note that, according to 2.11 (b) and 2.12, for every semiselective ultrafilter p , $\xi(p)$ is a strictly $\text{WFU}(T(p))$ -space which is not a strictly $\text{FU}(p)$ -space.

2.13 Problem. Let $p \in \omega^*$ be selective. Is there any $q \in \omega^*$ such that $p <_{\text{RK}} q$ and $\xi(p)$ is a strictly $\text{FU}(q)$ -space?

3. Products, subspaces and sums

For a nonempty subset M of ω^* , S_M will denote one of the p -sequential like properties that we have considered in this work so far: weakly and strongly M -sequentiality, $\text{WFU}(M)$ and $\text{SFU}(M)$ -property and strictly $\text{WFU}(M)$ and strictly $\text{SFU}(M)$ -property. Let \mathfrak{J} be the set of all S_M with $\emptyset \neq M \subset \omega^*$ plus sequentiality and Fréchet-Urysohn property. In a similar way, C_M is one of the cover properties $W\gamma_M$, $S\gamma_M$, δ_M , strictly $W\gamma_M$ or strictly $S\gamma_M$, and $\mathfrak{C} = \{C_M : \emptyset \neq M \subset \omega^*\} \cup \{\varepsilon, \gamma\}$.

Properties in \mathfrak{J} are not invariants under continuous functions and products, even the square of a space X having $S \in \mathfrak{J}$ does not necessarily satisfy S . In fact, there exists a Fréchet-Urysohn space X whose square $X \times X$ has uncountable tightness ([AR 1], [AR 2]; see also Example 1 in [GN 1]).

On the other hand, the properties in \mathfrak{C} satisfy the following result, proof of which is similar to that given for Theorem 2.1 in [GT 1].

3.1 Theorem. Let $C \in \mathfrak{C}$.

- (a) C is preserved under continuous functions.
- (b) If F is an F_σ -subset of a space X having C , then F has C .
- (c) If X has C , then X_n satisfies C for every $n < \omega$.

3.2 Remarks. (1) Every countable space has C for every $C \in \mathfrak{C}$.

(2) Let X, Y be two spaces and $C \in \mathfrak{C}$. $X \times Y$ has C if and only if the disjoint sum $X \amalg Y$ has C . In fact, $X \amalg Y$ is homeomorphic to a closed subset of $X \times Y$, and $X \times Y$ is homeomorphic to a closed subset of $(X \amalg Y) \times (X \amalg Y)$.

(3) Malykhin and Shakhmatov have shown in [MS] that if we add a single Cohen real to a countable model of $\text{MA} + \neg\text{CH}$, then there exist two spaces X and Y in the generic extension satisfying γ such that $X \times Y$ does not even have ε . On the other hand, as was remarked by van Douwen (see p. 1222 in [C2]), there are (without assuming any additional set theoretic axioms) two spaces X and Y of weight not bigger than 2^ω satisfying ε such that $X \times Y$ is not Lindelöf. Thus, because of Theorem 2.3 in [GT1], X and Y have γ_p for some $p \in \omega^*$ but $X \times Y$ does not satisfy any of the properties in \mathfrak{C} .

3.3 Theorem. *If X satisfies $C \in \mathfrak{C}$ and N is a countable space, then $X \times N$ has C .*

PROOF: Without loss of generality we may suppose that X and N are infinite. The proof for the case $C = \gamma_p$ suffices here. Let $\{a_n : n < \omega\}$ be a faithful indexation of N , let $\{x_n : n < \omega\} \subset X$ such that $x_i \neq x_j$ if $i \neq j$ and let \mathcal{G} be an open ω -cover of $X \times N$. For each $n < \omega$ set $\mathcal{H}_n = \{V \subset X : V \text{ is open and there is } G \in \mathcal{G} \text{ such that } V \times \{a_j : j < n\} \subset G\}$ and $\mathcal{H}'_n = \{V \setminus \{x_n\} : V \in \mathcal{H}_n\}$. We claim that $\mathcal{H} = \bigcup_{n < \omega} \mathcal{H}'_n$ is an open ω -cover of X . In fact, if $\{y_0, \dots, y_r\} \subset X$ there are $s < \omega$ and $G \in \mathcal{G}$ such that $x_s \notin \{y_0, \dots, y_r\}$ and $\{y_0, \dots, y_r\} \times \{a_0, \dots, a_s\} \subset G$. Then, there is an open subset V of X containing $\{y_0, \dots, y_r\}$ such that $V \times \{a_0, \dots, a_s\} \subset G$. So, $V \in \mathcal{H}_s$ and $\{y_0, \dots, y_r\} \subset V \setminus \{x_s\} \in \mathcal{H}'_s$. Thus, \mathcal{H} is an open ω -cover of X . By hypothesis, there is a sequence $(H'_k)_{k < \omega}$ in \mathcal{H} such that $X = \lim_p H'_k$, where $H'_k = V_k \setminus \{x_{n_k}\}$ for some open subset V_k of X and some natural number n_k . For each k there is $G_k \in \mathcal{G}$ such that $V_k \times \{a_j : j < n_k\} \subset G_k$. Now, we are going to prove that $X \times N = \lim_p G_n$. In order to achieve this goal, we take a point $(x, a_n) \in X \times N$. If $D = \{k < \omega : n_k \leq n\} \in p$, then there is $m \leq n$ such that $E = \{k < \omega : n_k = m\} \in p$; but this implies that $X = \bigcup_{k \in E} V_k \setminus \{x_m\}$, which is a contradiction. So, $\{k < \omega : x \in V_k \setminus \{x_{n_k}\}, n < n_k\} \in p$. It follows that $\{k < \omega : (x, a_n) \in V_k \times \{a_j : j < n_k\} \subset G_k \text{ and } n < n_k\} \in p$. \square

3.4 Corollary. *Let $S \in \mathfrak{I} \setminus \{\text{weakly } M\text{-sequentiality, strongly } M\text{-sequentiality} : \emptyset \neq M \subset \omega^*\}$.*

- (a) *If $C_\pi(X)$ satisfies S , then $C_\pi(X^n)$ satisfies S .*
- (b) *If $C_\pi(X)$ satisfies S , then $C_\pi(X)^\omega$ has S .*
- (c) *$C_\pi(X) \times C_\pi(Y)$ has S if and only if $C_\pi(X \times Y)$ has S .*
- (d) *If Y is a quotient space of a space X and Y satisfies $C \in \mathfrak{C}$, then X has C .*

PROOF: We obtain (a) as a consequence of Theorem 3.1; (b) follows from 3.1, 3.3 and the fact that $C(X)^\omega \cong C(X \times \omega)$ (see Corollary 2.4.7 in [MN]); 3.2(2) and

$C_\pi(X) \times C_\pi(Y) \cong C_\pi(X \amalg Y)$ implies (c); and (d) results from Theorems 2.2.8 and 2.2.10 in [MN]. □

It is pointed out in [G, p. 258] that if a space X has γ , then X must be zero-dimensional. For $\emptyset \neq M \subset \omega^*$, the following holds.

3.5 Corollary. *If all RK-predecessors of each $p \in M \subset \omega^*$ are rapid, then every space X with $W\gamma_M$ is zero-dimensional.*

PROOF: Assume that X has property $W\gamma_M$. Let $x \in X$ and U an open neighborhood of x . Choose a continuous function $f : X \rightarrow [0, 1]$ such that $f(x) = 0$ and $f(y) = 1$ for all $y \in X \setminus U$. From 3.1 (a) and 1.18 it follows that $f[X]$ has measure zero. Hence, there is $r \in [0, 1]$ such that $r \notin f[X]$. Then $x \in f^{-1}((0, r)) \subset f^{-1}([0, r]) \subset U$ and $f^{-1}([0, r])$ is a clopen subset of X . □

We know ([GT 1, Theorem 2.3]) that there is $p \in \omega^*$ such that $\beta\omega$ and ω^* have property γ_p . However, if $p \in \omega^*$ is selective, then $\beta\omega$ and ω^* cannot have property γ_p since none of them have property C'' . In fact, by induction we may define a partition $\{A_{t(0), \dots, t(n-1)} : t : n \rightarrow \{0, 1\} \text{ is a function}\}$ of ω in infinite subsets such that $\{A_{t(0), \dots, t(n-1), 0}, A_{t(0), \dots, t(n-1), 1}\}$ is a partition of $A_{t(0), \dots, t(n-1)}$ in infinite sets, for each $n < \omega$ and each function $t : n \rightarrow \{0, 1\}$. Now define, for each $n < \omega$, $\mathcal{G}_n = \{\hat{A}_{t(0), \dots, t(n-1)} : t : n \rightarrow \{0, 1\} \text{ is a function}\}$. We have that \mathcal{G}_n is a cover of $\beta\omega$. Assume that $\beta\omega$ has property C'' . Then, for every $n < \omega$ there is $t_n : n \rightarrow \{0, 1\}$ such that $\beta\omega = \bigcup_{n < \omega} \hat{A}_{t_n(0), \dots, t_n(n-1)}$. We may find $\sigma : \omega \rightarrow \{0, 1\}$ so that $\sigma(n-1) \neq t_n(n-1)$ for all $1 < n < \omega$. We may choose $q \in [\bigcap_{n < \omega} \hat{A}_{\sigma(0), \dots, \sigma(n-1)}] \cap \omega^*$. It is then evident that $q \notin \hat{A}_{t_n(0), \dots, t_n(n-1)}$ for all $1 < n < \omega$, a contradiction. □

Using analogous proofs of those given for Theorems 3.18 and 3.20 in [GT 1], we have

3.6 Theorem. *Let $C \in \mathfrak{C}$.*

- (1) *the following are equivalent.*
 - (a) \mathbb{R} satisfies C .
 - (b) The Cantor space 2^ω satisfies C .
 - (c) Every metrizable separable locally compact space has C .
- (2) *the following are equivalent.*
 - (a) \mathbb{R}^ω satisfies C .
 - (b) The set of irrational numbers ω^ω satisfies C .
 - (c) Every completely metrizable space has C .

If $|X| > 1$, then the Cantor set 2^ω is homeomorphic to a closed subspace of X^ω . On the other hand, 2^ω does not have γ ([GN 2, Theorem 6]). Thus, we obtain

3.7. *If X has more than one point, then X^ω does not satisfy γ .*

Note also that if X^ω has $C \in \mathfrak{C}$, then X has C , because X is homeomorphic to some closed subset of X^ω . But the converse is not necessarily true, for example

the discrete space $\{0, 1\}$ satisfies C for every $C \in \mathfrak{C}$ but 2^ω does not have γ_p if p is semiselective ([GT 1, Theorems 3.9 and 3.18]). We obtain something different when we take X into some convenient class of spaces:

3.8 Theorem. *Let X be a compact second countable and non-countable space and let $C \in \mathfrak{C} \setminus \{\gamma\}$, then X has C if and only if X^ω has C .*

PROOF: Since X is second countable and non-countable, there is $\emptyset \neq F \subset X$ which is perfect (see [E, p. 59, 1.7.11]). Furthermore, F is completely metrizable because X is, then F contains a copy of the Cantor set ([E, p. 290, 4.5.5]). Thus, 2^ω satisfies C . By Theorem 3.6, every compact metric space has C ; in particular X^ω has C . \square

3.9 Corollary. *Let X be a compact second countable non-countable space, and let $S \in \mathfrak{J} \setminus \{\text{weakly } M\text{-sequentiality, strongly } M\text{-sequentiality} : \emptyset \neq M \subset \omega^*\}$. Then $C_\pi(X)$ has S if and only if $C_\pi(X^\omega)$ satisfies S .*

3.10 Problems. (1) Let $C \in \mathfrak{C}$. Does \mathbb{R}^ω satisfy C if \mathbb{R} does?

(2) Let $p \in \omega^*$ be a P -point and let X be a space having γ . Does X^ω satisfy γ_p ?

3.11 Examples. Let $C \in \mathfrak{C}$.

(1) If X contains a closed discrete set of cardinality $> \aleph_0$, then X does not satisfy C . So, the Moore Plane and $L \times L$, where L is the Sorgenfrey Line do not satisfy C , and by Theorem 3.1, L does not satisfy C either. Also for every cardinal $\alpha \geq \omega$, $\alpha^{(\alpha^+)}$ does not have C . In particular, if X contains a closed copy of ω , X^α does not satisfy C if $\alpha > \omega$.

(2) Let X be a space satisfying ε and such that X contains a point x_0 with the property that for every neighborhood V of x_0 , $|X \setminus V| \leq \aleph_0$. Then X satisfies C . Thus, the one point compactification of a discrete space and $[o, \omega_1]$ (with the order topology) satisfy C .

Because of Theorem 2.12 in [GT 1] we know that 2^{2^ω} has γ_p for some $p \in \omega^*$. So it is natural to ask:

3.12 Problem. Does $2^{(2^\omega)^+}$ satisfy γ_p for any $p \in \omega^*$?

For a space X , $n(X)$ denotes the Novak number of X (that is, $n(X)$ is the smaller power of a family of nowhere dense sets covering X). Our last result is a consequence of Theorem 1 in [Ma].

3.13 Theorem ($n(\omega^*) > c$). *Let X be a space. The following statements are equivalent.*

- (a) X satisfies γ .
- (b) X satisfies γ_p for every $p \in \omega^*$.
- (c) X satisfies strictly γ_p for every $p \in \omega^*$.
- (d) $C_\pi(X)$ is p -sequential for every $p \in \omega^*$.

- 3.14 Problems.** (1) Is it consistent with ZFC that every space satisfying δ_p for every $p \in \omega^*$ has γ_p for every $p \in \omega^*$?
- (2) Is it consistent with ZFC that there exists a space X satisfying γ_p (resp., δ_p , strictly γ_p) for every $p \in \omega^*$ and $\neg\gamma$?
- (3) Is it consistent with ZFC that \mathbb{R} satisfies γ_p for every $p \in \omega^*$?

REFERENCES

- [AR 1] Arhangel'skii A.V., *The spectrum of frequencies of topological spaces and their classification* (in Russian), Dokl. Akad. Nauk SSSR **206** (1972), 265–268.
- [AR 2] Arhangel'skii A.V., *The spectrum of frequencies of a topological space and the product operation* (in Russian), Trudy Moskov. Mat. Obsc. **40** (1979), 171–266.
- [AR 3] Arhangel'skii A.V., *Structure and classification of topological spaces and cardinal invariants*, Russian Math. Surveys **33** (1978), 33–96.
- [AF] Arhangel'skii A.V., Franklin S.P., *Ordinal invariants for topological spaces*, Michigan Math. J. **15** (1968), 313–320.
- [B] Bernstein A.R., *A new kind of compactness for topological spaces*, Fund. Math. **66** (1970), 185–193.
- [Bo] Booth D., *Ultrafilters on a countable set*, Ann. Math. Logic **2** (1970), 1–24.
- [C 1] Comfort W.W., *Ultrafilters: some old and some new results*, Bull. Amer. Math. Soc. **83** (1977), 417–455.
- [C 2] ———, *Topological groups*, in K. Kunen and J.E. Vaughan, editors, Handbook of Set-Theoretic Topology, North-Holland, 1984.
- [CN] Comfort W.W., Negrepointis S., *The Theory of Ultrafilters*, Springer Verlag, Berlin-Heidelberg-New York, 1974.
- [D] Daniels P., *Pixley-Roy spaces over subsets of reals*, Top. Appl. **29** (1988), 93–106.
- [E] Engelking R., *General Topology*, Sigma Series in Pure Math., vol. 6, Heldermann Verlag, Berlin, 1989.
- [F] Frolík Z., *Sums of ultrafilters*, Bull. Amer. Math. Soc. **73** (1967), 87–91.
- [G-F 1] García-Ferreira S., *Comfort types of ultrafilters*, Proc. Amer. Math. Soc. **120** (1994), 1251–1260.
- [G-F 2] ———, *On $FU(p)$ -spaces and p -sequential spaces*, Comment. Math. Univ. Carolinae **32** (1991), 161–171.
- [G-T 1] García-Ferreira S., Tamariz-Mascarúa A., *p -Fréchet-Urysohn property of function spaces*, to be published in Top. and Appl.
- [G-T 2] ———, *On p -sequential p -compact spaces*, Comment. Math. Univ. Carolinae **34** (1993), 347–356.
- [G-T 3] ———, *Some generalizations of rapid ultrafilters in topology and Id -fan tightness*, to be published in Tsukuba Journal of Math.
- [G] Gerlitz J., *Some properties of $C(X)$, II*, Topology Appl. **15** (1983), 255–262.
- [GN 1] Gerlitz J., Nagy Zs., *Products of convergence properties*, Comment. Math. Univ. Carolinae **23** (1982), 747–766.
- [GN 2] ———, *Some properties of $C(X)$, I*, Topology Appl. **14** (1982), 151–161.
- [K] Katětov M., *Products of filters*, Comment. Math. Univ. Carolinae **9** (1968), 173–189.
- [Ko] Kocinac L.D., *A generalization of chain-net spaces*, Publ. Inst. Math. (Beograd) **44** (58), 1988, pp. 109–114.
- [Km] Kombarov A.P., *On a theorem of A.H. Stone*, Soviet Math. Dokl. **27** (1983), 544–547.
- [Ku] Kunen K., *Some points in βN* , Proc. Cambridge Philos. Soc. **80** (1976), 358–398.
- [L] Laflamme C., *Forcing with filters and complete combinatorics*, Ann. Math. Logic **42** (1989), 125–163.

- [Ma] Malykhin V.I., *The sequentiality and Fréchet-Urysohn property with respect to ultrafilters*, Acta Univ. Carolinae Math. et Phy. **31** (1990), 65–69.
- [MS] Malykhin V.I., Shakmatov D.D., *Cartesian products of Fréchet topological groups and function spaces*, Acta Math. Hung. **60** (1992), 207–215.
- [MN] McCoy R.A., Ntanty I., *Topological Properties of Spaces of Continuous functions*, Lecture Notes in Math. **1315**, Springer Verlag, 1980.
- [M] Miller A.W., *There are no Q -points in Laver's model for the Borel conjecture*, Proc. Amer. Math. Soc. **78** (1980), 103–106.
- [N] Nyikos P.J., *Metrizability and the Fréchet-Urysohn property in topological groups*, Proc. Amer. Math. Soc. **83** (1981), 793–801.
- [Py] Pytkeev E.G., *On sequentiality of spaces of continuous functions*, Russian Math. Surveys **37** (1982), 190–191.
- [S] Sakai M., *Property C'' and function spaces*, Proc. Amer. Math. Soc. **104** (1988), 917–919.
- [V] Vopěnka P., *The construction of models of set theory by the method of ultraproducts*, Z. Math. Logik Grundlagen Math. **8** (1962), 293–304.

INSTITUTO DE MATEMÁTICAS, CIUDAD UNIVERSITARIA (UNAM), D.F. 04510 MEXICO, MEXICO

E-mail: garcia@redvax1.dgsca.unam.mx

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, CIUDAD UNIVERSITARIA (UNAM), D.F. 04510 MEXICO, MEXICO

E-mail: atamarif@condor.dgsca.unam.mx

(Received June 15, 1992, revised May 20, 1993)