

Jiří Jelínek

Equality of two diffeomorphism invariant Colombeau algebras

Commentationes Mathematicae Universitatis Carolinae, Vol. 45 (2004), No. 4, 633--662

Persistent URL: <http://dml.cz/dmlcz/119491>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2004

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Equality of two diffeomorphism invariant Colombeau algebras

JIŘÍ JELÍNEK

Abstract. The two diffeomorphism invariant algebras introduced in Grosser M., Farkas E., Kunzinger M., Steinbauer R., *On the foundations of nonlinear generalized functions I, II*, Mem. Amer. Math. Soc. **153** (2001), no. 729, 93 pp., are identical.

Keywords: Colombeau algebra of generalized functions, representative, diffeomorphism invariance

Classification: 46F, 46F05

The paper is a continuation of [9] and its only aim is to prove that both diffeomorphism invariant Colombeau-type algebras \mathcal{G}^d and \mathcal{G}^2 introduced in [8] and [6] (Grosser et al.) coincide. In [6] diverse possibilities to define Colombeau-type algebras are researched; our result shows that there is only one diffeomorphism invariant Colombeau-type algebra among them for a given domain $\Omega \subset \mathbb{R}^d$.

§1. In this paper, we use notations introduced in [9] and mostly we refer to [9]. This paper is devoted to prove the following

Theorem. *For every open $\Omega \subset \mathbb{R}^d$, the algebras $\mathcal{G}^2(\Omega)$ and $\mathcal{G}^d(\Omega)$ coincide.*

As the algebras are quotient algebras $\mathcal{G}^2 := \mathcal{E}_M^2/\mathcal{N}$ and $\mathcal{G}^d := \mathcal{E}_M^d/\mathcal{N} \cap \mathcal{E}_M^d$, the theorem says that for any representative $R \in \mathcal{E}_M^2$ another representative $\tilde{R} \in \mathcal{E}_M^d$ can be found with $R - \tilde{R} \in \mathcal{N}$. To prove it, in all what follows, we assume that $R \in \mathcal{E}_M^2$ is given and we are going to construct \tilde{R} . This will be done in several steps. In every step functions of variables φ, x are constructed and their properties are presented with the aim to construct at last the required representative \tilde{R} . First we show that it is sufficient to do it for representatives with compact support.

Proposition. *Suppose that for any representative $R \in \mathcal{E}_M^2(\mathbb{R}^d)$ such that there is a compact $K \Subset \mathbb{R}^d$ fulfilling $R(\varphi, x) = 0$ whenever $x \in \mathbb{R}^d \setminus K$, a representative*

Partially supported by the grant GAČR 201/00/0767 and partially by the grant MSM 113200007.

$\tilde{R} \in \mathcal{E}_M^d(\mathbb{R}^d)$ can be found with $R - \tilde{R} \in \mathcal{N}$. Then for any open $\Omega \subset \mathbb{R}^d$ the algebras $\mathcal{G}^2(\Omega)$ and $\mathcal{G}^d(\Omega)$ coincide.

PROOF: Let $R \in \mathcal{E}_M^2(\Omega)$. Recall that unlike in [8] and [6] here a representative is defined on $\mathcal{E}(\Omega) = \mathcal{A}_0(\mathbb{R}^d) \times \Omega$ and we do not lose generality with this assumption. Choose a locally finite covering

$$\Omega = \bigcup \Omega_m \quad \text{with} \quad \overline{\Omega_m} \Subset \Omega$$

and a partition of unity $1 = \sum \chi_m$ on Ω subordinated to this covering, $\chi_m \in \mathcal{D}(\Omega_m)$, $K_m := \text{supp } \chi_m$. Then $R = \sum R_m$ if we denote $R_m(\varphi, x) := R(\varphi, x) \cdot \chi_m(x)$ and we have $R_m(\varphi, x) = 0$ whenever $x \in \Omega \setminus K_m$. R_m can be easily extended to belong to $\mathcal{E}_M^2(\mathbb{R}^d)$ putting $R_m(\varphi, x) = 0$ whenever $x \in \mathbb{R}^d \setminus K_m$. By hypothesis, we can find $\tilde{R}_m \in \mathcal{E}_M^d(\mathbb{R}^d)$ with $\tilde{R}_m - R_m \in \mathcal{N}$. Then, for every m , we choose a test function $\sigma_m \in \mathcal{D}(\Omega_m)$ that is $= 1$ on a neighbourhood of K_m . The functions χ_m and σ_m are considered to be elements of $\mathcal{E}(\mathbb{R}^d)$ as functions independent of the first variable. Consequently, $\sigma_m \in \mathcal{E}_M^d$. Considering all representatives to be elements of $\mathcal{E}_M^2(\Omega)$ (i.e. restricted to $\mathcal{A}_0(\mathbb{R}^d) \times \Omega$), we have (note that \mathcal{N} is an ideal) $R - \sum \tilde{R}_m \sigma_m = \sum (R_m - \tilde{R}_m \sigma_m) = \sum (R_m - \tilde{R}_m) \sigma_m \in \mathcal{N}$, the sum being locally finite. From the same reason, $\tilde{R} := \sum \tilde{R}_m \sigma_m \in \mathcal{E}_M^d$. \tilde{R} is thus a required representative. \square

§2. Remark. For $B \in \mathbb{R}^d$, it is known that $\mathcal{D}(B)$ is a Fréchet space. Its topology can be generated by a countable system of norms defined by continuous scalar products, e.g.

$$\varphi, \psi \mapsto \int \frac{\partial^{dm}}{\partial \xi_1^m \dots \partial \xi_d^m} \varphi(\xi) \cdot \frac{\partial^{dm}}{\partial \xi_1^m \dots \partial \xi_d^m} \overline{\psi}(\xi) \, d\xi.$$

A continuous scalar product $\varphi, \psi \mapsto (\varphi, \psi)$ is \mathcal{C}^∞ , being sesqui-linear, and we have

$$\begin{aligned} d_\psi(\varphi, \varphi) &= 2\Re(\varphi, \psi), \\ d_{\psi_1, \psi_2}^2(\varphi, \varphi) &= 2\Re(\psi_1, \psi_2); \end{aligned}$$

the derivatives of higher orders are zero. Hence the norm generated by a continuous scalar product is \mathcal{C}^∞ in all points except of origin. The function $\psi \mapsto (\psi, \psi) = \|\psi\|^2$ is \mathcal{C}^∞ always.

Notation. The function V_N , used in Equivalent Definition [9, §7, (4°) and (5°)] of \mathcal{E}_M^2 , can be

$$(1) \quad V_N(\varphi) = \left(\sum_{\substack{\beta \in \mathbb{N}_0^d \\ 1 \leq |\beta| \leq N}} \left| \int \xi^\beta \varphi(\xi) \, d\xi \right|^2 \right)^{1/2} \quad (\varphi \in \mathcal{A}_0).$$

Evidently, this function fulfills [9, (6)]:

$$\forall N \in \mathbb{N}, \mathcal{B} \text{ (bounded)} \subset \mathcal{A}_0 \quad \exists C_1, C_2 > 0 \quad \forall \varphi \in \mathcal{B} :$$

$$C_2 \sum_{\substack{\beta \in \mathbb{N}_0^d \\ 1 \leq |\beta| \leq N}} \left| \int \xi^\beta \varphi(\xi) \, d\xi \right| \leq V_N(\varphi) \leq C_1 \sum_{\substack{\beta \in \mathbb{N}_0^d \\ 1 \leq |\beta| \leq N}} \left| \int \xi^\beta \varphi(\xi) \, d\xi \right|.$$

Evidently every multiple $c \cdot V_N(\varphi)$ satisfies these inequalities, so it can be used in Equivalent Definition [9, §7, (4°) and (5°)]. Also the function

$$V'_N(\varphi) = \left(\sum_{\substack{\beta \in \mathbb{N}_0^d \\ 1 \leq |\beta| \leq N}} \|\varphi\|_{\mathcal{L}^2}^{4|\beta|/d} \left| \int \xi^\beta \varphi(\xi) \, d\xi \right|^2 \right)^{1/2} \quad (\varphi \in \mathcal{A}_0)$$

fulfills the above inequalities [9, (6)] that allows us to use it in Equivalent Definitions [9, §7, (4°), (5°)]. Indeed, a bounded set is relatively compact, so $\exists c_1, c_2 > 0$ (depending on \mathcal{B}) $\forall \varphi \in \mathcal{B}$ we have $c_2 \leq \|\varphi\|_{\mathcal{L}^2} \leq c_1$; [9, (6)] follows easily. It can be checked that

$$(2) \quad \|S_\varepsilon \varphi\|^{-2/d} = \varepsilon \cdot \|\varphi\|^{-2/d}, \quad \int S_\varepsilon \varphi(\xi) \xi^\beta \, d\xi = \varepsilon^{|\beta|} \int \varphi(\xi) \xi^\beta \, d\xi \quad (\beta \in \mathbb{N}_0^d),$$

so $V'_N(S_\varepsilon \varphi) = V'_N(\varphi)$.

For all what follows, a function $\rho \in \mathcal{A}_0([-1, 1])$ is fixed such that

$$\rho(\xi) > 0 \text{ iff } \xi \in]-1, 1[,$$

$$\rho_\varepsilon := S_\varepsilon \rho, \quad \vartheta := \rho_{1/2} * \chi_{[-3/2, 3/2]}$$

(convolution with the characteristic function),

$$\vartheta_m(\xi) := \vartheta(2^{-m}\xi) \quad (m \text{ integer}).$$

So $\vartheta_m(\xi) = 1$ iff $\xi \in [-2^m, 2^m]$, $0 < \vartheta_m(\xi) \leq 1$ iff $\xi \in]-2^{m+1}, 2^{m+1}[$ and ϑ_m is decreasing on $[2^m, 2^{m+1}]$. Denote furthermore

$$K_m := [-2^m, 2^m]^d$$

and for $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$, denote $\vartheta_m^{\otimes d}(\xi) := \vartheta_m(\xi_1) \dots \vartheta_m(\xi_d)$. If there is no danger of confusion, we will write simply $\vartheta_m(\xi)$ instead of $\vartheta_m^{\otimes d}(\xi)$.

§3. Thanks to §1, Proposition we can assume that the given representative R belongs to $\mathcal{E}_M^2(\mathbb{R}^d)$ and that there is a compact $K \Subset \Omega$ fulfilling $R(\varphi, x) = 0$ for $x \in \mathbb{R}^d \setminus K$. In this case, in the equivalent definitions of \mathcal{E}_M^2 and \mathcal{N} ([9, §§7, 8]) we can omit $\forall K \Subset \Omega$ and replace the uniformity on K with the uniformity on the whole of \mathbb{R}^d . Denote by N_L the number N from Equivalent Definition [9, §7, (5°)] holding at the same time for all $|\alpha| \leq L$ and for all differentials of order $k \leq L$. Certainly, this equivalent definition remains valid if we take any greater number for N_L . We replace our representative with another one determining the same generalized function, if needed, to obtain the following

Properties of R .

- (1°) There is an increasing sequence $\{N_L\}_{L \in \mathbb{N}} \subset \mathbb{N}$, $N_L \geq L$, fulfilling:
 $\forall B \Subset \mathbb{R}^d$, \mathcal{B} (bounded) $\subset \mathcal{A}_0(B)$, $L \in \mathbb{N} \quad \exists \mathcal{U}$ (absolutely convex open neighbourhood of zero) $\subset \mathcal{A}(B)$, $C > 0 \quad \forall \ell = 1, 2, \dots, L$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$, $\varphi \in \mathcal{B} + 2\mathcal{U}$, $\varepsilon \in]0, 1]$, $\varepsilon^{N_L} \geq V_{N_L}(\varphi)$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L$, $x \in \mathbb{R}^d$:

$$(3) \quad \begin{aligned} |\partial^\alpha (d_{S_\varepsilon \psi_1, \dots, S_\varepsilon \psi_\ell}^\ell R)(S_\varepsilon \varphi, x)| &= |\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R(S_\varepsilon \varphi, x)| \leq \varepsilon^{-N_L}, \\ |\partial^\alpha R(S_\varepsilon \varphi, x)| &\leq C \varepsilon^{-N_L}. \end{aligned}$$

- (2°) The first inequality in (3) can be written in the form

$$(4) \quad |\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R(S_\varepsilon \varphi, x)| \leq \varepsilon^{-N_L} \|\psi_1\|_{\mathcal{U}} \dots \|\psi_\ell\|_{\mathcal{U}}$$

if we omit the hypothesis $\psi_1, \dots, \psi_\ell \in \mathcal{U}$, only supposing $\psi_1, \dots, \psi_\ell \in \mathcal{A}(B)$ ($\|\bullet\|_{\mathcal{U}}$ denotes the Minkowski functional assigned to \mathcal{U}).

- (3°) If $L = 1$, the hypothesis $\varepsilon^{N_L} \geq V_{N_L}(\varphi)$ can be omitted, so that (3) and (4) hold for every $\varepsilon \in]0, 1]$.
 (4°) Consequently, if \mathcal{B} is convex, $\varphi_1, \varphi_2 \in \mathcal{B} + 2\mathcal{U}$, we have to consider two cases.

If $L = 1$, $|\alpha| \leq 1$ then

$$(5) \quad |\partial^\alpha (R(S_\varepsilon \varphi_2, x) - R(S_\varepsilon \varphi_1, x))| \leq \varepsilon^{-N_1} \|\varphi_2 - \varphi_1\|_{\mathcal{U}}.$$

Otherwise if $|\alpha| \leq L$ and $\varepsilon^{N_L} \geq V_{N_L}(\varphi_1)$, $\varepsilon^{N_L} \geq V_{N_L}(\varphi_2)$, $\ell = 1, \dots, L$, $\psi_1, \dots, \psi_{\ell-1} \in \mathcal{A}(B)$, then

$$(6) \quad \begin{aligned} |\partial^\alpha d_{\psi_1, \dots, \psi_{\ell-1}}^{\ell-1} (R(S_\varepsilon \varphi_2, x) - R(S_\varepsilon \varphi_1, x))| \\ \leq \varepsilon^{-N_L} \|\psi_1\|_{\mathcal{U}} \dots \|\psi_{\ell-1}\|_{\mathcal{U}} \|\varphi_2 - \varphi_1\|_{\mathcal{U}}. \end{aligned}$$

PROOF OF (3°): The items (1°) and (2°) are consequences of [9, §7, (5°)]. The equality in (3) follows from the chain rule where the inner function is linear, so

its higher derivatives vanish. (2°) follows from the linearity of differentials. This holds for any representative with compact support. However, for (3°) we have to choose a suitable representative determining a given generalized function and possibly we have to choose the number N_1 , too. Let R be a given representative. Applying $\frac{1}{2}\|\chi_B\|^{-2N_1/d} V'_{N_1}$ instead of V_{N_1} in Equivalent Definition [9, §7, (5°)] (or in Properties, item (1°)) for $L = 1$, we get \mathcal{U} and $C > 0$ such that (3) holds if $\ell = 1$, $|\alpha| \leq 1$, $\psi_1 \in \mathcal{U}$, $\varphi \in \mathcal{B} + 2\mathcal{U}$ and

$$(7) \quad \varepsilon^{N_1} \geq \frac{1}{2}\|\chi_B\|^{-2N_1/d} V'_{N_1}(\varphi).$$

Let us define $R'(\varphi, x) := \vartheta(\|\varphi\|^{2N_1/d} V'_{N_1}(\varphi)) \cdot R(\varphi, x)$ (\mathcal{L}^2 norms).

First we prove that $R' \in \mathcal{E}_M^2$ and $R - R' \in \mathcal{N}$. Let $\varepsilon \mapsto \varphi^\varepsilon$ (see [9, §7, (2°), §8, (4°)]) be a bounded path with asymptotically vanishing moments of order $N_1 + 1$. This means that the set $\{\varphi^\varepsilon; \varepsilon \in]0, 1]\}$ is bounded and

$$V'_{N_1+1}(\varphi^\varepsilon) = O(\varepsilon^{N_1+1}) \quad (\varepsilon \searrow 0).$$

Consequently the set $\{\|\varphi^\varepsilon\|; \varepsilon \in]0, 1]\}$ is bounded and by (2), as $V'_{N_1} \leq V'_{N_1+1}$, we have

$$\|S_\varepsilon \varphi^\varepsilon\|^{2N_1/d} V'_{N_1}(S_\varepsilon \varphi^\varepsilon) \leq \|\varphi^\varepsilon\|^{2N_1/d} \varepsilon^{-N_1} V'_{N_1+1}(\varphi^\varepsilon) = O(\varepsilon).$$

It follows that for a sufficiently small ε , $S_\varepsilon \varphi^\varepsilon$ belongs to the open set (independent of x) $\{\varphi; \|\varphi\|^{2N_1/d} V'_{N_1}(\varphi) < 1\}$ where $R'(\bullet, x) = R(\bullet, x)$. Hence the assertions $R' \in \mathcal{E}_M^2$ and $R - R' \in \mathcal{N}$ are proved.

Now we want to prove that R' fulfills (3°). This means that the relations (3) with R' hold for all $\varepsilon \in]0, 1]$, provided $L = \ell = 1$ ($|\alpha| \leq 1$). To this aim, for $\psi \in \mathcal{A}(B)$ we first estimate

$$\begin{aligned} d_\psi \partial^\alpha R'(S_\varepsilon \varphi, x) &= d_\psi \partial^\alpha \left(\vartheta(\|S_\varepsilon \varphi\|^{2N_1/d} V'_{N_1}(S_\varepsilon \varphi)) \cdot R(S_\varepsilon \varphi, x) \right) \\ &= d_\psi \vartheta \left(\|S_\varepsilon \varphi\|^{2N_1/d} V'_{N_1}(S_\varepsilon \varphi) \right) \cdot \partial^\alpha R(S_\varepsilon \varphi, x) \\ (8) \quad &+ \vartheta \left(\|S_\varepsilon \varphi\|^{2N_1/d} V'_{N_1}(S_\varepsilon \varphi) \right) \cdot d_\psi \partial^\alpha R(S_\varepsilon \varphi, x) \\ &= d_\psi \vartheta \left(\|\varphi\|^{2N_1/d} \varepsilon^{-N_1} V'_{N_1}(\varphi) \right) \cdot \partial^\alpha R(S_\varepsilon \varphi, x) \\ &+ \vartheta \left(\|\varphi\|^{2N_1/d} \varepsilon^{-N_1} V'_{N_1}(\varphi) \right) \cdot d_\psi \partial^\alpha R(S_\varepsilon \varphi, x). \end{aligned}$$

If $\|\varphi\|^{2N_1/d} \varepsilon^{-N_1} V'_{N_1}(\varphi) > 2$ then $R'(S_\varepsilon \varphi, x) = 0$, hence we have to estimate (8) only if

$$\frac{1}{2}\|\varphi\|^{2N_1/d} V'_{N_1}(\varphi) \leq \varepsilon^{N_1}.$$

By the Hölder inequality we have (χ denotes the characteristic function):

$$(9) \quad 1 = \int \varphi \chi_B \leq \|\varphi\| \|\chi_B\|, \quad \text{i.e.} \quad \|\varphi\| \geq \|\chi_B\|^{-1}.$$

Consequently

$$\frac{1}{2} \|\chi_B\|^{-2N_1/d} V'_{N_1}(\varphi) \leq \varepsilon^{N_1}$$

and this is exactly our hypothesis (7) assuring that (3) holds. Hence two terms of (8) are estimated:

$$|\partial^\alpha R(S_\varepsilon \varphi, x)| \leq C\varepsilon^{-N_1}, \quad |\partial^\alpha d_\psi R(S_\varepsilon \varphi, x)| \leq C\varepsilon^{-N_1}.$$

It remains to estimate

$$(10) \quad d_\psi \vartheta \left(\|\varphi\|^{2N_1/d} \varepsilon^{-N_1} V'_{N_1}(\varphi) \right) \leq \max_t \left| \frac{d}{dt} \vartheta(t) \right| \varepsilon^{-N_1} \cdot d_\psi \left(\|\varphi\|^{2N_1/d} V'_{N_1}(\varphi) \right).$$

By [9, §2, Proposition] about local equicontinuity of differentials there is an absolutely convex open neighbourhood of zero $\mathcal{U} \subset \mathcal{A}(B)$ such that

$$(11) \quad d_\psi \left(\|\varphi\|^{2N_1/d} V'_{N_1}(\varphi) \right) \leq 1$$

whenever $\varphi \in \mathcal{B} + 2\mathcal{U}$, $\psi \in \mathcal{U}$. Under these hypotheses, we have got

$$d_\psi \vartheta \left(\|\varphi\|^{2N_1/d} \varepsilon^{-N_1} V'_{N_1}(\varphi) \right) \leq C_1 \varepsilon^{-N_1}$$

with a constant C_1 depending only on ϑ . Due to (8) and (3), it follows

$$d_\psi \partial^\alpha R'(S_\varepsilon \varphi, x) \leq C_1 C \varepsilon^{-2N_1} + C \varepsilon^{-N_1} \leq (C_1 + 1) C \varepsilon^{-2N_1}.$$

Replacing \mathcal{U} with a smaller one, we get $\leq \varepsilon^{-2N_1}$. It remains to estimate $\partial^\alpha R'(S_\varepsilon \varphi, x)$. This is similar or simpler, so we let it to the reader. □

PROOF OF (4°): Using the mean value theorem, we have for some $\tau \in]0, 1[$

$$\begin{aligned} & \left| \partial^\alpha d_{\psi_1, \dots, \psi_{\ell-1}}^{\ell-1} R(S_{2^{-n}} \varphi_2, x) - \partial^\alpha d_{\psi_1, \dots, \psi_{\ell-1}}^{\ell-1} R(S_{2^{-n}} \varphi_1, x) \right| \\ & \leq \left| \partial^\alpha d^\ell R(S_{2^{-n}}(\tau \varphi_1 + (1-\tau)\varphi_2), x) [\psi_1, \dots, \psi_{\ell-1}, \varphi_2 - \varphi_1] \right|. \end{aligned}$$

The function $\tau \mapsto V_N(\tau \varphi_1 + (1-\tau)\varphi_2)$ is convex because $V_N(\varphi)$ is the Euclidean norm of the point with coordinates $\int \xi^\beta \varphi(\xi) d\xi$. Thus in the second case of (4°), $V_{N_L}(\tau \varphi_1 + (1-\tau)\varphi_2) \leq \varepsilon^{N_L}$ holds for all $\tau \in [0, 1]$ and we can apply (4). □

§4. Notation. Let a number $r \in \mathbb{N}$ be given, let us consider the Fourier series of a function $\psi \in \mathcal{A}(K_r)$ on the cube $K_{r+1} \subset \mathbb{R}^d$ (§2, Notation, the function $\vartheta_r^{\otimes d}$ will be denoted simply by ϑ_r)

$$(12) \quad \begin{aligned} \psi(\xi) &= \sum_{\beta \in \mathbb{Z}^d} c'_\beta e^{2^{-r-1}\pi i \beta \cdot \xi} \quad \text{where } \beta \cdot \xi := \beta_1 \xi_1 + \dots + \beta_d \xi_d, \\ c'_\beta &= 2^{-d(r+2)} \int \psi(\xi) e^{-2^{-r-1}\pi i \beta \cdot \xi} d\xi. \end{aligned}$$

As $\int \psi = 0$, we have $c'_0 = 0$. As $\vartheta_r = 1$ on K_r , we have as well

$$\psi(\xi) = \sum_{\beta \neq 0} c'_\beta e^{2^{-r-1}\pi i \beta \cdot \xi} \vartheta_r(\xi).$$

We will use another expansion $\psi = \sum_{\beta \neq 0} c'_\beta \gamma'_\beta$ where the functions γ'_β are defined:

$$\gamma'_\beta(\xi) = e^{2^{-r-1}\pi i \beta \cdot \xi} \vartheta_r(\xi) - c''_\beta \vartheta_r(\xi)$$

with constants c''_β such that $\gamma'_\beta \in \mathcal{A}$. This means that

$$(13) \quad c''_\beta \int \vartheta_r(\xi) d\xi = \int e^{2^{-r-1}\pi i \beta \cdot \xi} \vartheta_r(\xi) d\xi.$$

It is known that the Fourier coefficients (12) of a test function tend rapidly to zero if $|\beta| \rightarrow \infty$. By (13), c''_β tend rapidly to zero as well.

We arrange the multi-indices $\beta \neq 0$ into a sequence $\{\beta_j\}_{j=1}^\infty$ in such a way that the sequence $\{|\beta_j|\}$ is non-decreasing; then we change the notation writing γ'_j, c'_j, \dots rather than $\gamma'_\beta, c'_\beta, \dots$. Then the above expansion takes the form

$$\psi = \sum_{j=1}^\infty c'_j \gamma'_j.$$

Evidently $|\beta_j| \leq j \leq (2|\beta_j|+1)^d$ ($\Leftarrow j$ does not exceed the number of the indices β with $|\beta| \leq |\beta_j|$). Hence any multi-sequence $\{a_\beta\}_\beta$ is moderated (i.e. $|a_\beta| \leq c|\beta|^m$ for some c and m) iff the sequence $\{a_{\beta_j}\}_j$ is moderated. $\{a_\beta\}_\beta$ tends rapidly to zero iff $\{a_{\beta_j}\}_j$ tends rapidly to zero.

If $\mathcal{U} \subset \mathcal{A}(K_{r+1})$ is an absolutely convex open neighbourhood of zero, then $\|\gamma'_j\|_{\mathcal{U}}$ is a moderate sequence (this can be calculated e.g. if $\|\gamma'_\beta\|_{\mathcal{U}}$ is the norm $\|\gamma'_\beta\|$ from §2, Remark). So we get the following

Result. If $\mathcal{U} \subset \mathcal{A}(K_{r+1})$ is an absolutely convex open neighbourhood of zero, then there are $\gamma_j \in \mathcal{A}(K_{r+1})$ ($j \in \mathbb{N}$) such that

$$(14) \quad \sum_{j=1}^{\infty} \|\gamma_j\|_{\mathcal{U}} \leq 1$$

and any function $\psi \in \mathcal{A}(K_r)$ has an expansion

$$\psi = \sum c_j \gamma_j$$

with coefficients c_j tending rapidly to zero.

Indeed, choose a moderate sequence $\lambda_j \nearrow \infty$ such that the functions $\gamma_j := \gamma'_j/\lambda_j \in \mathcal{A}(K_{r+1})$ fulfill (14) and then put $c_j = c'_j \cdot \lambda_j$.

Definition of $R_{krn\omega}$. Let to any $k, r, n \in \mathbb{N}$ and $\omega \in \mathcal{A}_0(K_r)$, a neighbourhood of zero \mathcal{U} in the space $\mathcal{A}(K_{r+1})$ be assigned which is the unit ball for a smooth norm (see §2, Remark), independent of n , following Properties of R (§3) with $\mathcal{B} = \{\omega\} \subset \mathcal{A}(K_{r+1})$ and all $L \leq k$. Assume furthermore that \mathcal{U} is as small as $|\mathrm{d}_\psi V_{N_L}(\varphi)| \leq 1$ whenever $L = 1, \dots, k$, $\varphi \in \omega + 2\mathcal{U}$, $\psi \in \mathcal{U}$, due to the local equicontinuity of the differentials of \mathcal{C}^∞ functions, [9, §2, Proposition].

Then the function $\varphi, x \mapsto R_{krn\omega}(\varphi, x)$ is defined on the domain

$$(15) \quad S_{2-n}(\omega + (\mathcal{U} \cap \mathcal{A}(K_r))) \times \mathbb{R}^d = S_{2-n}(\omega + \mathcal{U}) \cap \mathcal{A}_0(K_{r-n}) \times \mathbb{R}^d$$

as follows.

$$R_{krn\omega} := \lim_{\substack{J \in \mathbb{N} \\ J \rightarrow \infty}} R_J,$$

$$(16) \quad R_J(S_{2-n} \varphi, x) := \int \cdots \int R\left(S_{2-n}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \prod_{j=1}^J \rho_\delta(t_j) dt_j,$$

$$\varphi \in (\omega + \mathcal{U}) \cap \mathcal{A}_0(K_r), \quad \rho_\delta := S_\delta \rho \quad (\S 2, \text{following (2)}), \quad \delta = \delta_{kn} := 2^{-n(k+1)N_k}.$$

For the sake of simplicity of the notation, we do not indicate the dependence of R_J on k, r, n, ω .

Properties of $R_{krn\omega}$.

(1°) If $k, r, n \in \mathbb{N}$, $\omega \in \mathcal{A}_0(K_r)$, then $R_{krn\omega}$ is well defined on its domain (15). If $x \in \mathbb{R}^d$, $\varphi \in \omega + (\mathcal{U} \cap \mathcal{A}(K_r))$, then

$$(17) \quad R_{krn\omega}(S_{2-n} \varphi, x) = \lim_{J \rightarrow \infty} R_J(S_{2-n} \varphi, x)$$

uniformly with respect to φ, x and

$$(18) \quad |R_{krn\omega}(S_{2^{-n}}\varphi, x)| \leq C \cdot 2^{nN_1},$$

with a constant C not depending on n .

(2°) If $n \in \mathbb{N}$, $x \in \mathbb{R}^d$, $\ell = 0, 1, \dots, L - 1$, $L \leq k$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L$, $\varphi \in \omega + (\mathcal{U} \cap \mathcal{A}(K_r))$, $2^{-nN_L-1} > V_{N_L}(\varphi)$, $\psi_1, \dots, \psi_\ell \in \mathcal{U} \cap \mathcal{A}(K_r)$, then

$$(19) \quad \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{krn\omega}(S_{2^{-n}}\varphi, x) = \lim_{J \rightarrow \infty} \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_J(S_{2^{-n}}\varphi, x)$$

uniformly with respect to $x, \varphi, \psi_1, \dots, \psi_\ell$ under the above conditions (k, r, n fixed). Consequently $\varphi, x \mapsto \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{krn\omega}(S_{2^{-n}}\varphi, x)$ is continuous, hence the order of derivatives (under the above conditions) does not matter. Furthermore we have

$$(20) \quad \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{krn\omega}(S_{2^{-n}}\varphi, x) \right| \leq C 2^{nN_L}$$

with a constant C not depending on n , and

$$(21) \quad \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (R_{krn\omega}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x)) \right| \leq 2^{nN_L} \cdot \delta_{kn}.$$

(3°) $R_{krn\omega}$ is \mathcal{C}^∞ with respect to the first variable on its domain (15) and there is an absolutely convex open neighbourhood of zero $\mathcal{V} = \mathcal{V}_{kr\omega} \subset \mathcal{A}(K_r)$ not depending on n such that if $x \in \mathbb{R}^d$, $n \in \mathbb{N}$, $L \in \mathbb{N}$, $\psi_\ell \in \mathcal{V}$ ($\ell = 1, \dots, L$), $\varphi \in \omega + \mathcal{U}$, then

$$(22) \quad |d_{\psi_1, \dots, \psi_L}^L R_{krn\omega}(S_{2^{-n}}\varphi, x)| \leq b_L \cdot 2^{nN_1} \cdot \delta_{kn}^{-L}$$

with a constant b_L depending only on L and ρ .

PROOF OF (17) AND (19): By the definition (16) we have

$$\begin{aligned} & \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_J(S_{2^{-n}}\varphi, x) \\ &= \int \dots \int \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \prod_{j=1}^J \rho_\delta(t_j) dt_j \end{aligned}$$

and we have as well

$$\begin{aligned} & \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_J(S_{2^{-n}}\varphi, x) \\ &= \int \dots \int \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \prod_{j=1}^{J+1} \rho_\delta(t_j) dt_j, \end{aligned}$$

because $\int \rho_\delta(t_{J+1}) dt_{J+1} = 1$. It follows

$$\begin{aligned}
 & \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{J+1}(S_{2^{-n}}\varphi, x) - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_J(S_{2^{-n}}\varphi, x) \right| \\
 (23) \quad &= \left| \int_{-\delta}^\delta \cdots \int_{-\delta}^\delta \left[\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^{J+1} t_j \gamma_j\right), x\right) \right. \right. \\
 & \quad \left. \left. - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \right] \cdot \prod_{j=1}^{J+1} \rho_\delta(t_j) dt_j \right|.
 \end{aligned}$$

Now we want to apply §3, Property of R (4°) for $\varepsilon = 2^{-n}$. By hypotheses of §3, (4°), there are two cases. For $L = 1$ this gives estimation

$$\begin{aligned}
 & |R_{J+1}(S_{2^{-n}}\varphi, x) - R_J(S_{2^{-n}}\varphi, x)| \\
 (24) \quad & \leq \left| \int_{-\delta}^\delta \cdots \int_{-\delta}^\delta 2^{nN_1} \|t_{j+1}\gamma_{j+1}\|_{\mathcal{U}} \cdot \prod_{j=1}^{J+1} \rho_\delta(t_j) dt_j \right| \\
 & \leq 2^{nN_1} \delta_{kn} \|\gamma_{J+1}\|_{\mathcal{U}} \leq \|\gamma_{J+1}\|_{\mathcal{U}}
 \end{aligned}$$

(δ defined in (16)), so by (14) the limit in (17) is uniform. Thus $R_{k\ell n\omega}$ is well defined. Similarly (19) can be deduced from (23): By the local equicontinuity of $dV_{N_L}(\varphi)$ noted in the definition of $R_{k\ell n\omega}$, we get using the mean value theorem

$$\left| V_{N_L}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right) - V_{N_L}(\varphi) \right| \leq \left\| \sum_{j=1}^J t_j \gamma_j \right\|_{\mathcal{U}} \leq \delta = 2^{-n(k+1)N_k} \leq 2^{-nN_L-1}.$$

From the hypothesis in (2°) $V_{N_L}(\varphi) < 2^{-nN_L-1}$, we obtain $\left| V_{N_L}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right) \right| \leq 2^{-nN_L}$ that is the hypothesis in §3, Property (4°). Thus, by §3, (4°) (for $\ell = 0, 1, \dots, L - 1$) we get from (23):

$$\begin{aligned}
 & \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{J+1}(S_{2^{-n}}\varphi, x) - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_J(S_{2^{-n}}\varphi, x) \right| \\
 (25) \quad & \leq 2^{nN_L} \|\psi_1\|_{\mathcal{U}} \cdots \|\psi_\ell\|_{\mathcal{U}} \cdot \delta_{kn} \|\gamma_{J+1}\|_{\mathcal{U}}.
 \end{aligned}$$

As above, thanks to (14), the uniform convergence of the limit in (19) and then the equality (19) follows. □

PROOF OF (18) AND (20): In all cases where the uniform convergence is already proved, we have

$$\begin{aligned}
 & \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{k\ell n\omega}(S_{2^{-n}}\varphi, x) = \lim_{J \rightarrow \infty} \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_J(S_{2^{-n}}\varphi, x) \\
 & = \lim_{J \rightarrow \infty} \int \cdots \int \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \prod_{j=1}^J \rho_\delta(t_j) dt_j.
 \end{aligned}$$

It was shown while proving (19) that the hypothesis in (2°) $V_{N_L}(\varphi) < 2^{-nN_L-1}$ implies $V_{N_L}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right) \leq 2^{-nN_L}$, and this is the hypothesis in Properties of R (§3) allowing us to use (3) and (4) for estimating the term $\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right)$ in the last integral. By §3, Property (3°) this hypothesis is not needed for proving (18). Thus (18) and (20) follow from the corresponding properties of R . □

PROOF OF (21): (25) holds for $J = 0$ as well with $R_J = R$. Adding the inequalities (25), we get

$$\begin{aligned} & \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R_{J+1}(S_{2^{-n}}\varphi, x) - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R(S_{2^{-n}}\varphi, x) \right| \\ & \leq 2^{nN_L} \|\psi_1\|_{\mathcal{U}} \dots \|\psi_\ell\|_{\mathcal{U}} \cdot \delta_{kn} \sum_{j=1}^{J+1} \|\gamma_j\|_{\mathcal{U}} \leq 2^{nN_L} \|\psi_1\|_{\mathcal{U}} \dots \|\psi_\ell\|_{\mathcal{U}} \cdot \delta_{kn} \end{aligned}$$

due to (14). Hence the inequality (21) is proved. □

PROOF OF 3°: For $L \in \mathbb{N}$ let $\psi_1, \dots, \psi_L \in \mathcal{A}(K_r)$ be given functions, let

$$(26) \quad \psi_\ell = \sum_{j=1}^\infty c_{\ell j} \gamma_j, \quad \text{i.e.} \quad S_{2^{-n}}\psi_\ell = \sum_{j=1}^\infty c_{\ell j} S_{2^{-n}}\gamma_j \quad (\ell = 1, \dots, L)$$

be their expansions by §4, Notation with γ_j fulfilling (14). As $\lim_{j \rightarrow \infty} c_{\ell j} = 0$ (rapidly), there is an $A > 0$ for which

$$(27) \quad |c_{\ell j}| \leq A \quad (\forall \ell = 1, \dots, L, j \in \mathbb{N}).$$

In the following calculation, h_1, \dots, h_L are real variables with

$$(28) \quad |h_\ell| < \frac{1 - \delta}{LA}$$

and we have to put $h_1, \dots, h_L = 0$ to obtain the following equality:

$$\begin{aligned} d_{\psi_1, \dots, \psi_L}^L R_{krn\omega}(S_{2^{-n}}\varphi, x) &= \frac{\partial^L}{\partial h_1 \dots \partial h_L} \lim_{J \rightarrow \infty} R_J\left(S_{2^{-n}}\left(\varphi + \sum_{\ell=1}^L h_\ell \psi_\ell\right), x\right) \\ &= \frac{\partial^L}{\partial h_1 \dots \partial h_L} \lim_{J \rightarrow \infty} \int \dots \int R\left(S_{2^{-n}}\left(\varphi + \sum_{\ell=1}^L h_\ell \psi_\ell + \sum_{j=1}^J t_j \gamma_j\right), x\right) \prod_{j=1}^J \rho_\delta(t_j) dt_j. \end{aligned}$$

By (26) this is equal to

$$\begin{aligned}
 & \frac{\partial^L}{\partial h_1 \dots \partial h_L} \lim_{J \rightarrow \infty} \int \dots \int R\left(S_{2^{-n}}\left(\varphi + \sum_{\ell=1}^L h_\ell \sum_{j=1}^\infty c_{\ell j} \gamma_j + \sum_{j=1}^J t_j \gamma_j\right), x\right) \\
 & \qquad \qquad \qquad \cdot \prod_{j=1}^J \rho_\delta(t_j) dt_j \\
 (29) \quad & = \frac{\partial^L}{\partial h_1 \dots \partial h_L} \lim_{J \rightarrow \infty} \int \dots \int R\left(S_{2^{-n}}\left(\varphi + \sum_{\ell=1}^L h_\ell \sum_{j=1}^J c_{\ell j} \gamma_j + \sum_{j=1}^J t_j \gamma_j\right), x\right) \\
 & \qquad \qquad \qquad \cdot \prod_{j=1}^J \rho_\delta(t_j) dt_j
 \end{aligned}$$

because by §3, Property (4°), (27) and (28), the difference of both expressions after $\lim_{J \rightarrow \infty}$ is estimated by

$$2^{nN_1} \sum_{\ell=1}^L |h_\ell| \sum_{j=J+1}^\infty |c_{\ell j}| \cdot \|\gamma_j\|_{\mathcal{U}} \leq 2^{nN_1} (1 - \delta) \sum_{j=J+1}^\infty \|\gamma_j\|_{\mathcal{U}}.$$

This tends to zero thanks to (14), only we have to verify the hypothesis in §3, (4°) that $\varphi + \sum_{\ell=1}^L h_\ell \sum_{j=1}^\infty c_{\ell j} \gamma_j + \sum_{j=1}^J t_j \gamma_j$ and $\varphi + \sum_{\ell=1}^L h_\ell \sum_{j=1}^J c_{\ell j} \gamma_j + \sum_{j=1}^J t_j \gamma_j$ are elements of $\omega + 2\mathcal{U}$. Indeed, $\varphi \in \omega + \mathcal{U}$ and for the other terms we have by (28), (27) and (14)

$$\left\| \sum_{\ell=1}^L h_\ell \sum_{j=1}^\infty c_{\ell j} \gamma_j + \sum_{j=1}^J t_j \gamma_j \right\|_{\mathcal{U}} < \sum_{\ell=1}^L \frac{1 - \delta}{LA} \sum_{j=1}^\infty A \|\gamma_j\|_{\mathcal{U}} + \sum_{j=1}^J \delta \|\gamma_j\|_{\mathcal{U}} \leq 1.$$

Thus (29) is verified. After a substitution in (29) (and putting $h_1, \dots, h_L = 0$) we get

$$\begin{aligned}
 (30) \quad & d_{\psi_1, \dots, \psi_L}^L R_{krn\omega}(S_{2^{-n}} \varphi, x) = \frac{\partial^L}{\partial h_1 \dots \partial h_L} \\
 & \lim_{J \rightarrow \infty} \int \dots \int R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \cdot \prod_{j=1}^J \rho_\delta\left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j}\right) dt_j \\
 & = \lim_{J \rightarrow \infty} \int \dots \int R\left(S_{2^{-n}}\left(\varphi + \sum_{j=1}^J t_j \gamma_j\right), x\right) \\
 & \qquad \qquad \qquad \cdot \frac{\partial^L}{\partial h_1 \dots \partial h_L} \prod_{j=1}^J \rho_\delta\left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j}\right) dt_j
 \end{aligned}$$

provided the last limit is uniform with respect to h_ℓ , $|h_\ell| < \frac{1-\delta}{LA}$ ($\ell = 1, \dots, L$).

Now we are going to prove it. Let us denote the last integral by I_J . Using the Leibniz rule for the derivation of a product:

$$\frac{\partial}{\partial h_\ell} \prod_{j=1}^J \rho_\delta \left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j} \right) = \sum_{j_\ell=1}^J (-c_{\ell, j_\ell}) \frac{\partial}{\partial t_{j_\ell}} \prod_{j=1}^J \rho_\delta \left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j} \right),$$

we obtain

$$I_J = \int \cdots \int R \left(S_{2-n} \left(\varphi + \sum_{j=1}^J t_j \gamma_j \right), x \right) \cdot \sum_{j_1, \dots, j_L=1}^J \left(\prod_{\ell=1}^L -c_{\ell, j_\ell} \frac{\partial}{\partial t_{j_\ell}} \right) \prod_{j=1}^J \rho_\delta \left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j} \right) dt_j.$$

Using the Kronecker delta ($\delta_j^{j'}$ is the truth value of the statement $j = j'$) we can write

$$I_J = \int \cdots \int R \left(S_{2-n} \left(\varphi + \sum_{j=1}^J t_j \gamma_j \right), x \right) \cdot \sum_{j_1, \dots, j_L=1}^J \left(\prod_{\ell=1}^L -c_{\ell, j_\ell} \right) \prod_{j=1}^J \left(\frac{\partial}{\partial t_j} \right)^{\sum_{\ell=1}^L \delta_j^{j_\ell}} \rho_\delta \left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j} \right) dt_j$$

and we have as well

$$I_J = \int \cdots \int R \left(S_{2-n} \left(\varphi + \sum_{j=1}^J t_j \gamma_j \right), x \right) \cdot \sum_{j_1, \dots, j_L=1}^{J+1} \left(\prod_{\ell=1}^L -c_{\ell, j_\ell} \right) \prod_{j=1}^{J+1} \left(\frac{\partial}{\partial t_j} \right)^{\sum_{\ell=1}^L \delta_j^{j_\ell}} \rho_\delta \left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j} \right) dt_j.$$

Indeed, if some $j_\ell = J+1$, then the term $\rho_\delta \left(t_{J+1} - \sum_{\ell=1}^L h_\ell c_{\ell, J+1} \right)$ is differentiated and so its integral is equal to 0; else its integral is is equal to 1. It follows

$$I_{J+1} - I_J = \int \cdots \int \left[R \left(S_{2-n} \left(\varphi + \sum_{j=1}^{J+1} t_j \gamma_j \right), x \right) - R \left(S_{2-n} \left(\varphi + \sum_{j=1}^J t_j \gamma_j \right), x \right) \right] \cdot \sum_{j_1, \dots, j_L=1}^{J+1} \left(\prod_{\ell=1}^L -c_{\ell, j_\ell} \right) \prod_{j=1}^{J+1} \left(\frac{\partial}{\partial t_j} \right)^{\sum_{\ell=1}^L \delta_j^{j_\ell}} \rho_\delta \left(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j} \right) dt_j.$$

By (27) and (28) we have $|\sum_{\ell=1}^L h_\ell c_{\ell j}| \leq 1 - \delta$, so $|t_j| \leq 1$ or $\rho_\delta(t_j - \sum_{\ell=1}^L h_\ell c_{\ell j}) = 0$, and we can apply §3, Property of R (4°). We get

$$|I_{J+1} - I_J| \leq 2^{nN_1} \|\gamma_{J+1}\|_{\mathcal{U}} \sum_{j_1, \dots, j_L=1}^{J+1} \prod_{\ell=1}^L |c_{\ell, j_\ell}| \prod_{j=1}^{J+1} \|\partial^{\sum_{\ell=1}^L \delta_j^{j_\ell}} \rho_\delta\|_{\mathcal{F}^1}.$$

It is $\|\rho_\delta\|_{\mathcal{F}^1} = 1$ (§2, following (2)) and, in the last product, for given j_1, \dots, j_L there are at most L indices j for which ρ_δ is differentiated. Thus this product can be estimated

$$\begin{aligned} \prod_{j=1}^{J+1} \|\partial^{\sum_{\ell=1}^L \delta_j^{j_\ell}} \rho_\delta\|_{\mathcal{F}^1} &= \prod_{j=1}^{J+1} \delta^{-\sum_{\ell=1}^L \delta_j^{j_\ell}} \|\partial^{\sum_{\ell=1}^L \delta_j^{j_\ell}} \rho\|_{\mathcal{F}^1} \\ &\leq \delta^{-\sum_{j=1}^{J+1} \sum_{\ell=1}^L \delta_j^{j_\ell}} \left(\max_{0 \leq \ell \leq L} \|\partial^\ell \rho\|_{\mathcal{F}^1} \right)^L = \delta^{-L} \cdot b_L \end{aligned}$$

with a constant b_L depending only on ρ and L . It follows

$$\begin{aligned} |I_{J+1} - I_J| &\leq 2^{nN_1} \delta^{-L} b_L \|\gamma_{J+1}\|_{\mathcal{U}} \sum_{j_1, \dots, j_L=1}^{J+1} \prod_{\ell=1}^L |c_{\ell, j_\ell}| \\ (31) \quad &= 2^{nN_1} \delta^{-L} b_L \|\gamma_{J+1}\|_{\mathcal{U}} \prod_{\ell=1}^L \sum_{j=1}^{J+1} |c_{\ell j}| \leq 2^{nN_1} \delta^{-L} b_L \|\gamma_{J+1}\|_{\mathcal{U}} \prod_{\ell=1}^L \sum_{j=1}^{\infty} |c_{\ell j}| \\ &= 2^{nN_1} c^L \delta^{-L} b_L \|\gamma_{J+1}\|_{\mathcal{U}} \end{aligned}$$

where the constant $c = \max_{\ell=1, \dots, L} \sum_{j=1}^{\infty} |c_{\ell j}|$ depends only on ψ_1, \dots, ψ_L and r .

Now the uniform convergence of (30) can be deduced from (14). The smoothness is verified; it remains to deduce the estimations.

The inequality (31) holds for $J = 0$ as well with $I_0 = 0$. Adding these inequalities we get

$$|I_J| \leq 2^{nN_1} c^L \delta^{-L} b_L \sum_{j=1}^J \|\gamma_j\|_{\mathcal{U}} \leq 2^{nN_1} c^L \delta^{-L} b_L.$$

This is an estimation for the last integral in (30). Hence

$$|d_{\psi_1, \dots, \psi_L}^L R_{k r n \omega}(S_{2^{-n}} \varphi, x)| \leq b_L c^L 2^{nN_1} \cdot \delta^{-L}.$$

b_L depends only on ρ and L , c is the constant in (31), $c = 1$ if ψ_1, \dots, ψ_L belong to

$$\mathcal{V} = \left\{ \psi \in \mathcal{A}(K_r); \psi = \sum c_j \gamma_j, \sum |c_j| \leq 1 \right\}.$$

By §4, Result γ_j depends on \mathcal{U} not depending on n . It remains to prove that \mathcal{V} is a neighbourhood of zero in $\mathcal{A}(K_r)$. It is known that the Fourier coefficients of functions ψ running over a bounded set in $\mathcal{A}(K_r)$ tend uniformly rapidly to zero. Evidently the same holds for the coefficients c_j in §4, Notation. Hence any bounded set is absorbed by \mathcal{V} . In a metric vector space such sets \mathcal{V} are neighbourhoods of zero. □

§5. Partition of unity. The space \mathcal{D} has the property of smooth partition of unity expressed by the following

Theorem. For $B \in \mathbb{R}^d$, let $\{\omega_s + \mathcal{U}_s\}_{s \in S}$ be an open covering of the space $\mathcal{D}(B)$, where S is an arbitrary set of indices, $\omega_s \in \mathcal{D}(B)$ ($\forall s \in S$), \mathcal{U}_s are open neighbourhoods of zero in $\mathcal{D}(B)$. Then there is a locally finite smooth (i.e. \mathcal{C}^∞) partition of unity on $\mathcal{D}(B)$

$$1 = \sum_{m=1}^{\infty} \Phi_m$$

subordinated to this covering.

This means:

1° The functions $\Phi_m : \mathcal{D}(B) \rightarrow [0, 1]$, fulfilling this equality, are \mathcal{C}^∞ and

$$\forall m \quad \exists s \in S : \text{supp } \Phi_m \subset \omega_s + \mathcal{U}_s.$$

2° For every $\omega \in \mathcal{D}(B)$ there is an absolutely convex open neighbourhood of zero $\mathcal{U} \subset \mathcal{D}(B)$ such that $\omega + \mathcal{U}$ meets only a finite number of supports of functions Φ_m .

For the proof, we refer to [13, (5.3.8)], where a more general theorem is proved concerning several categories of smoothness, not only \mathcal{C}^∞ . Hypotheses: \mathcal{D} is a Lindelöf locally convex space and there are sufficiently many \mathcal{C}^∞ functions on \mathcal{D} so that they generate the original topology on \mathcal{D} . This is fulfilled, see §2, Remark.

Corollary. For $B \in \mathbb{R}^d$, let $\{\omega_s + \mathcal{U}_s\}_{s \in S}$ be an open covering of the space $\mathcal{A}_0(B)$, where ($\forall s \in S$) $\omega_s \in \mathcal{A}_0(B)$ and \mathcal{U}_s is an open neighbourhood of zero in $\mathcal{A}(B)$. Then there is a locally finite smooth partition of unity on $\mathcal{A}_0(B)$

$$1 = \sum_{m=1}^{\infty} \Phi_m$$

subordinated to this covering.

PROOF: We can write $\mathcal{U}_s = \tilde{\mathcal{U}}_s \cap \mathcal{A}(B)$ where $\tilde{\mathcal{U}}_s$ are neighbourhoods of zero in the space $\mathcal{D}(B)$. Then we apply Theorem to the covering $\{(\omega_s + \tilde{\mathcal{U}}_s)\}_s \cup \{\mathcal{D}(B) \setminus \mathcal{A}_0(B)\}$ of $\mathcal{D}(B)$. \square

§6. Notation. Let us have chosen $k, r \in \mathbb{N}$. Then, for every $\omega \in \mathcal{A}_0(K_r)$, we have a neighbourhood of zero $\mathcal{U}_\omega \subset \mathcal{A}(K_r)$ (independent of n) such that $\forall n \in \mathbb{N}$ the function $R_{krn\omega}$ is defined by §4, on $S_{2^{-n}}(\omega + \mathcal{U}_\omega) \times \mathbb{R}^d$. Thus we have a covering of $\mathcal{A}_0(K_r)$ with the sets $\omega + \mathcal{U}_\omega$. We choose a partition of unity $1 = \sum_{m=1}^\infty \Phi_m$ on $\mathcal{A}_0(K_r)$ by the above corollary. For every m , we choose a test function ω_m for which $\text{supp } \Phi_m \subset \omega_m + \mathcal{U}_{\omega_m}$; we will use the notation \mathcal{U}_m rather than \mathcal{U}_{ω_m} .

Definition of R_{krn} . With the above notation, for $\varphi \in \mathcal{A}_0(K_r)$ (so $S_{2^{-n}} \varphi \in \mathcal{A}_0(K_{r-n})$) we define

$$(32) \quad R_{krn}(S_{2^{-n}} \varphi, x) := \sum_{m=1}^\infty \Phi_m(\varphi) \cdot R_{krn\omega_m}(S_{2^{-n}} \varphi, x).$$

If φ does not belong to $\text{supp } \Phi_m$, the term of this sum is considered to be zero even if $R_{krn\omega_m}(S_{2^{-n}} \varphi, x)$ is not defined.

Properties of R_{krn} . (1°) For every $k, r, n \in \mathbb{N}$, the function R_{krn} is defined on $\mathcal{A}_0(K_{r-n}) \times \mathbb{R}^d$.

Moreover, for every $\omega \in \mathcal{A}_0(K_r)$ and $L \in \mathbb{N}_0$, there exist an absolutely convex open neighbourhood of zero $\mathcal{U} \subset \mathcal{A}(K_r)$ and a constant $C > 0$, both independent of n , such that for every $n \in \mathbb{N}$ and $x \in \mathbb{R}^d$ the following hold.

(2°) If $1 \leq L \leq k$, $\ell = 0, 1, \dots, L - 1$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L$, $\varphi \in \omega + \mathcal{U}$, $2^{-nN_L-1} > V_{N_L}(\varphi)$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$, then $\varphi, x \mapsto d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R_{krn}(S_{2^{-n}} \varphi, x)$ is continuous and

$$(33) \quad \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R_{krn}(S_{2^{-n}} \varphi, x) \right| \leq C \cdot 2^{nN_L},$$

$$(34) \quad \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (R_{krn}(S_{2^{-n}} \varphi, x) - R(S_{2^{-n}} \varphi, x)) \right| \leq C 2^{nN_L} \cdot \delta_{kn}$$

($\delta = \delta_{kn}$ by (16)).

(3°) R_{krn} is \mathcal{C}^∞ with respect to the first variable on its domain $\mathcal{A}_0(K_{r-n})$ and if $\varphi \in \omega + \mathcal{U}$ and $\psi_\ell \in \mathcal{U}$ ($\ell = 1, \dots, L$), then

$$\left| d_{\psi_1, \dots, \psi_L}^L R_{krn}(S_{2^{-n}} \varphi, x) \right| \leq C 2^{nN_1} \cdot \delta_{kn}^{-L}.$$

PROOF OF (2°): If $\omega \in \mathcal{A}_0(K_r)$, we choose a neighbourhood of zero $\mathcal{U} \subset \mathcal{A}(K_r)$ such that $\omega + \mathcal{U}$ meets only a finite number of supports of the functions Φ_m (by 5.2°). So \mathcal{U} can be chosen as small as ($\forall m$)

$$(35) \quad \text{either } (\omega + \mathcal{U}) \cap \text{supp } \Phi_m = \emptyset \quad \text{or } (\omega + \mathcal{U}) \subset \omega_m + \mathcal{U}_m.$$

Furthermore, thanks to [9, §2, Proposition], let \mathcal{U} be as small as

$$(36) \quad \left| d_{\psi_1, \dots, \psi_\ell}^\ell \Phi_m(\varphi) \right| \leq 1$$

whenever $1 \leq \ell \leq k - 1$, $m \in \mathbb{N}$, $\varphi \in \omega + \mathcal{U}$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$; for $\ell = 0$ this is fulfilled, too. Now we use the following Leibniz rule for derivation of a product of two functions: if F_1, F_2 are two smooth functions on (a part of) a locally convex space, then

$$d^\ell (F_1(\varphi)F_2(\varphi))[\psi_1, \dots, \psi_\ell] = \sum_{\substack{I_1 \\ I_1 \cup I_2 = \{1, \dots, \ell\} \\ \text{disjoint}}} d_{\psi_{I_1}}^{\#I_1} F_1(\varphi) \cdot d_{\psi_{I_2}}^{\#I_2} F_2(\varphi),$$

where, for $I = \{i_1, \dots, i_{\#I}\} \subset \{1, \dots, \ell\}$, ψ_I denotes the finite sequence $\psi_{i_1}, \dots, \psi_{i_{\#I}}$ and the summation is extended over all ordered decompositions of multi-index $(1, \dots, \ell)$ in two disjoint multi-indices I_1, I_2 that are written in the increasing order. Unlike the chain rule (theorem on the derivative of the composition), here the multi-index I_1 or I_2 can be empty. The proof of the Leibniz formula can be deduced easily from the chain rule if the outer function is $F(u, v) = uv$, $u = F_1$, $v = F_2$.

We apply the Leibniz rule for differentiating the product to the defining formula (32):

$$\begin{aligned} d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R_{krn}(S_{2-n}\varphi, x) \\ = \sum_{m=1}^\infty \sum_{\substack{I_1 \\ I_1 \cup I_2 = \{1, \dots, \ell\} \\ \text{disjoint}}} d_{\psi_{I_1}}^{\#I_1} \Phi_m(\varphi) \cdot d_{\psi_{I_2}}^{\#I_2} \partial^\alpha R_{krn\omega_m}(S_{2-n}\varphi, x) \end{aligned}$$

and, by (35) as $\varphi \in \omega + \mathcal{U}$, the first sum is extended only over a finite number of m for which $(\omega + \mathcal{U}) \subset \omega_m + \mathcal{U}_m$. Then $(\omega + \mathcal{U}) - (\omega + \mathcal{U}) \subset (\omega_m + \mathcal{U}_m) - (\omega_m + \mathcal{U}_m)$; for absolutely convex sets it follows $\mathcal{U} \subset \mathcal{U}_m$. Hence the functions $R_{krn\omega_m}$, fulfilling §4, Properties on $\omega_m + \mathcal{U}_m$ fulfil it the more on $\omega + \mathcal{U}$. By the above Leibniz formula, we estimate $d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R_{krn}(S_{2-n}\varphi, x)$ using (36) for estimating the term $d_{\psi_{I_1}}^{\#I_1} \Phi_m(\varphi)$ and using §4, Properties of $R_{krn\omega}$ for estimating the term $d_{\psi_{I_2}}^{\#I_2} \partial^\alpha R_{krn\omega_m}(S_{2-n}\varphi, x)$. So we deduce (33) from the corresponding inequality (20) in Properties of $R_{krn\omega}$. The constant C in (33) depends only on the constants assigned by §4, Properties to the functions $R_{krn\omega_m}$ and on the used finite set of terms of the sum \sum_m , so it is independent of n .

For estimating $\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (R_{krn}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x))$, we apply the Leibniz formula to the product $\sum_{m=1}^\infty \Phi_m(\varphi) \cdot (R_{krn\omega_m}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x))$ and proceed similarly. \square

PROOF OF (3°): Unlike in §4, Properties of $R_{krn\omega}$, here the neighbourhood \mathcal{U} depends on L . We chose \mathcal{U} as small as (36) hold whenever $1 \leq \ell \leq L$, $m \in \mathbb{N}$, $\varphi \in \omega + \mathcal{U}$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$. Denote by \mathcal{V}_m the neighbourhood \mathcal{V} defined by §4 (3°) for the function $R_{krn\omega_m}$ and chose furthermore \mathcal{U} such that we have instead of (35):

$$\text{either } (\omega + \mathcal{U}) \cap \text{supp } \Phi_m = \emptyset \quad \text{or} \quad (\omega + \mathcal{U}) \subset \omega_m + (\mathcal{V}_m \cap \mathcal{U}_m).$$

Then we follow the above proof and deduce (3°) from the corresponding properties of $R_{krn\omega}$, §4: (18) and the item (3°). \square

§7. Notations. Choose test functions $\psi_\alpha \in \mathcal{D}(K_r \setminus K_{r-1})$ ($\alpha \in \mathbb{N}_0^d$, $0 \leq |\alpha| \leq N_k$), fulfilling (like in [8, (22), (23)])

$$\begin{aligned} \int \psi_\alpha(\xi) \cdot \xi^\alpha \, d\xi &= 1 \\ \int \psi_\alpha(\xi) \cdot \xi^\beta \, d\xi &= 0 \quad \text{for } \beta \neq \alpha, 0 \leq |\beta| \leq N_k. \end{aligned}$$

Let $\Lambda_r : \mathcal{D}(K_{r+1}) \rightarrow \mathcal{D}(K_r)$ be a continuous (hence smooth) linear mapping defined (see end of §2, ϑ_r means $\vartheta_r^{\otimes d}$):

$$\Lambda_r \varphi := \varphi \cdot \vartheta_{r-1} + \sum_{0 \leq |\alpha| \leq N_k} c_\alpha \psi_\alpha$$

with such constants c_α depending on φ that

$$\forall \beta \in \mathbb{N}_0^d, 0 \leq |\beta| \leq N_k : \quad \int \Lambda_r \varphi(\xi) \xi^\beta \, d\xi = \int \varphi(\xi) \xi^\beta \, d\xi.$$

This means $\int \varphi(\xi) \xi^\beta \, d\xi = \int \vartheta_{r-1}(\xi) \varphi(\xi) \xi^\beta \, d\xi + c_\beta$, hence c_β are well determined; Λ_r maps $\mathcal{A}_0(K_{r+1})$ into $\mathcal{A}_0(K_r)$ and $\mathcal{A}(K_{r+1})$ into $\mathcal{A}(K_r)$. Λ_r is identical on $\mathcal{D}(K_{r-1})$; for $\varphi \in \mathcal{A}_0(K_{r+1})$ and $N \leq N_k$ we have $V_N(\Lambda_r \varphi) = V_N(\varphi)$.

Definition of R'_{krn} . Let $k, n \in \mathbb{N}$ be given. For $r \in \mathbb{N}$, we define the functions R'_{krn} on $\mathcal{A}_0(K_{r-n}) \times \mathbb{R}^d$ by induction as follows. $R'_{k1n} = R_{k1n}$. If R'_{krn} is already defined on $\mathcal{A}_0(K_{r-n}) \times \mathbb{R}^d$, we define

$$\begin{aligned} (37) \quad R'_{k,r+1,n}(S_{2^{-n}}\varphi, x) &:= \\ R_{k,r+1,n}(S_{2^{-n}}\varphi, x) + R'_{krn}(S_{2^{-n}}(\Lambda_r \varphi), x) - R_{k,r+1,n}(S_{2^{-n}}(\Lambda_r \varphi), x) \\ &\text{for } \varphi \in \mathcal{A}_0(K_{r+1}), x \in \mathbb{R}^d. \end{aligned}$$

Properties of R'_{krn} . (1°) For every $k, r, n \in \mathbb{N}$ the function R'_{krn} is defined on $\mathcal{A}_0(K_{r-n}) \times \mathbb{R}^d$ and for $\varphi \in \mathcal{A}_0(K_{r-n-1})$ it is $R'_{k,r,n}(\varphi, x) = R'_{k,r+1,n}(\varphi, x)$.

Moreover, for every $\omega' \in \mathcal{A}_0(K_r)$ and $L \in \mathbb{N}_0$ there is an absolutely convex open neighbourhood of zero $\mathcal{U}' \subset \mathcal{A}(K_r)$ and a constant $C'_{kr} > 0$, both independent of n , such that for every $n \in \mathbb{N}$, $x \in \mathbb{R}^d$ the following hold.

(2°) If $1 \leq L \leq k$, $\ell = 0, 1, \dots, L-1$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L$, $\varphi' \in \omega' + \mathcal{U}'$, $2^{-nN_L-1} > V_{N_L}(\varphi')$, $\psi'_1, \dots, \psi'_\ell \in \mathcal{U}'$, then $\varphi', x \mapsto d_{\psi'_1, \dots, \psi'_\ell}^\ell \partial^\alpha R'_{krn}(S_{2^{-n}}\varphi', x)$ is continuous and

$$(38) \quad \left| d_{\psi'_1, \dots, \psi'_\ell}^\ell \partial^\alpha R'_{krn}(S_{2^{-n}}\varphi', x) \right| \leq C'_{kr} 2^{nN_L},$$

$$(39) \quad \left| \partial^\alpha d_{\psi'_1, \dots, \psi'_\ell}^\ell (R'_{krn}(S_{2^{-n}}\varphi', x) - R(S_{2^{-n}}\varphi', x)) \right| \leq C'_{kr} 2^{nN_L} \cdot \delta_{kn}.$$

(3°) R'_{krn} is \mathcal{C}^∞ with respect to the first variable on its domain $\mathcal{A}_0(K_{r-n})$ and $\forall \varphi' \in \omega' + \mathcal{U}'$, $\psi'_\ell \in \mathcal{U}'$ ($\ell = 1, \dots, L$), it is

$$\left| d_{\psi'_1, \dots, \psi'_L}^L R'_{krn}(S_{2^{-n}}\varphi', x) \right| \leq C'_{kr} 2^{nN_1} \cdot \delta_{kn}^{-L}.$$

PROOF OF (38): (1°) follows easily from the fact that Λ_r is identical on $\mathcal{A}_0(K_{r-1})$. The other properties will be proved by induction. For $r = 1$ this is affirmed by §6, Properties of R_{krn} . Assuming that (38) is satisfied for un certain r , we have to prove it for $r + 1$. So we have to prove that every one of the three terms on the right-hand side of the defining equality (37) satisfies (38). This is clear for $R_{k,r+1,n}(S_{2^{-n}}\varphi, x)$ due to Properties of R_{krn} , (33). We are going to prove it for $R'_{krn}(S_{2^{-n}}(\Lambda_r\varphi), x)$. Let, by the hypothesis, $\omega \in \mathcal{A}_0(K_{r+1})$. Then $\omega' := \Lambda_r(\omega) \in \mathcal{A}_0(K_r)$ and by the induction assumption we have $\mathcal{U}' \subset \mathcal{A}(K_r)$ and $C'_{kr} > 0$, both independent of n , fulfilling (2°). Now, $\mathcal{U} := \Lambda_r^{-1}\mathcal{U}'$ is a neighbourhood of zero in $\mathcal{A}(K_{r+1})$. For $\psi_j \in \mathcal{U}$ ($j = 1, \dots, \ell$) and $\varphi \in \omega + \mathcal{U}$ we have $\psi'_j := \Lambda_r \psi_j \in \mathcal{U}'$, $\varphi' := \Lambda_r \varphi \in \omega' + \mathcal{U}'$, hence (chain rule with the inner function Λ_r linear)

$$\left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R'_{krn}(S_{2^{-n}}(\Lambda_r\varphi), x) \right| = \left| d_{\psi'_1, \dots, \psi'_\ell}^\ell \partial^\alpha R'_{krn}(S_{2^{-n}}\varphi', x) \right| \leq C'_{kr} 2^{nN_L}.$$

Exactly by the same way we deduce the same estimation for the last term in (37), i.e.

$$\left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R_{k,r+1,n}(S_{2^{-n}}(\Lambda_r\varphi), x) \right| \leq C 2^{nN_L},$$

only we have to start with (33) instead of the induction assumption. Thus (38) is proved by induction. □

PROOF OF (39): Taking (39) as the induction assumption, we get by the recurrent definition (37) of R'_{krn} :

$$\begin{aligned} & \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha (R'_{k,r+1,n}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x)) \right| \\ & \leq \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha (R_{k,r+1,n}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x)) \right| \\ & \quad + \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha [R'_{k,r,n}(S_{2^{-n}}(\Lambda_r\varphi), x) - R(S_{2^{-n}}(\Lambda_r\varphi), x)] \right| \\ & \quad - d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha [R_{k,r+1,n}(S_{2^{-n}}(\Lambda_r\varphi), x) - R(S_{2^{-n}}(\Lambda_r\varphi), x)] \Big|. \end{aligned}$$

As above, we estimate every one of these three terms using (34) and the induction assumption. □

PROOF OF (3°): by §6 (3°) can be the same as the proof of (38). □

§8. **Definition of R'_{kn} .** We define

$$R'_{kn}(S_{2^{-n}}\varphi, x) = \lim_{r \rightarrow \infty} R'_{krn}(S_{2^{-n}}\varphi, x)$$

for $\varphi \in \mathcal{A}_0(\mathbb{R}^d)$ and $x \in \mathbb{R}^d$. Every φ belongs to $\mathcal{A}_0(K_{r-1})$ for some $r \in \mathbb{N}$; up from this r the sequence $\{R'_{krn}\}_r$ is constant thanks to §7, Property 1°, so it is $R'_{kn}(S_{2^{-n}}\varphi, x) = R'_{krn}(S_{2^{-n}}\varphi, x)$.

Properties of R'_{kn} . (1°). For every $k, n \in \mathbb{N}$ the function R'_{kn} is defined on $\mathcal{A}_0(\mathbb{R}^d) \times \mathbb{R}^d$.

Moreover, if $B \in \mathbb{R}^d$, \mathcal{B} a bounded set $\subset \mathcal{A}_0(B)$ and $L \in \mathbb{N}_0$, then there is an absolutely convex open neighbourhood of zero $\mathcal{U}' = \mathcal{U}'_k \subset \mathcal{A}(B)$ and a constant $C'_k > 0$, both independent of n , such that for every $n \in \mathbb{N}$, $x \in \mathbb{R}^d$ the following hold.

(2°) If $1 \leq L \leq k$, $\ell = 0, 1, \dots, L - 1$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L$, $\varphi \in \mathcal{B}$, $2^{-nN_L-1} > V_{N_L}(\varphi)$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}'$, then $\varphi, x \mapsto d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R'_{kn}(S_{2^{-n}}\varphi, x)$ is continuous and

$$(40) \quad \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha R'_{kn}(S_{2^{-n}}\varphi, x) \right| \leq C'_k 2^{nN_L},$$

$$(41) \quad \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (R'_{kn}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x)) \right| \leq C'_k 2^{nN_L} \cdot \delta_{kn}.$$

(3°) R'_{kn} is \mathcal{C}^∞ with respect to the first variable and if $\varphi \in \mathcal{B}$, $\psi_\ell \in \mathcal{U}'$ ($\ell = 1, \dots, L$), then

$$(42) \quad \left| d_{\psi_1, \dots, \psi_L}^L R'_{kn}(S_{2^{-n}}\varphi, x) \right| \leq C'_k 2^{nN_1} \cdot \delta_{kn}^{-L}.$$

PROOF: We have $R'_{kn}(S_{2^{-n}}\varphi, x) = R'_{krn}(S_{2^{-n}}\varphi, x)$ for $\varphi \in \mathcal{A}_0(K_{r-1})$ so by §7, Property (3°), R'_{kn} is smooth with respect to the first variable on every $\mathcal{A}_0(K_{r-1})$. As smoothness depends only on the behaviour of R'_{kn} on bounded sets, R'_{kn} is smooth on $\mathcal{A}_0(\mathbb{R}^d)$. For proving the estimations, we can assume without loss of generality that $B = K_{r-1}$ for some $r \in \mathbb{N}$. It is known that the bounded sets in \mathcal{D} are relatively compact; thus, for a given L , the set \mathcal{B} can be covered with a finite number of sets $\omega'_m + \mathcal{U}'_m$ where \mathcal{U}'_m is assigned to ω'_m by §7, Properties of R'_{krn} . Putting $\mathcal{U}' = \bigcap \mathcal{U}'_m$, we get the properties of R'_{kn} from Properties of R'_{krn} . \square

§9. Up to now, we have constructed functions that were \mathcal{C}^∞ with respect to the first variable. Now we are going to regularize the function R'_{kn} by convolution with respect to the second variable to obtain a simultaneously \mathcal{C}^∞ function.

Notation. The function ρ_δ is introduced in §2, Notation. If k, n are chosen, we have still $\delta = \delta_{kn} = 2^{-n(k+1)N_k}$. Denote furthermore

$$\begin{aligned} \rho^{\otimes d}(x) &:= \rho(x_1) \cdot \dots \cdot \rho(x_d), & \rho_\delta^{\otimes d}(x) &:= \rho_\delta(x_1) \cdot \dots \cdot \rho_\delta(x_d) \\ &\text{for } x = (x_1, \dots, x_d) \in \mathbb{R}^d. \end{aligned}$$

Definition. We define a function \tilde{R}_{kn} on $\mathcal{A}_0(\mathbb{R}^d) \times \mathbb{R}^d$ by convolution as follows.

$$\tilde{R}_{kn}(\varphi, x) := R'_{kn}(\varphi, x) * \rho_\delta^{\otimes d}(x) = \int R'_{kn}(\varphi, y) \rho_\delta^{\otimes d}(x - y) dy.$$

Properties of \tilde{R}_{kn} . For every $k, n \in \mathbb{N}$, \tilde{R}_{kn} is a \mathcal{C}^∞ function on $\mathcal{A}_0(\mathbb{R}^d) \times \mathbb{R}^d$. That is: $\tilde{R}_{kn} \in \mathcal{E}(\mathbb{R}^d)$. Moreover, if $B \in \mathbb{R}^d$, \mathcal{B} a bounded set $\subset \mathcal{A}_0(B)$ and $L \in \mathbb{N}$, then there is an absolutely convex open neighbourhood of zero $\mathcal{U} = \mathcal{U}_k \subset \mathcal{A}(B)$ and a constant $\tilde{C}_k > 0$, both independent of n , such that for every $n \in \mathbb{N}$, $x \in \mathbb{R}^d$, $\ell = 0, 1, \dots, L$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$, $\varphi \in \mathcal{B}$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L$, we have

$$(43) \quad \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha \tilde{R}_{kn}(S_{2^{-n}}\varphi, x) \right| \leq \tilde{C}_k 2^{nN_1} \cdot \delta_{kn}^{-2L}.$$

If in addition $L \leq k$, $2^{-nN_L-1} > V_{N_L}(\varphi)$ and $\ell \leq L - 1$, then

$$(44) \quad \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha \tilde{R}_{kn}(S_{2^{-n}}\varphi, x) \right| \leq \tilde{C}_k 2^{nN_L}.$$

If in addition $|\alpha| \leq L - 1$, then

$$(45) \quad \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (\tilde{R}_{kn}(S_{2^{-n}}\varphi, x) - R(S_{2^{-n}}\varphi, x)) \right| \leq \tilde{C}_k 2^{nN_L} \cdot \delta_{kn}.$$

PROOF OF (43):

$$\begin{aligned} \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell \tilde{R}_{kn}(S_{2^{-n}}\varphi, x) \right| &= \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (R'_{kn}(S_{2^{-n}}\varphi, x) * \rho_\delta^{\otimes d}(x)) \right| \\ &= \left| (d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, x)) * \partial^\alpha \rho_\delta^{\otimes d}(x) \right|. \end{aligned}$$

By (42), this is

$$\leq C'_k 2^{nN_1} \cdot \delta^{-L} \|\partial^\alpha \rho_\delta^{\otimes d}\|_{\mathcal{L}^1} = C'_k 2^{nN_1} \cdot \delta^{-L} \delta^{-|\alpha|} \|\partial^\alpha \rho^{\otimes d}\|_{\mathcal{L}^1}.$$

As $|\alpha| \leq L$, we obtain (43).

We see that for given k, n the derivatives $|d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha \tilde{R}_{kn}(S_{2^{-n}}\varphi, x)|$ are equibounded if $\varphi \in \mathcal{B}$, $x \in \mathbb{R}^d$, hence they are continuous on $\mathcal{B} \times \mathbb{R}^d$ for any bounded $\mathcal{B} \subset \mathcal{A}_0(B) \times \mathbb{R}^d$ ($B \in \mathbb{R}^d$). They are continuous on $\mathcal{A}_0(B) \times \mathbb{R}^d$ because they are continuous on convergent sequences in a metric space. Thus the order of taking derivatives does not matter and \tilde{R}_{kn} is smooth ([13, 1.11.5.(2^o)]). \square

PROOF OF (44):

$$\begin{aligned} \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell \tilde{R}_{kn}(S_{2^{-n}}\varphi, x) \right| &= \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell (R'_{kn}(S_{2^{-n}}\varphi, x) * \rho_\delta^{\otimes d}(x)) \right| \\ &= \left| (\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, x)) * \rho_\delta^{\otimes d}(x) \right|. \end{aligned}$$

Then we deduce easily (44) from (40). \square

PROOF OF (45): We first estimate

$$\begin{aligned} &\left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell \tilde{R}_{kn}(S_{2^{-n}}\varphi, x) - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, x) \right| \\ &= \left| \int \left(\partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, y) - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, x) \right) \rho_\delta^{\otimes d}(x - y) \, dy \right| \\ &\leq \sup \left\{ \left| \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, y) - \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, x) \right| ; \right. \\ &\quad \left. |x_1 - y_1| \leq \delta, \dots, |x_d - y_d| \leq \delta \right\} \leq C'_k 2^{nN_L} \cdot d \delta \end{aligned}$$

because the function $x \mapsto \partial^\alpha d_{\psi_1, \dots, \psi_\ell}^\ell R'_{kn}(S_{2^{-n}}\varphi, x)$ has its derivatives of order 1 estimated by (40). Then (45) follows from (41). \square

§10. Lemma. *Let \mathcal{B} be a bounded set in $\mathcal{D}(\mathbb{R}^d)$. Then there is a natural number k such that $V'_{N_k}(\varphi) \geq 2^{-k}$ for all $\varphi \in \mathcal{B}$ (V'_N defined by §2, Notation).*

As V'_N is non-decreasing, this inequality holds for all sufficiently large k .

PROOF: If not, there would be a sequence $\{\varphi_k\}_{k=1}^\infty \subset \mathcal{B}$ such that $V'_{N_k}(\varphi_k) < 2^{-k}$ ($\forall k$). As $\overline{\mathcal{B}}$ is a metrizable compact, a subsequence $\{\varphi_{k_n}\}_{n=1}^\infty$ is convergent in \mathcal{A}_0 , $\lim \varphi_{k_n} = \varphi \in \mathcal{A}_0$. We have $V'_N(\varphi_k) \leq V'_{N_k}(\varphi_k) < 2^{-k}$ for $N \leq N_k$. As $N_k \nearrow \infty$ (see §3, Properties of R), we have $0 \leq V'_N(\varphi) \leq \lim 2^{-k} = 0$ for all $N \in \mathbb{N}$, that is impossible. (Proof: If φ has all moments of order ≥ 1 equal to 0, then its Fourier transform has zero derivatives at origin; being holomorphic, it must be constant). \square

Notation. Denote $\vartheta_{kn}(\varphi) := \vartheta(2^{nN_k+k+1} \cdot V'_{N_k}(\varphi))$ (ϑ by §2, Notation). ϑ_{kn} is \mathcal{C}^∞ on \mathcal{A}_0 and

$$\begin{aligned} \vartheta_{kn}(\varphi) &= 0 & \text{if } 2^{nN_k+k+1} \cdot V'_{N_k}(\varphi) \geq 2, \\ \vartheta_{kn}(\varphi) &= 1 & \text{if } 2^{nN_k+k+1} \cdot V'_{N_k}(\varphi) \leq 1. \end{aligned}$$

Definition. We define

$$(46) \quad \tilde{R}_n(\varphi, x) := \sum_{k=1}^\infty (\vartheta_{kn}(\varphi) - \vartheta_{k+1,n}(\varphi)) \cdot \tilde{R}_{kn}(\varphi, x)$$

$$(47) \quad = \sum_{k=1}^\infty \vartheta_{kn}(\varphi) \cdot (\tilde{R}_{kn}(\varphi, x) - \tilde{R}_{k-1,n}(\varphi, x))$$

if we set $\tilde{R}_{0,n} = 0$.

Remark. For a given n and φ , at most two terms of the sum (46) are nonzero. Only a finite number of terms of the sum (47) are nonzero: if k satisfies the above lemma, then $\vartheta_{kn}(\varphi) = 0$.

$$(48) \quad \sum_{k=1}^\infty (\vartheta_{kn}(\varphi) - \vartheta_{k+1,n}(\varphi)) = 1$$

is a smooth partition of unity on $\{\varphi \in \mathcal{A}_0; V'_{N_1}(\varphi) < 2^{-nN_1-2}\}$. Indeed, the sequence $\left\{2^{nN_k+k+1}V'_{N_k}(\varphi)\right\}_k$ is non-decreasing and its first member is $2^{nN_1+2}V'_{N_1}(\varphi) < 1$. Thanks to Lemma, there is the greatest index k' for which $2^{nN_{k'}+k'+1}V'_{N_{k'}}(\varphi) \leq 1$. Then $2^{nN_{k'+1}+k'+2}V'_{N_{k'+1}}(\varphi) > 1$, $2^{nN_{k'+2}+k'+3}V'_{N_{k'+2}}(\varphi) > 2$, hence

$$(\vartheta_{k',n}(\varphi) - \vartheta_{k'+1,n}(\varphi)) + (\vartheta_{k'+1,n}(\varphi) - \vartheta_{k'+2,n}(\varphi)) = 1$$

and the other terms of (48) are zero.

Properties of \tilde{R}_n . $\tilde{R}_n \in \mathcal{E}(\mathbb{R}^d)$. If $\tilde{B} \in \mathbb{R}^d$, $\tilde{\mathcal{B}} \subset \mathcal{A}_0(\tilde{B})$ is a bounded set and $L \in \mathbb{N}$, then there is an absolutely convex open neighbourhood of zero $\tilde{\mathcal{U}} \subset \mathcal{A}(\tilde{B})$ and a constant $\tilde{C} > 0$, both independent of n , such that for every $n \in \mathbb{N}$, $x \in \mathbb{R}^d$, the following hold:

If $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L - 1$, $\ell \in \mathbb{N}_0$, $\ell \leq L - 1$, $\tilde{\varphi} \in \tilde{\mathcal{B}}$ with $\|\tilde{\varphi}\|_{\mathcal{L}^2} \geq 1$, $\tilde{\psi}_1, \dots, \tilde{\psi}_\ell \in \tilde{\mathcal{U}}$, then

$$(49) \quad \left| d_{\tilde{\psi}_1, \dots, \tilde{\psi}_\ell}^\ell \partial^\alpha \tilde{R}_n(S_{2^{-n}} \tilde{\varphi}, x) \right| \leq \tilde{C} 2^{nN_L(L+1)(2L+1)}.$$

If $(\forall q \in \mathbb{N}) \tilde{\varphi} \in \tilde{\mathcal{B}} \cap \mathcal{A}_{N_q}$ with $\|\tilde{\varphi}\|_{\mathcal{L}^2} \geq 1$ $q \in \mathbb{N}$, then

$$(50) \quad \left| \tilde{R}_n(S_{2^{-n}} \tilde{\varphi}, x) - R(S_{2^{-n}} \tilde{\varphi}, x) \right| \leq \tilde{C} 2^{-nq}.$$

PROOF OF (49): For a nonzero term of (46) or (47), we have $2^{nN_k+k+1} \cdot V'_{N_k}(\tilde{\varphi}) < 2$, i.e. $V'_{N_k}(\tilde{\varphi}) < 2^{-nN_k-k}$. If $\|\tilde{\varphi}\|_{\mathcal{L}^2} > 1$, then (§2, Notation) $V_{N_k}(\tilde{\varphi}) \leq V'_{N_k}(\tilde{\varphi}) < 2^{-nN_k-k}$, so the hypothesis $V_{N_L}(\tilde{\varphi}) < 2^{-nN_L-1}$ ($L \leq k$) in §9, Properties of \tilde{R}_{kn} for (44) and (45) is always satisfied. By the Leibniz rule (formulated in the proof of §6, Property (2°)) applied to the definition (47) of \tilde{R}_n (recall that $V'_N(\tilde{\varphi}) = V'_N(S_{2^{-n}} \tilde{\varphi})$), we have

$$(51) \quad d_{\tilde{\psi}_1, \dots, \tilde{\psi}_\ell}^\ell \partial^\alpha \tilde{R}_n(S_{2^{-n}} \tilde{\varphi}, x) = \sum_{k=1}^\infty \sum_{\substack{I_1 \cup I_2 = \{1, \dots, \ell\} \\ \text{disjoint}}} d_{\tilde{\psi}_{I_1}}^{\#I_1} \vartheta_{kn}(\tilde{\varphi}) \cdot d_{\tilde{\psi}_{I_2}}^{\#I_2} \partial^\alpha \left(\tilde{R}_{kn}(S_{2^{-n}} \tilde{\varphi}, x) - \tilde{R}_{k-1, n}(S_{2^{-n}} \tilde{\varphi}, x) \right)$$

(If $I = (i_1, \dots, i_{\#I})$, then $\tilde{\psi}_I$ denotes $(\tilde{\psi}_{i_1}, \dots, \tilde{\psi}_{i_{\#I}})$). Due to Lemma, the sum $\sum_{k=1}^\infty$ can be replaced with $\sum_{k=1}^{k_0}$ with a number k_0 depending on $\tilde{\mathcal{B}}$ but not on $\tilde{\varphi} \in \tilde{\mathcal{B}}$.

First we estimate

$$\begin{aligned} d_{\tilde{\psi}_I}^{\#I} (\vartheta_{kn}(\tilde{\varphi})) &= d_{\tilde{\psi}_I}^{\#I} \vartheta(2^{nN_k+k+1} \cdot V'_{N_k}(\tilde{\varphi})) \\ &= \sum_{M=1}^{\#I} \sum_{\substack{I = I_1 \cup \dots \cup I_M \\ \neq \emptyset, \text{ disjoint}}} (\partial^M \vartheta)(2^{nN_k+k+1} \cdot V'_{N_k}(\tilde{\varphi})) \prod_{m=1}^M d_{\tilde{\psi}_{I_m}}^{\#I_m} (2^{nN_k+k+1} \cdot V'_{N_k}(\tilde{\varphi})) \end{aligned}$$

(chain rule [13, 1.8.3] or [8, Theorem 12], the summation is extended over all decompositions $I = I_1 \cup \dots \cup I_M$ on non-empty disjoint parts). Let us choose

(by [9, §2, Proposition]) an absolutely convex open neighbourhood of zero $\tilde{\mathcal{U}} \subset \mathcal{A}(\tilde{B})$ such that for every $k = 1, \dots, k_0, \ell = 1, \dots, L, \tilde{\varphi} \in \tilde{\mathcal{B}}, \tilde{\psi}_1, \dots, \tilde{\psi}_\ell \in \tilde{\mathcal{U}}$, we have $|\mathfrak{d}_{\tilde{\psi}_1, \dots, \tilde{\psi}_\ell}^\ell V'_{N_k}(\tilde{\varphi})| \leq 1$. We obtain:

$$(52) \quad \left| \mathfrak{d}_{\tilde{\psi}_I}^{\#I} (\vartheta_{kn}(\tilde{\varphi})) \right| \leq C_k 2^{nN_k \cdot \#I} \leq C_k 2^{nN_k L}$$

with a constant C_k depending only on ϑ, L and k , not on n .

We are going to estimate the second term in (51)

$$\mathfrak{d}_{\psi_{I_2}}^{\#I_2} \partial^\alpha \left(\tilde{R}_{kn}(S_{2^{-n}}\tilde{\varphi}, x) - \tilde{R}_{k-1,n}(S_{2^{-n}}\tilde{\varphi}, x) \right).$$

We distinguish two cases. If $L \leq k - 1$ then $\#I_2 \leq k - 1$ and we can use the estimation (45) as follows:

$$\begin{aligned} & \left| \mathfrak{d}_{\psi_{I_2}}^{\#I_2} \partial^\alpha \left(\tilde{R}_{kn}(S_{2^{-n}}\tilde{\varphi}, x) - \tilde{R}_{k-1,n}(S_{2^{-n}}\tilde{\varphi}, x) \right) \right| \\ &= \left| \mathfrak{d}_{\psi_{I_2}}^{\#I_2} \partial^\alpha \left(\left(\tilde{R}_{kn}(S_{2^{-n}}\tilde{\varphi}, x) - R(S_{2^{-n}}\tilde{\varphi}, x) \right) \right. \right. \\ & \quad \left. \left. - \left(\tilde{R}_{k-1,n}(S_{2^{-n}}\tilde{\varphi}, x) - R(S_{2^{-n}}\tilde{\varphi}, x) \right) \right) \right| \\ &\leq 2^{nN_L} \cdot (\tilde{C}_k \delta_{kn} + \tilde{C}_{k-1} \delta_{k-1,n}) \leq 2^{nN_L} \cdot (\tilde{C}_k + \tilde{C}_{k-1}) \delta_{k-1,n}. \end{aligned}$$

Together with (52), a term of the sum in (51) for $L \leq k - 1$ fulfills

$$\begin{aligned} & \left| \mathfrak{d}_{\psi_{I_1}}^{\#I_1} \vartheta_{kn}(\tilde{\varphi}) \cdot \mathfrak{d}_{\psi_{I_2}}^{\#I_2} \partial^\alpha \left(\tilde{R}_{kn}(S_{2^{-n}}\tilde{\varphi}, x) - \tilde{R}_{k-1,n}(S_{2^{-n}}\tilde{\varphi}, x) \right) \right| \\ &\leq C_k 2^{nN_k L} \cdot 2^{nN_L} (\tilde{C}_k + \tilde{C}_{k-1}) \delta_{k-1,n} \leq C_k (\tilde{C}_k + \tilde{C}_{k-1}) \end{aligned}$$

(δ is defined in §9, Notation), that is a constant independent of n , however it depends on \mathcal{B} and the number of nonzero terms of the sum in (51) depends on \mathcal{B} , too.

If $L \geq k$, we use the estimations (43) valid for all L and we obtain:

$$\mathfrak{d}_{\psi_{I_2}}^{\#I_2} \partial^\alpha \tilde{R}_{kn}(S_{2^{-n}}\tilde{\varphi}, x) \leq \tilde{C}_k 2^{nN_1} \cdot \delta_{kn}^{-2L} \leq \tilde{C}_k 2^{nN_1} \cdot \delta_{L,n}^{-2L}.$$

Together with (52), a term of the sum in (51) for $L \geq k$ fulfills

$$\begin{aligned} & \left| \mathfrak{d}_{\psi_{I_1}}^{\#I_1} \vartheta_{kn}(\tilde{\varphi}) \cdot \mathfrak{d}_{\psi_{I_2}}^{\#I_2} \partial^\alpha \left(\tilde{R}_{kn}(S_{2^{-n}}\tilde{\varphi}, x) - \tilde{R}_{k-1,n}(S_{2^{-n}}\tilde{\varphi}, x) \right) \right| \\ &\leq C_k 2^{nN_k L} (\tilde{C}_k + \tilde{C}_{k-1}) 2^{nN_1} \cdot \delta_{L,n}^{-2L} \leq C_k (\tilde{C}_k + \tilde{C}_{k-1}) 2^{nN_L(2L+1)(L+1)}. \end{aligned}$$

So in both cases we can use the last estimation and (49) follows. □

PROOF OF (50): If $\tilde{\varphi} \in \mathcal{A}_{N_q}$ then $V'_{N_q}(\tilde{\varphi}) = 0$ and $k \geq q$ for all nonzero terms of the sum in (46). In that case, it follows from the definition and (48)

$$\begin{aligned} & \left| \tilde{R}_n(S_\varepsilon \tilde{\varphi}, x) - R(S_\varepsilon \tilde{\varphi}, x) \right| \\ &= \left| \sum_{k=q}^{k_0} (\vartheta_{kn}(\tilde{\varphi}) - \vartheta_{k+1,n}(\tilde{\varphi})) \cdot \tilde{R}_{kn}(S_\varepsilon \tilde{\varphi}, x) - \sum_{k=q}^{k_0} (\vartheta_{kn}(\tilde{\varphi}) - \vartheta_{k+1,n}(\tilde{\varphi})) \cdot R(S_\varepsilon \tilde{\varphi}, x) \right| \\ &\leq \sum_{k=q}^{k_0} (\vartheta_{kn}(\tilde{\varphi}) - \vartheta_{k+1,n}(\tilde{\varphi})) \cdot \left| \tilde{R}_{kn}(S_\varepsilon \tilde{\varphi}, x) - R(S_\varepsilon \tilde{\varphi}, x) \right| \\ &\leq \sum_{k=q}^{k_0} \tilde{C}_k 2^{nN_1} \delta_{kn} = \sum_{k=q}^{k_0} \tilde{C}_k 2^{nN_1 - n(k+1)N_k} \leq \tilde{C} 2^{-nq} \end{aligned}$$

by (45), as $V_{N_1}(\tilde{\varphi}) = 0$. □

§11. Now we have all tools for defining the desired representative \tilde{R} .

Definition of \tilde{R} . We define

$$\tilde{R}(\varphi, x) := \sum_{n=1}^{\infty} \left(\vartheta_{n+1} \left(\|\varphi\|_{\mathcal{L}^2}^{2/d} \right) - \vartheta_n \left(\|\varphi\|_{\mathcal{L}^2}^{2/d} \right) \right) \tilde{R}_n(\varphi, x)$$

for $\varphi \in \mathcal{A}_0(\mathbb{R}^d)$ and $x \in \mathbb{R}^d$ (ϑ by §2, Notation).

Remark. Note that $\vartheta_{n+1} \left(\|\varphi\|_{\mathcal{L}^2}^{2/d} \right) - \vartheta_n \left(\|\varphi\|_{\mathcal{L}^2}^{2/d} \right) \neq 0$ iff $2^n < \|\varphi\|_{\mathcal{L}^2}^{2/d} < 2^{n+2}$ and, for a given φ , at most 2 terms of this sum are $\neq 0$.

$$\sum_{n=1}^{\infty} \left(\vartheta_{n+1} \left(\|\varphi\|_{\mathcal{L}^2}^{2/d} \right) - \vartheta_n \left(\|\varphi\|_{\mathcal{L}^2}^{2/d} \right) \right) = 1$$

is a smooth partition of unity on $\{\varphi \in \mathcal{D}; \|\varphi\|_{\mathcal{L}^2} > 4\}$.

Properties of \tilde{R} . $\tilde{R} \in \mathcal{E}(\mathbb{R}^d)$. If $B \in \mathbb{R}^d$, \mathcal{B} a bounded set $\subset \mathcal{A}_0(B)$ and $L \in \mathbb{N}$, then there is an absolutely convex open neighbourhood of zero $\mathcal{U} \subset \mathcal{A}(B)$ and a constant $C > 0$, such that for every $x \in \mathbb{R}^d$ the following hold.

(1°) If $\varepsilon \in]0, 1]$, $\alpha \in \mathbb{N}_0^d$, $|\alpha| \leq L - 1$, $\ell \in \mathbb{N}_0$, $\ell \leq L - 1$, $\varphi \in \mathcal{B}$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$, then

$$(53) \quad \left| d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha \tilde{R}(S_\varepsilon \varphi, x) \right| \leq C \varepsilon^{-N_L(2L+1)(L+1)}.$$

Consequently, $\tilde{R} \in \mathcal{E}_M^d$.

(2°) $\forall q \in \mathbb{N}$ it holds: If $\varphi \in \mathcal{B} \cap \mathcal{A}_{N_q}$ and $0 < \varepsilon < \min \left\{ 1, \frac{1}{4} \|\chi_B\|_{\mathcal{L}^2}^{-2/d} \right\}$, then

$$(54) \quad \left| \tilde{R}(S_\varepsilon \varphi, x) - R(S_\varepsilon \varphi, x) \right| \leq C \cdot \varepsilon^q.$$

Due to [9, §8, (0°)], we have $\tilde{R} - R \in \mathcal{N}$.

PROOF OF (53): We write simply $\|\bullet\|$ instead of $\|\bullet\|_{\mathcal{L}^2}$ and we assume B to be convex and balanced. For proving (53), we calculate (see (2)):

$$(55) \quad \begin{aligned} \tilde{R}(S_\varepsilon \varphi, x) &= \sum_{n=1}^{\infty} \left(\vartheta_{n+1}(\|S_\varepsilon \varphi\|^{2/d}) - \vartheta_n(\|S_\varepsilon \varphi\|^{2/d}) \right) \tilde{R}_n(S_\varepsilon \varphi, x) \\ &= \sum_{n=1}^{\infty} \left(\vartheta_1(2^{-n} \|S_\varepsilon \varphi\|^{2/d}) - \vartheta_0(2^{-n} \|S_\varepsilon \varphi\|^{2/d}) \right) \tilde{R}_n(S_{2^{-n}}(S_{2^n \varepsilon} \varphi), x) \\ &= \sum_{n=1}^{\infty} \left(\vartheta_1(\|S_{2^n \varepsilon} \varphi\|^{2/d}) - \vartheta_0(\|S_{2^n \varepsilon} \varphi\|^{2/d}) \right) \tilde{R}_n(S_{2^{-n}}(S_{2^n \varepsilon} \varphi), x). \end{aligned}$$

By the definition of ϑ , a term of this sum can be nonzero only if

$$(56) \quad 1 < \|S_{2^n \varepsilon} \varphi\|^{2/d} < 4, \quad \text{i.e.} \quad 1 < \frac{1}{2^n \varepsilon} \|\varphi\|^{2/d} < 4.$$

$\overline{\mathcal{B}} \in \mathcal{A}_0(B)$, hence there are constants $c_1, c_2 > 0$ such that

$$c_2 \leq \|\varphi\|^{2/d} \leq c_1 \quad (\forall \varphi \in \mathcal{B}).$$

Due to (56), it follows that for nonzero terms of the sum in (55), we have

$$(57) \quad \frac{1}{4} c_2 < 2^n \varepsilon < c_1.$$

By the Leibniz rule (formulated in the proof of §6, Property 2°) applied to (55), we have

$$(58) \quad \begin{aligned} d_{\psi_1, \dots, \psi_\ell}^\ell \partial^\alpha \tilde{R}(S_\varepsilon \varphi, x) &= \sum_{n=1}^{\infty} \sum_{\substack{I_1 \cup I_2 = \{1, \dots, \ell\} \\ \text{disjoint}}} d_{\psi_{I_1}}^{\#I_1} \left(\vartheta_1(\|S_{2^n \varepsilon} \varphi\|^{2/d}) - \vartheta_0(\|S_{2^n \varepsilon} \varphi\|^{2/d}) \right) \\ &\quad \cdot d_{\psi_{I_2}}^{\#I_2} \partial^\alpha \tilde{R}_n(S_{2^{-n}}(S_{2^n \varepsilon} \varphi), x). \end{aligned}$$

The function $\varphi \mapsto \vartheta_1(\|S_{2^n \varepsilon} \varphi\|^{2/d}) - \vartheta_0(\|S_{2^n \varepsilon} \varphi\|^{2/d})$ is composed of functions

$$[t \mapsto \vartheta_1(t^{1/d}) - \vartheta_0(t^{1/d})] \in \mathcal{D}([(1, 4^d)])$$

and

$$\varphi \mapsto \|S_{2^n \varepsilon} \varphi\|^2.$$

Hence, for proving that the term $d_{\psi_{I_1}}^{\#I_1}(\vartheta_1(\|S_{2^n \varepsilon} \varphi\|^{2/d}) - \vartheta_0(\|S_{2^n \varepsilon} \varphi\|^{2/d}))$ in (58) is equi-bounded under the hypotheses (1°) (for a fixed L), it is sufficient to prove the same for derivatives of $\|S_{2^n \varepsilon} \varphi\|^2$ up to a certain order. We have

$$\begin{aligned} d_{\psi} \|S_{2^n \varepsilon} \varphi\|^2 &= d_{\psi}(S_{2^n \varepsilon} \varphi, S_{2^n \varepsilon} \varphi) = 2\Re(S_{2^n \varepsilon} \varphi, S_{2^n \varepsilon} \psi), \\ d_{\psi_1, \psi_2}^2 \|S_{2^n \varepsilon} \varphi\|^2 &= 2\Re(S_{2^n \varepsilon} \psi_1, S_{2^n \varepsilon} \psi_2) \end{aligned}$$

and the higher derivatives are zero. So we have using the Hölder inequality

$$\left| d_{\psi} \|S_{2^n \varepsilon} \varphi\|^2 \right| = \left| 2\Re \int S_{2^n \varepsilon} \varphi \cdot S_{2^n \varepsilon} \bar{\psi} \right| \leq \|S_{2^n \varepsilon} \varphi\| \|S_{2^n \varepsilon} \psi\| = \frac{1}{(2^n \varepsilon)^d} \|\varphi\| \|\psi\|.$$

Thanks to (57), this will be equi-bounded if $\mathcal{U} \subset \{\psi; \|\psi\| < 1\}$, $\psi \in \mathcal{U}$, $\varphi \in \mathcal{B}$, as the bounded set \mathcal{B} is absorbed by \mathcal{U} . The same can be deduced for the second derivative.

Now, we have to estimate the term $d_{\psi_{I_2}}^{\#I_2} \partial^\alpha \tilde{R}_n(S_{2^{-n}}(S_{2^n \varepsilon} \varphi), x)$ in (58). We apply §10, Properties of \tilde{R}_n , namely the estimation (49), to the bounded set

$$\tilde{\mathcal{B}} := \{S_\eta \varphi; \varphi \in \mathcal{B}, \frac{1}{4}c_2 \leq \eta \leq c_1\}.$$

The supports of the functions $\tilde{\varphi} \in \tilde{\mathcal{B}}$ are contained in $\tilde{B} := c_1 B$ (for B convex and balanced). Thanks to (57) we have $S_{2^n \varepsilon} \varphi \in \tilde{\mathcal{B}}$ for $\varphi \in \mathcal{B}$. Thus we get $\tilde{\mathcal{U}}$ and \tilde{C} by §10, Properties of \tilde{R}_n . Let

$$\mathcal{U} := \left\{ \varphi \in \mathcal{A}(B); S_\eta \varphi \in \tilde{\mathcal{U}} \quad \forall \eta \quad \text{with} \quad \frac{1}{4}c_2 \leq \eta \leq c_1 \right\}.$$

\mathcal{U} is a neighbourhood of zero in $\mathcal{A}(B)$ because it absorbs bounded sets in a metric vector space. If $\varphi \in \mathcal{B}$, $\psi_1, \dots, \psi_\ell \in \mathcal{U}$, then $\tilde{\varphi} := S_{2^n \varepsilon} \varphi \in \tilde{\mathcal{B}}$ and $\tilde{\psi}_j := S_{2^n \varepsilon} \psi_j \in \tilde{\mathcal{U}}$ ($j = 1, \dots, \ell$) due to (57). If $\|\tilde{\varphi}\| = \|S_{2^n \varepsilon} \varphi\| < 1$, then (by (56)) the term $d_{\psi_{I_2}}^{\#I_2} \partial^\alpha \tilde{R}_n(S_{2^{-n}}(S_{2^n \varepsilon} \varphi), x)$ in (58) is multiplied by zero, so we have to estimate this term only if $\|\tilde{\varphi}\| \geq 1$, hence we can use (49). It follows using the chain rule for the inner function $S_{2^n \varepsilon}$ linear:

$$\begin{aligned} \left| d_{\psi_{I_2}}^{\#I_2} \partial^\alpha \tilde{R}_n(S_{2^{-n}}(S_{2^n \varepsilon} \varphi), x) \right| &= \left| d_{\psi_{I_2}}^{\#I_2} \partial^\alpha \tilde{R}_n(S_{2^{-n}} \tilde{\varphi}, x) \right| \\ &\leq \tilde{C} \cdot 2^n N_L(2L+1)(L+1) \leq \tilde{C} c_1^{N_L(2L+1)(L+1)} \cdot \varepsilon^{-N_L(2L+1)(L+1)} \end{aligned}$$

due to (57). From (58) we deduce (53). □

PROOF OF (54): By (9), $\|\varphi\| \geq \|\chi_B\|^{-1}$ (norms in \mathcal{L}^2). If $\varepsilon < \frac{1}{4}\|\chi_B\|^{-2/d}$, then

$$\|S_\varepsilon\varphi\|^{2/d} = \|\varphi\|^{2/d} \cdot \varepsilon^{-1} > \|\varphi\|^{2/d} \cdot 4 \|\chi_B\|^{2/d} \geq 4.$$

Under this hypothesis we have by §11, Remark similarly to (55):

$$\begin{aligned} R(S_\varepsilon\varphi, x) &= \sum_{n=1}^{\infty} \left(\vartheta_{n+1}(\|S_\varepsilon\varphi\|^{2/d}) - \vartheta_n(\|S_\varepsilon\varphi\|^{2/d}) \right) R(S_\varepsilon\varphi, x) \\ &= \sum_{n=1}^{\infty} \left(\vartheta_1(\|S_{2^n\varepsilon}\varphi\|^{2/d}) - \vartheta_0(\|S_{2^n\varepsilon}\varphi\|^{2/d}) \right) R(S_{2^{-n}}(S_{2^n\varepsilon}\varphi), x). \end{aligned}$$

With (55) it gives

$$\begin{aligned} &\left| \tilde{R}(S_\varepsilon\varphi, x) - R(S_\varepsilon\varphi, x) \right| \\ &\leq \sum_{n=1}^{\infty} \left(\vartheta_1(\|S_{2^n\varepsilon}\varphi\|^{2/d}) - \vartheta_0(\|S_{2^n\varepsilon}\varphi\|^{2/d}) \right) \\ &\quad \cdot \left| \tilde{R}_n(S_{2^{-n}}(S_{2^n\varepsilon}\varphi), x) - R(S_{2^{-n}}(S_{2^n\varepsilon}\varphi), x) \right|. \end{aligned}$$

If $\varphi \in \mathcal{B} \cap \mathcal{A}_{N_q}$ then $\tilde{\varphi} := S_{2^n\varepsilon}\varphi \in \tilde{\mathcal{B}} \cap \mathcal{A}_{N_q}$. By (50) this is $\leq \tilde{C} \cdot 2^{-nq}$ and by (57) this is $\leq \tilde{C} \cdot \left(\frac{4}{c_2}\right)^q \varepsilon^q$. By the above lemma, $\mathcal{B} \cap \mathcal{A}_{N_q} = \emptyset$ for sufficiently large q , so the constant in our estimation can be independent of q . Hence, (54) is proved. □

REFERENCES

- [1] Colombeau J.F., *Differential Calculus and Holomorphy*, North Holland MathStudies 64, 1982.
- [2] Colombeau J.F., *New Generalized Functions and Multiplication of Distributions*, North Holland MathStudies 84, 1984.
- [3] Colombeau J.F., *Elementary Introduction to New Generalized Functions*, North Holland MathStudies 113, 1985.
- [4] Colombeau J.F., Meril A., *Generalized Functions and Multiplication of Distributions on \mathcal{C}^∞ Manifolds*, J. Math. Anal. Appl. **186** (1994), 357–364.
- [5] Deville R., Godefroy G., Zizler V., *Smoothness and Renormings in Banach Spaces*, John Wiley, New York, 1993.
- [6] Grosser M., Farkas E., Kunzinger M., Steinbauer R., *On the foundations of nonlinear generalized functions I, II*, Mem. Amer. Math. Soc. **153** (2001), no. 729, 93 pp.
- [7] Grosser M., Kunzinger M., Steinbauer R., Vickers J., *A global theory of algebras of generalized functions*, Adv. Math. **166** (2002), no. 1, 50–72.
- [8] Jelínek J., *An intrinsic definition of the Colombeau generalized functions*, Comment. Math. Univ. Carolinae **40.1** (1999), 71–95.

- [9] Jelínek J., *On introduction of two diffeomorphism invariant Colombeau algebras*, Comment. Math. Univ. Carolinae **45.4** (2004), 615–632.
- [10] Kuratowski, *Topology*, Academic Press, New York-London; PWN, Warsaw, 1966.
- [11] Robertson A.P., Robertson W.J., *Topological Vector Spaces*, Cambridge Univ. Press, New York, 1964.
- [12] Schwartz L., *Théorie des Distributions*, Hermann, Paris, 1966.
- [13] Yamamuro S., *Differential Calculus in Topological Linear Spaces*, Lecture Notes in Math. 374, Springer, Berlin-New York, 1974.

DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS,
CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

E-mail: jelinek@karlin.mff.cuni.cz

(Received July 6, 2001, revised December 13, 2002)