

# Applications of Mathematics

---

Haishen Lü; Donal O'Regan; Ravi P. Agarwal

On the existence of multiple periodic solutions for the vector  $p$ -Laplacian via critical point theory

*Applications of Mathematics*, Vol. 50 (2005), No. 6, 555–568

Persistent URL: <http://dml.cz/dmlcz/134623>

## Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE EXISTENCE OF MULTIPLE PERIODIC SOLUTIONS  
FOR THE VECTOR  $p$ -LAPLACIAN VIA  
CRITICAL POINT THEORY\*

HAISHEN LÜ, Nanjing, DONAL O'REGAN, Galway,  
and RAVI P. AGARWAL, Melbourne

(Received November 20, 2003, in revised version June 24, 2004)

*Abstract.* We study the vector  $p$ -Laplacian

$$(*) \quad \begin{cases} -(|u'|^{p-2}u')' = \nabla F(t, u) & \text{a.e. } t \in [0, T], \\ u(0) = u(T), \quad u'(0) = u'(T), & 1 < p < \infty. \end{cases}$$

We prove that there exists a sequence  $(u_n)$  of solutions of  $(*)$  such that  $u_n$  is a critical point of  $\varphi$  and another sequence  $(u_n^*)$  of solutions of  $(*)$  such that  $u_n^*$  is a local minimum point of  $\varphi$ , where  $\varphi$  is a functional defined below.

*Keywords:*  $p$ -Laplacian equation, periodic solution, critical point theory

*MSC 2000:* 34B15

## 1. INTRODUCTION AND MAIN RESULTS

Consider the second order system

$$(1.1) \quad \begin{cases} -(|u'|^{p-2}u')' = \nabla F(t, u) & \text{a.e. } t \in [0, T], \\ u(0) = u(T), \quad u'(0) = u'(T), & 1 < p < \infty, \end{cases}$$

where  $T > 0$  and  $F: [0, T] \times \mathbb{R}^N \rightarrow \mathbb{R}$  satisfies the following assumption:

---

\* The research is supported by NNSF of China (10301033).

(A)  $F(t, x)$  is measurable in  $t$  for every  $x \in \mathbb{R}^N$  and continuously differentiable in  $x$  for a.e.  $t \in [0, T]$ , and there exist  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $b \in L^1(0, T; \mathbb{R}^+)$  such that

$$|F(t, x)| \leq a(|x|)b(t), \quad |\nabla F(t, x)| \leq a(|x|)b(t)$$

for all  $x \in \mathbb{R}^N$  and a.e.  $t \in [0, T]$ ; here  $\mathbb{R}^+ = [0, \infty)$ .

Recently some papers have appeared discussing scalar periodic problems driven by the one-dimensional  $p$ -Laplacian. We refer the reader to the works of Del Pino-Manasevich-Murua [3], Fabry-Fayyad [4], Gao [5] and Dang-Opppenheimer [6]. In all of these works the approach is degree theoretical and the existence of one solution is established. In [2], J. Mawhin generalized the Hartman-Knobloch results to perturbations of a vector  $p$ -Laplacian ordinary operator.

For  $p = 2$ , Mawhin-Willem [1] proved the existence of solutions for the problem (1.1) under the conditions

(M1) there exists  $g \in L^1(0, T; \mathbb{R}^+)$  such that  $|\nabla F(t, x)| \leq g(t)$  for all  $x \in \mathbb{R}^N$  and a.e.  $t \in [0, T]$ , and

(M2)  $\int_0^T F(t, x) dt \rightarrow +\infty$  as  $|x| \rightarrow \infty$  or  $\int_0^T F(t, x) dt \rightarrow -\infty$  as  $|x| \rightarrow \infty$ .

In this paper we study the existence of periodic solutions of a vector  $p$ -Laplacian with potential oscillating around the first eigenvalue. Our arguments are based on a result by Habets-Manasevich-Zanolin [8] related to the two-point boundary value problem

$$u'' + u + f(t, u) = 0, \quad u(0) = u(1) = 0.$$

Our main result is the following theorem.

**Theorem 1.1.** *Suppose*

(H1) *there exists  $g \in L^q(0, T; \mathbb{R}^+)$  (here  $q$  is the conjugate of  $p$ ) such that  $|\nabla F(t, x)| \leq g(t)$  for all  $x \in \mathbb{R}^N$  and a.e.  $t \in [0, T]$ ,*

(H2)

$$(1.2) \quad \limsup_{R \rightarrow +\infty} \inf_{a \in \mathbb{R}^N, |a|=R} \int_0^T F(s, a) ds = +\infty$$

and

$$(1.3) \quad \liminf_{r \rightarrow +\infty} \sup_{b \in \mathbb{R}^N, |b|=r} \int_0^T F(s, b) ds = -\infty$$

*hold. Then*

- i) *there exists a sequence  $(u_n)$  of solutions of (1.1) such that  $u_n$  is a critical point of  $\varphi$  and  $\lim_{n \rightarrow \infty} \varphi(u_n) = +\infty$ ;*

ii) *there exists a sequence  $(u_n^*)$  of solutions of (1.1) such that  $u_n^*$  is a local minimum point of  $\varphi$  and  $\lim_{n \rightarrow \infty} \varphi(u_n^*) = -\infty$ ; here  $\varphi$  is a functional defined below.*

Throughout the paper, for  $N \geq 1$  and  $I = [0, T]$ , we will set  $C = C(I, \mathbb{R}^N)$ ,  $L^p = L^p(I, \mathbb{R}^N)$ ,  $W^{1,p} = W^{1,p}(I, \mathbb{R}^N)$  and  $X = W_T^{1,p} = \{u \in W^{1,p}; u(0) = u(T)\}$ . The norm in  $C$  will be defined by  $\|u\|_\infty = \max_{t \in [0, T]} |u(t)|$ , the norm in  $L^p$  by

$$\|u\|_p = \left( \int_0^T |u(t)|^p dt \right)^{1/p},$$

and the norm in  $X$  by  $\|u\|_X = \|u\|_p + \|u'\|_p$ . Note that for each  $p > 1$ ,  $X$  is compact embedded in  $C$ . Let  $B_r = \{x \in \mathbb{R}^N : |x| \leq r\}$ .

Each  $u \in L^1$  can be written as  $u(t) = \bar{u} + \tilde{u}(t)$  with

$$\bar{u} := \frac{1}{T} \int_0^T u(t) dt, \quad \int_0^T \tilde{u}(t) dt = 0.$$

We will use the Sobolev inequality

$$\|\tilde{u}\|_\infty \leq T^{1/q} \|u'\|_p \quad \text{for each } u \in X \text{ (here } q \text{ is the conjugate of } p)$$

and Wirtinger's inequality

$$\|\tilde{u}\|_p^p \leq T^p \|\tilde{u}'\|_p^p \quad \text{for each } u \in X.$$

The proof of the theorem is given in Section 3. In Section 2 we present some preliminary results on the variational setting of  $p$ -Laplacian equations in  $X$  and the related Palais-Smale compactness.

## 2. PRELIMINARIES

We first recall some facts about the eigenvalue problem for the  $p$ -Laplacian, see [9]. A  $\lambda \in \mathbb{R}$  is said to be an "eigenvalue" of the  $p$ -Laplacian with periodic boundary conditions, if the problem

$$\begin{cases} (|u'|^{p-2}u')' + \lambda|u|^{p-2}u = 0 & \text{a.e. on } [0, T], \\ u(0) - u(T) = u'(0) - u'(T) = 0 \end{cases}$$

has a nontrivial solution  $u \in C^1(I, \mathbb{R}^N)$ , known as the corresponding one to  $\lambda$  "eigenfunction". Let  $S$  denote the set of these eigenvalues. Evidently  $0 \in S$ , each element

of  $S$  is nonnegative and 0 is the smallest (first) eigenvalue. If  $N = 1$  (scalar case), by direct integration of the equation we see that all the eigenvalues are

$$\lambda_n = \left( \frac{2\pi_p n}{T} \right)^p, \quad n = 0, 1, 2, 3, \dots$$

where

$$\pi_p = 2(p-1)^{1/p} \int_0^1 (1-t^p)^{-1/p} dt.$$

In the case  $N > 1$  (vector case),  $\{\lambda_n\}_{n \geq 1} \subseteq S$  but  $S$  contains more elements.

It follows from assumption (A) that the functional  $\varphi$  on  $X$  given by

$$\varphi(u) = \frac{1}{p} \int_0^T |u'(t)|^p dt - \int_0^T F(t, u(t)) dt$$

is continuously differentiable and weakly lower semicontinuous on  $X$  (the proof is similar to the case  $p = 2$ , see [1]). Moreover, one has

$$\langle \varphi'(u), v \rangle = \int_0^T |u'(t)|^{p-2} (u'(t), v'(t)) dt - \int_0^T (\nabla F(t, u(t)), v(t)) dt$$

for all  $u, v \in X$ . It is easy to see that the solutions of problem (1.1) correspond to the critical points of  $\varphi$ .

Let us write

$$I(u) = \frac{1}{p} \int_0^T |u'|^p dt, \quad G(u) = \int_0^T F(t, u(t)) dt, \quad \forall u \in X.$$

**Proposition 2.1.** *The mapping  $I': X \rightarrow X^*$  is of type  $(S_+)$  (see [7]), i.e. any sequence  $\{u_n\}$  in  $X$  satisfying  $u_n \rightharpoonup u$  in  $X$  and*

$$\overline{\lim}_{n \rightarrow \infty} \langle I'(u_n), u_n - u \rangle \leq 0$$

*contains a convergent subsequence.*

*Proof.* Assume that  $u_n \rightharpoonup u$  in  $X$  and  $\overline{\lim}_{n \rightarrow \infty} \langle I'(u_n), u_n - u \rangle \leq 0$ . Then we get

$$\overline{\lim}_{n \rightarrow \infty} \langle I'(u_n) - I'(u), u_n - u \rangle \leq 0,$$

and together with the monotonicity property of  $I'$  this implies

$$\lim_{n \rightarrow \infty} \langle I'(u_n) - I'(u), u_n - u \rangle = 0,$$

i.e.,

$$(2.1) \quad \int_0^T \langle |u'_n|^{p-2} u'_n - |u'|^{p-2} u', u'_n - u' \rangle dt \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Recall that for all  $x, y \in \mathbb{R}^N$  the following inequalities hold:

$$\langle |x|^{p-2} x - |y|^{p-2} y, x - y \rangle \geq \left(\frac{1}{2}\right)^{p-1} |x - y|^p \quad \text{when } p \geq 2,$$

and

$$(|x| + |y|)^{2-p} \langle |x|^{p-2} x - |y|^{p-2} y, x - y \rangle \geq (p-1) |x - y|^2 \quad \text{when } 1 < p \leq 2.$$

Hence, by (2.1), one has that  $u'_n$  converges to  $u'$  in  $[0, T]$  in measure. Thus there exists a subsequence (without loss of generality assume it to be the whole sequence) with

$$u'_n(t) \rightarrow u'(t) \quad \text{for a.e. } t \in [0, T], \quad n \rightarrow \infty,$$

and so

$$(2.2) \quad \frac{1}{p} |u'_n(t) - u'(t)|^p \rightarrow 0 \quad \text{for a.e. } t \in [0, T], \quad n \rightarrow \infty.$$

One also has

$$\langle I'(u_n), u_n - u \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Moreover,

$$\begin{aligned} \langle I'(u_n), u_n - u \rangle &= \int_0^T |u'_n|^p dt - \int_0^T |u'_n|^{p-2} u'_n u' dt \\ &\geq \int_0^T |u'_n|^p dt - \int_0^T |u'_n|^{p-1} |u'| dt \\ &\geq \int_0^T |u'_n|^p dt - \int_0^T \left[ \frac{p-1}{p} |u'_n|^p + \frac{1}{p} |u'|^p \right] dt \\ &= \frac{1}{p} \int_0^T (|u'_n|^p - |u'|^p) dt. \end{aligned}$$

Now the lower semi-continuity of  $I$  yields

$$\liminf_{n \rightarrow \infty} \frac{1}{p} \int_0^T |u'_n|^p dt \geq \frac{1}{p} \int_0^T |u'|^p dt.$$

Thus

$$\frac{1}{p} \int_0^T |u'_n|^p dt \rightarrow \frac{1}{p} \int_0^T |u'|^p dt \quad \text{as } n \rightarrow \infty.$$

Consequently, the sequence  $\{p^{-1} \int_0^t |u'_n|^p dt\}$  is equi-absolutely continuous on  $[0, T]$ . From the inequality

$$\frac{1}{p} |u'_n(t) - u'(t)|^p \leq \frac{2^p}{p} (|u'_n|^p + |u'|^p)$$

we obtain that

$$(2.3) \quad \left\{ \frac{1}{p} \int_0^t |u'_n(t) - u'(t)|^p dt \right\} \text{ is equi-absolutely continuous on } [0, T].$$

Now (2.2), (2.3) and the convergence theorem of Vitali (see [7], p. 1017) guarantee that

$$\int_0^T \frac{1}{p} |u'_n(t) - u'(t)|^p dt \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus  $u'_n \rightarrow u'$  in  $L^p$ .

Also  $u_n \rightarrow u$  in  $X$ , which implies that  $u_n \rightarrow u$  in  $L^p$ . Hence  $u_n \rightarrow u$  in  $X$ , and the proof is complete.  $\square$

Since the sum of a mapping of type  $(S_+)$  with a weakly-strongly continuous mapping is still a mapping of type  $(S_+)$ , we obtain the following result.

**Proposition 2.2.**  $\varphi' : X \rightarrow X^*$  is a mapping of type  $(S_+)$ .

**Proposition 2.3.** Suppose that if a sequence  $\{u_n\}$  in  $X$  is such that

$$\{\varphi(u_n)\} \text{ is bounded and } \varphi'(u_n) \rightarrow 0$$

as  $n \rightarrow \infty$ , then  $\{u_n\}$  has a bounded subsequence. Then  $\varphi$  satisfies the PS-condition.

*Proof.* Assume that  $\{u_n\} \subset X$ ,  $\{\varphi(u_n)\}$  is bounded and  $\varphi'(u_n) \rightarrow 0$ . Now we know that  $\{u_n\}$  contains a bounded subsequence and for simplicity we denote it again by  $\{u_n\}$ . Since  $X$  is reflexive we can extract a subsequence (again denoting it by  $\{u_n\}$ ) such that  $u_n \rightarrow u$  in  $X$ . Since  $\varphi'(u_n) \rightarrow 0$ , one has

$$\langle \varphi'(u_n), u_n - u \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Since  $\varphi'$  is a mapping of type  $(S_+)$  we have that  $u_n \rightarrow u$  in  $X$ . Hence  $\varphi$  satisfies the PS-condition.  $\square$

### 3. PROOFS OF THEOREMS

We shall prove Theorem 1.1 from a sequence of claims.

Consider the direct sum decomposition  $X = \mathbb{R}^N \oplus V$  with  $V = \{v \in X : \int_0^T v(t) dt = 0\}$ . So for all  $u \in X$  we can write  $u = \bar{u} + \tilde{u}$  with  $\bar{u} \in \mathbb{R}^N$  and  $\tilde{u} \in V$ .

**Claim 1.**  $\varphi$  is coercive on  $V$ .

*Proof.* Combining hypotheses (H1) with the Mean Value Theorem, we see that for almost all  $t \in I$  and all  $x \in \mathbb{R}^N$ ,  $|F(t, x)| \leq g(t)|x|$ . If  $\tilde{u} \in V$ , then

$$\varphi(\tilde{u}) \geq \frac{1}{p} \|\tilde{u}'\|_p^p - \int_0^T g(t)|\tilde{u}(t)| dt \geq \frac{1}{p} \|\tilde{u}'\|_p^p - \|g\|_q \|\tilde{u}\|_p \geq \frac{1}{p} \|\tilde{u}'\|_p^p - T \|g\|_q \|\tilde{u}'\|_p$$

by Wirtinger's inequality (here  $q$  is the conjugate of  $p$ ). The above inequality and  $\tilde{u} \in V$  imply that  $\varphi(\tilde{u}) \rightarrow \infty$  as  $\|\tilde{u}'\|_p \rightarrow \infty$ . Now it follows that  $\varphi(\tilde{u}) \rightarrow \infty$ , as  $\|\tilde{u}\|_X \rightarrow 0$ .  $\square$

**Claim 2.** There exists a sequence  $(R_n)$  such that

$$\lim_{n \rightarrow \infty} R_n = +\infty, \quad \lim_{n \rightarrow \infty} \left[ \sup_{a \in \mathbb{R}^N, |a|=R_n} \varphi(a) \right] = -\infty.$$

*Proof.* This follows from (1.2) since

$$\varphi(a) = - \int_0^T F(t, a) dt.$$

$\square$

**Claim 3.** There exists a sequence  $(r_m)$  such that

$$\lim_{m \rightarrow \infty} r_m = +\infty, \quad \text{and} \quad \lim_{m \rightarrow \infty} \left[ \inf_{b \in \mathbb{R}^N, |b|=r_m, \tilde{u} \in V} \varphi(b + \tilde{u}) \right] = +\infty.$$

*Proof.* For any  $b \in \mathbb{R}^N$ ,  $|b| = r_m$ , and  $\tilde{u} \in V$ , let  $u = b + \tilde{u}$ . Note that

$$\begin{aligned} \varphi(u) &= \frac{1}{p} \|u'\|_p^p - \int_0^T F(t, u) dt \\ &= \frac{1}{p} \|\tilde{u}'\|_p^p - \int_0^T [F(t, u) - F(t, b)] dt - \int_0^T F(t, b) dt \\ &= \frac{1}{p} \|\tilde{u}'\|_p^p - \int_0^T \int_0^1 (\nabla F(t, b + s\tilde{u}(t)), \tilde{u}(t)) ds dt - \int_0^T F(t, b) dt \\ &\geq \frac{1}{p} \|\tilde{u}'\|_p^p - \left( \int_0^T g(t) dt \right) \|\tilde{u}\|_\infty - \int_0^T F(t, b) dt \\ &\geq \frac{1}{p} \|\tilde{u}'\|_p^p - C_1 \|\tilde{u}'\|_p - \int_0^T F(t, b) dt \end{aligned}$$



by Sobolev's inequality; here  $C_1 = (\int_0^T g(t) dt) T^{1/q}$ . Thus

$$\inf_{b \in \mathbb{R}^N, |b|=r_m, \tilde{u} \in V} \varphi(b + \tilde{u}) \geq \inf_{\tilde{u} \in V} \left( \frac{1}{p} \|\tilde{u}'\|_p^p - C_1 \|\tilde{u}'\|_p \right) + \inf_{b \in \mathbb{R}^N, |b|=r_m} \left( - \int_0^T F(t, b) dt \right).$$

On the other hand, there exists  $\beta \in \mathbb{R}$  such that  $\inf_{\tilde{u} \in V} (p^{-1} \|\tilde{u}'\|_p^p - C_1 \|\tilde{u}'\|_p) \geq \beta$ . The claim now follows from (1.3).  $\square$

Consider now the set

$$S_n = \{ \gamma \in C(B_{R_n}, X), \gamma|_{\partial B_{R_n}} = i|_{\partial B_{R_n}} \}$$

and define

$$(3.1) \quad c_n = \inf_{\gamma \in S_n} \left[ \max_{x \in B_{R_n}} \varphi(\gamma(x)) \right].$$

We prove that each  $\gamma$  intersects the hyperplane  $V$ . Let  $\pi: X \rightarrow \mathbb{R}^N$  be the (continuous) projection of  $X$  onto  $\mathbb{R}^N$ , defined by

$$\pi(u) = \frac{1}{T} \int_0^T u dt \quad \text{for } u \in X.$$

Let  $\gamma$  be any continuous map such that  $\gamma|_{\partial B_{R_n}} = i|_{\partial B_{R_n}}$ . We have to show that  $0 \in \pi(\gamma(B_{R_n}))$ .

For  $t \in [0, 1]$ ,  $u \in \mathbb{R}^N$  define

$$\gamma_t(u) = t\pi(\gamma(u)) + (1-t)u.$$

Note that  $\gamma_t \in C^0(\mathbb{R}^N; \mathbb{R}^N)$  defines a homotopy of  $\gamma_0 = \text{id}$  with  $\gamma_1 = \pi \circ \gamma$ . Moreover,  $\gamma_t|_{\partial B_{R_n}} = \text{id}$  for all  $t$ . By homotopy invariance and normalization of the degree (see for instance Deimling [10, Theorem 1.3.1]) we have

$$\deg(\pi \circ \gamma, B_{R_n}, 0) = \deg(\text{id}, B_{R_n}, 0) = 1.$$

Hence  $0 \in \pi(\gamma(B_{R_n}))$ . Thus each  $\gamma$  intersects the hyperplane  $V$ .

Now since  $\varphi$  is coercive on  $V$ , there is a constant  $K$  such that

$$\max_{x \in B_{R_n}} \varphi(\gamma(x)) \geq \inf_{\tilde{u} \in V} \varphi(\tilde{u}) \geq K.$$

Hence  $c_n \geq K$  and, for all large values of  $n$ ,

$$c_n > \sup_{a \in \mathbb{R}^N, |a|=R_n} \varphi(a).$$

Let us now fix such  $n$  and apply Theorem 4.3 in [1]. This proves the next claim (see also [1, Corollary 4.3]).

**Claim 4.** *If  $n$  is large enough there exist sequences  $(\gamma_k)$  in  $S_n$  and  $(v_k)$  in  $X$  such that*

$$\max_{x \in B_{R_n}} \varphi(\gamma_k(x)) \rightarrow c_n \quad (k \rightarrow \infty).$$

*Then there exists a sequence  $(v_k)$  in  $X$  such that*

$$\varphi(v_k) \rightarrow c_n, \quad \text{dist}(v_k, \gamma_k(B_{R_n})) \rightarrow 0, \quad |\varphi'(v_k)| \rightarrow 0$$

*when  $k \rightarrow \infty$ .*

**Claim 5.** *The sequence  $(v_k)$  is bounded in  $X$ .*

*Proof.* For any  $k$  large enough,

$$c_n \leq \max_{x \in B_{R_n}} \varphi(\gamma_k(x)) \leq c_n + 1$$

and we can find  $w_k \in \gamma_k(B_{R_n})$  such that

$$(3.2) \quad \|v_k - w_k\|_X \leq 1.$$

Using Claim 3, for a fixed  $n \in \mathbb{N}$  there exists  $m \in \mathbb{N}$  such that  $r_m > R_n$ . As in the proof above, let  $\pi: X \rightarrow \mathbb{R}^N$  be the (continuous) projection of  $X$  onto  $\mathbb{R}^N$ . Then  $|\pi(\gamma_k(B_{R_n}))| \leq R_n$ . Let  $H_{r_m} = \{b \in \mathbb{R}^N: |b| = r_m\} + V$ . Then  $|\pi(H_{r_m})| = r_m$ , so  $\gamma_k(B_{R_n})$  cannot intersect the hyperplanes  $H_{r_m}$ . Hence, if we write  $w_k = \bar{w}_k + \tilde{w}_k$ , where  $\bar{w}_k \in \gamma_k(B_{R_n})$  and  $\tilde{w}_k \in V$ , we have

$$|\bar{w}_k| < r_m.$$

We also have

$$\begin{aligned}
c_n + 1 &\geq \varphi(w_k) = \frac{1}{p} \|(\bar{w}_k + \tilde{w}_k)'\|_p^p - \int_0^T F(t, w_k(t)) \, dt \\
&= \frac{1}{p} \|\tilde{w}_k'\|_p^p - \int_0^T [F(t, w_k(t)) - F(t, \bar{w}_k)] \, dt - \int_0^T F(t, \bar{w}_k) \, dt \\
&= \frac{1}{p} \|\tilde{w}_k'\|_p^p - \int_0^T \int_0^1 (\nabla F(t, \bar{w}_k + s\tilde{w}_k(t)), \tilde{w}_k(t)) \, ds \, dt - \int_0^T F(t, \bar{w}_k) \, dt \\
&\geq \frac{1}{p} \|\tilde{w}_k'\|_p^p - \left( \int_0^T g(t) \, dt \right) \|\tilde{w}_k\|_\infty - \int_0^T F(t, \bar{w}_k) \, dt \\
&\geq \frac{1}{p} \|\tilde{w}_k'\|_p^p - C_2 \|\tilde{w}_k'\|_p - \int_0^T F(t, \bar{w}_k) \, dt
\end{aligned}$$

by Sobolev's inequality; here  $C_2 = \left(\int_0^T g(t) \, dt\right) T^{1/q}$ . Notice also that since  $|\bar{w}_k| < r_m$ , the integral  $\int_0^T F(t, \bar{w}_k) \, dt$  is bounded. Consequently,  $\tilde{w}_k$  is bounded in  $X$ . Thus the sequence  $w_k$  is bounded and the claim follows from 3.2.  $\square$

**Claim 6.**  $c_n$  is a critical value.

*Proof.* From Proposition 2.3 and the last claim (recall also Claim 4) it follows that  $(v_k)$  contains a convergent subsequence, which we rename as  $(v_k)$ . Let  $u_n = \lim_{k \rightarrow \infty} v_k$ , then (see Claim 4)

$$\varphi'(u_n) = \lim_{k \rightarrow \infty} \varphi'(v_k) = 0 \quad \text{and} \quad \varphi(u_n) = \lim_{k \rightarrow \infty} \varphi(v_k) = c_n.$$

$\square$

*Proof of Theorem 1.1.* (a) Claim 6 proves that, for each  $n$  large enough, there exists at least one solution  $u_n$  of (1.1) such that  $\varphi(u_n) = c_n$ , where  $c_n$  is given by (3.1). If  $0 < r_k \leq R_n$  and  $H_{r_k} = \{b \in \mathbb{R}^N : |b| = r_k\} + V$ , then for any  $\gamma \in S_n$  we have that  $\gamma$  intersects the hyperplane  $H_{r_k}$  (the proof is similar to the above). Then

$$\max_{x \in B_{R_n}} \varphi(\gamma(x)) \geq \inf_{b \in \mathbb{R}^N, |b|=r_k, \tilde{u} \in V} \varphi(b + \tilde{u}).$$

Thus, using Claim 3, we obtain that

$$\lim_{n \rightarrow \infty} c_n = +\infty,$$

and (a) follows.

(b) For  $n \in \{1, 2, \dots\}$ , define a subset  $P_n$  of  $X$  by

$$P_n = \{u \in X : u = \bar{u} + \tilde{u}, \bar{u} \in \mathbb{R}^N, |\bar{u}| \leq r_n, \tilde{u} \in V\}.$$

We note that for  $u \in P_n$  we have, proceeding as in Claim 3,

$$(3.3) \quad \begin{aligned} \varphi(u) &= \frac{1}{p} \|u'\|_p^p - \int_0^T F(t, u) dt \\ &\geq \frac{1}{p} \|\tilde{u}'\|_p^p - C_1 \|\tilde{u}'\|_p - \int_0^T F(t, \bar{u}) dt \end{aligned}$$

by Sobolev's inequality; here  $C_1 = (\int_0^T g(t) dt) T^{1/q}$ . Notice also that  $\int_0^T F(t, \bar{u}) dt$  is bounded. Thus  $\varphi$  is bounded below on  $P_n$ .

Let us set

$$(3.4) \quad \mu_n = \inf_{u \in P_n} \varphi(u)$$

and let  $(u_k)$  be a sequence in  $P_n$  such that

$$(3.5) \quad \varphi(u_k) \rightarrow \mu_n \quad \text{as } k \rightarrow \infty.$$

We have

$$u_k = \bar{u}_k + \tilde{u}_k, \quad \bar{u}_k \in \mathbb{R}^N \quad \text{and} \quad |\bar{u}_k| \leq r_n.$$

Without loss of generality we assume that  $\bar{u}_k \rightarrow \bar{u} \in B_{r_n}$ . From (3.3) and (3.5) we obtain that  $(u_k)$  is a bounded sequence in  $X$  and thus passing to a subsequence, which we rename  $(u_k)$ , we have that

$$u_k \rightharpoonup u_n^* \quad \text{in } X.$$

Since  $P_n$  is a convex closed subset of  $X$ ,  $u_n^* \in P_n$ .

Note that  $\varphi$  is weakly lower semi-continuous, so

$$\mu_n = \lim_{k \rightarrow \infty} \varphi(u_k) \geq \varphi(u_n^*)$$

and, since  $u_n^* \in P_n$ , we must have

$$\mu_n = \varphi(u_n^*).$$

Next we want to show that  $u_n^* \in \text{Int } P_n$  for large  $n$ , where

$$\text{Int } P_n := \{u \in X : u = \bar{u} + \tilde{u}, |\bar{u}| < r_n\}.$$

Indeed, taking

$$0 < R_n < r_n,$$

it follows from Claim 2 and Claim 3 that  $u_n^* \notin \{b + \tilde{u}, b \in Y, |b| = r_n\}$  for large  $n$  and hence  $u_n^* \in \text{Int } P_n$ . This fact and (3.4) imply that

$$\varphi'(u_n^*) = 0$$

and  $u_n^*$  is a solution of (1.1).

Finally, from the fact that

$$\varphi(u_n^*) \leq \inf_{a \in \mathbb{R}^N, |a|=R_n} \varphi(a)$$

and from Claim 2 we obtain that

$$\lim_{n \rightarrow \infty} \varphi(u_n^*) = -\infty.$$

□

**Remark 1.** Going through the above proof, it is easy to see that the existence of critical points  $u_n$  still holds if assumption (1.2) is weakened to

$$\limsup_{R \rightarrow +\infty} \left[ \inf_{a \in \mathbb{R}^N, |a|=R} \int_0^T F(s, a) \, ds \right] \text{ is bounded from below.}$$

In that case, however, the existence of the minima  $u_n^*$  is no longer guaranteed.

**Example 1.** Let us consider the scalar problem

$$(3.6) \quad \begin{aligned} -(|u'|^{p-2}u')' &= a \sin(\ln(u^2 + 1)) + \frac{2au^2}{1+u^2} \cos(\ln(u^2 + 1)) + e(t), \\ u(0) - u(T) &= u'(0) - u'(T) = 0, \end{aligned}$$

where  $1 < p < \infty$ ,  $a > 0$ ,  $e \in L^1(0, T)$  and  $\int_0^T e(t) \, dt = 0$ . In this case

$$F(t, u) = au \sin(\ln(u^2 + 1)) + e(t)u$$

and hence

$$\int_0^T F(t, x) \, dt = ax \sin(\ln(x^2 + 1)).$$

Let  $R_k = \sqrt{e^{2k\pi+1/2\pi} - 1}$ ,  $r_k = R_k = \sqrt{e^{2k\pi+3/2\pi} - 1}$  for  $k = 1, 2, \dots$ . Then

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_0^T F(t, R_k) dt &= \lim_{k \rightarrow +\infty} aR_k \sin(\ln(R_k^2 + 1)) \\ &= \lim_{k \rightarrow +\infty} a\sqrt{e^{2k\pi+1/2\pi} - 1} = +\infty \end{aligned}$$

and

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_0^T F(t, r_k) dt &= \lim_{k \rightarrow +\infty} ar_k \sin(\ln(r_k^2 + 1)) \\ &= \lim_{k \rightarrow +\infty} -a\sqrt{e^{2k\pi+3/2\pi} - 1} = -\infty. \end{aligned}$$

As a result, (1.2) and (1.3) are satisfied. Moreover,  $|F'_u(t, u)|$  is clearly bounded by  $3a + |e(t)|$  and so we have the following result:

- (i) there exists a sequence  $(u_n)$  of solutions of (3.6) such that  $u_n$  is a critical point of  $\varphi$  and  $\lim_{n \rightarrow \infty} \varphi(u_n) = +\infty$ ;
- (ii) there exists a sequence  $(u_n^*)$  of solutions of (3.6) such that  $u_n^*$  is a local minimum point of  $\varphi$  and  $\lim_{n \rightarrow \infty} \varphi(u_n^*) = -\infty$ .

### References

- [1] *J. Mawhin, M. Willem: Critical Point Theory and Hamiltonian Systems.* Springer-Verlag, New York-Berlin-Heidelberg-London-Paris-Tokyo, 1989.
- [2] *J. Mawhin: Some boundary value problems for Hartman-type perturbations of the ordinary vector  $p$ -Laplacian.* *Nonlinear Anal., Theory Methods Appl.* 40A (2000), 497–503.
- [3] *M. Del Pino, R. Manasevich, and A. Murua: Existence and multiplicity of solutions with prescribed period for a second order O.D.E.* *Nonlinear Anal., Theory Methods Appl.* 18 (1992), 79–92.
- [4] *C. Fabry, D. Fayyad: Periodic solutions of second order differential equations with a  $p$ -Laplacian and asymmetric nonlinearities.* *Rend. Ist. Mat. Univ. Trieste* 24 (1992), 207–227.
- [5] *Z. Guo: Boundary value problems of a class of quasilinear ordinary differential equations.* *Differ. Integral Equ.* 6 (1993), 705–719.
- [6] *H. Dang, S. F. Oppenheimer: Existence and uniqueness results for some nonlinear boundary value problems.* *J. Math. Anal. Appl.* 198 (1996), 35–48.
- [7] *E. Zeidler: Nonlinear Functional Analysis and its Applications. II/B: Nonlinear Monotone Operators.* Springer-Verlag, New York-Berlin-Heidelberg, 1990.
- [8] *P. Habets, R. Manasevich, and F. Zanolin: A nonlinear boundary value problem with potential oscillating around the first eigenvalue.* *J. Differ. Equations* 117(1995), 428–445.
- [9] *J. Mawhin: Periodic solutions of systems with  $p$ -Laplacian-like operators.* In: *Nonlinear Analysis and Applications to Differential Equations. Papers from the Autumn School on Nonlinear Analysis and Differential Equations, Lisbon, September 14–October 23, 1998.* *Progress in Nonlinear Differential Equations and Applications.* Birkhäuser-Verlag, Boston, 1998, pp. 37–63.

- [10] *K. Deimling*: Nonlinear Functional Analysis. Springer-Verlag, Berlin-Heidelberg-New York-Tokyo, 1985.

*Authors' addresses:* *H. Lü*, Department of Applied Mathematics, Hohai University, Nanjing, 210098, China, e-mail: [haishen2001@yahoo.com.cn](mailto:haishen2001@yahoo.com.cn); *D. O'Regan*, Department of Mathematics, National University of Ireland, Galway, Ireland, e-mail: [Donal.ORegan@nuigalway.ie](mailto:Donal.ORegan@nuigalway.ie); *R.P. Agarwal*, Department of Mathematical Sciences, Florida Institute of Technology, Melbourne, Florida 32901, USA, e-mail: [agarwal@fit.edu](mailto:agarwal@fit.edu).