

Zhendong Gu; Daochun Sun

The growth of a class of random Dirichlet series on the horizontal zone

*Mathematica Bohemica*, Vol. 137 (2012), No. 1, 65–77

Persistent URL: <http://dml.cz/dmlcz/142788>

## Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE GROWTH OF A CLASS OF RANDOM DIRICHLET SERIES  
ON THE HORIZONTAL ZONE

ZHENDONG GU, DAOCHUN SUN, Guangzhou

(Received August 9, 2010)

*Abstract.* In the paper we obtain that, under some condition, the Rademacher-Dirichlet series or the Steinhaus-Dirichlet series on the whole plane and on the horizontal zone almost surely have the same growth.

*Keywords:* random Dirichlet series, Rademacher-Dirichlet series, Steinhaus-Dirichlet series, growth

*MSC 2010:* 30B20

1. INTRODUCTION AND MAIN RESULT

Let  $(\Omega, \mathcal{A}, P)$  be a probability space, where  $\Omega = [0, 1]$ ,  $\mathcal{A}$  is composed of all Lebesgue measurable sets  $E$  on  $\Omega$  and  $P(E)$  is the Lebesgue measure of  $E$ . Let  $\{\varepsilon_n(\omega)\}$  and  $\{\gamma_n(\omega)\}$  ( $n = 0, 1, 2, \dots$ ) (see [1], [2]), which can be considered to be random variables sequences in  $(\Omega, \mathcal{A}, P)$ , be respectively the Rademacher and Steinhaus sequence.  $\gamma_n(\omega) = \exp\{2\pi i\theta_n(\omega)\}$ , the value of  $\theta_n(\omega)$  uniformly distributes on  $[0, 1]$ .  $\{\varepsilon_n(\omega)\}$  ( $n = 0, 1, 2, \dots$ ) satisfies

$$P[\varepsilon_n(\omega) = 1] = P[\varepsilon_n(\omega) = -1] = \frac{1}{2}.$$

Consider the random Dirichlet series

$$f(s, \omega) = \sum_{n=0}^{+\infty} a_n X_n(\omega) e^{-\lambda_n s},$$

---

This work is supported by the National Natural Science Foundation of China (No. 11101096).

where  $X_n(\omega) = \varepsilon_n(\omega)$  or  $\gamma_n(\omega)$ ,  $s = \sigma + it$  ( $\sigma, t \in \mathbb{R}$ ) denotes the complex variable,  $\{a_n\}_{n=0}^{+\infty}$  is a sequence of complex numbers,  $0 = \lambda_0 < \lambda_1 < \lambda_2 < \dots < \lambda_n \uparrow +\infty$ . Following Bohr (see [3]), we define the quantities

$$\begin{aligned}\sigma_c(\omega) &= \inf \left\{ \sigma \in \mathbb{R} : \sum a_n X_n(\omega) e^{-\lambda_n \sigma} \text{ converges} \right\}, \\ \sigma_a(\omega) &= \inf \left\{ \sigma \in \mathbb{R} : \sum |a_n X_n(\omega)| e^{-\lambda_n \sigma} \text{ converges} \right\}, \\ \sigma_u(\omega) &= \inf \left\{ \sigma \in \mathbb{R} : \sum a_n X_n(\omega) e^{-\lambda_n(\sigma+it)} \text{ converges uniformly on } \mathbb{R} \right\}.\end{aligned}$$

Paley and Zygmund (see [3], 1932) were the first to study the convergence of  $f(s, \omega)$ . Yu Jiarong (see [4], 1978) studied the growth of  $f(s, \omega)$  under the conditions

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n} = 0, \quad \text{and} \quad \overline{\lim}_{n \rightarrow +\infty} \frac{n}{\lambda_n} < +\infty,$$

and obtained that  $f(s, \omega)$  on the right half plane and on any right horizontal zone a.s. (almost surely) has the same growth. In this paper, we discuss Yu's topic on the whole plane. We obtain that, under some condition,  $f(s, \omega)$  on the whole plane and on any horizontal zone a.s. has the same growth.

Let  $\sigma_u(\omega)$  be the uniformly convergence abscissa of  $f(s, \omega)$ . When  $\sigma_u(\omega) < +\infty$ , for any  $\sigma > \sigma_u(\omega)$  let

$$\begin{aligned}M(\sigma, \omega) &= \sup\{|f(s, \omega)| : -\infty < t < +\infty\}, \\ M(\sigma; \alpha, \beta; \omega) &= \sup\{|f(s, \omega)| : -\infty < \alpha \leq t \leq \beta < +\infty\}.\end{aligned}$$

Put

$$\Delta = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln(p_k + 1)}{k \ln k},$$

where  $p_k$  is given by  $[k, k+1) \cap \{\lambda_n\} = \{\lambda_{n_k}, \lambda_{n_k+1}, \dots, \lambda_{n_k+p_k}\}$ ,  $k \in \mathbb{N}$ . Our main result is

**Theorem 1.** *Assume that  $f(s, \omega)$  satisfies*

$$\sigma_u(\omega) = -\infty \text{ a.s. and } \Delta = 0.$$

*Then  $f(s, \omega)$  a.s. satisfies*

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma, \omega)}{-\sigma} = \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma; \alpha, \beta; \omega)}{-\sigma}.$$

## 2. LEMMAS

To give our lemmas, we define some symbols by the method of Knoop-Kojima (see [5]). Consider

$$f(s) = \sum_{n=0}^{+\infty} a_n e^{-\lambda_n s}.$$

For each  $k \in \mathbb{N}$ , when

$$(1) \quad [k, k+1) \cap \{\lambda_n\} = \{\lambda_{n_k}, \lambda_{n_k+1}, \dots, \lambda_{n_k+p_k}\} \neq \emptyset,$$

put

$$\bar{A}_k = \sup_{0 \leq p \leq p_k, t \in \mathbb{R}} \left| \sum_{j=0}^p a_{n_k+j} e^{-it\lambda_{n_k+j}} \right|, \quad A_k^* = \sum_{j=0}^{p_k} |a_{n_k+j}|;$$

when  $[k, k+1) \cap \{\lambda_n\} = \emptyset$ , put  $\ln \bar{A}_k = \ln A_k^* = -\infty$ . Then we have the formulas (see [5], [6]) for the abscissas  $\sigma_u, \sigma_a$  in terms of  $\bar{A}_k, A_k^*$ ,

$$\sigma_u = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln \bar{A}_k}{k}; \quad \sigma_a = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln A_k^*}{k}.$$

When  $\sigma_u < +\infty$ , for any  $\sigma > \sigma_u$  put

$$\bar{M}_u(\sigma) = \sup \left\{ \left| \sum_{j=0}^n a_j e^{-\lambda_j(\sigma+it)} \right| : n \in \mathbb{N}, t \in \mathbb{R} \right\}; \quad \bar{m}(\sigma) = \max\{\bar{A}_k e^{-k\sigma} : k \in \mathbb{N}\}.$$

Consider  $f(s, \omega)$ . Put

$$\bar{A}_k(\omega) = \sup_{0 \leq p \leq p_k, t \in \mathbb{R}} \left| \sum_{j=0}^p a_{n_k+j} X_j(\omega) e^{-it\lambda_{n_k+j}} \right|;$$

$$\bar{M}_u(\sigma, \omega) = \sup \left\{ \left| \sum_{j=0}^n a_j X_j(\omega) e^{-\lambda_j(\sigma+it)} \right| : n \in \mathbb{N}, -\infty < t < +\infty \right\}.$$

Chuji Tanaka (see [7]) studied the relation between  $\bar{m}(\sigma)$  and  $\bar{M}_u(\sigma)$ . Now we improve his results.

**Lemma 1.** *If  $f(s)$  is uniformly convergent on the whole plane, then*

- (I)  $\frac{\overline{m}(\sigma)}{4e^{|\sigma|}} \leq \overline{M}_u(\sigma) \leq \frac{2e^{|\sigma|\overline{m}(\sigma - \varepsilon)}}{1 - e^{-\varepsilon}}$  for any  $\varepsilon > 0$ ,
- (II)  $\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln \overline{M}_u(\sigma)}{-\sigma} = \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln \overline{m}(\sigma)}{-\sigma}$ .

*Proof.* Take  $p \in \mathbb{N}$ , such that  $n_k + p < n_{k+1}$ , where  $n_k$  is defined by (1). Using Abel's transformation, we arrive at

$$\begin{aligned} \sum_{j=n_k}^{n_k+p} a_j e^{-it\lambda_j} &= \sum_{j=n_k}^{n_k+p} a_j e^{-(\sigma+it)\lambda_j} e^{\sigma\lambda_j} \\ &= \sum_{j=n_k}^{n_k+p-1} \left( \sum_{q=n_k}^j a_q e^{-(\sigma+it)\lambda_q} \right) (e^{\sigma\lambda_j} - e^{\sigma\lambda_{j+1}}) \\ &\quad + \left( \sum_{q=n_k}^{n_k+p} a_q e^{-(\sigma+it)\lambda_q} \right) e^{\sigma\lambda_{n_k+p}}. \end{aligned}$$

Noting that

$$\begin{aligned} \left| \sum_{q=n_k}^j a_q e^{-(\sigma+it)\lambda_q} \right| &= \left| \sum_{q=0}^j a_q e^{-(\sigma+it)\lambda_q} - \sum_{q=0}^{n_k-1} a_q e^{-(\sigma+it)\lambda_q} \right| \\ &\leq \left| \sum_{q=0}^j a_q e^{-(\sigma+it)\lambda_q} \right| + \left| \sum_{q=0}^{n_k-1} a_q e^{-(\sigma+it)\lambda_q} \right| \leq 2\overline{M}_u(\sigma), \end{aligned}$$

we obtain

$$\begin{aligned} \overline{A}_k &\leq 2\overline{M}_u(\sigma) |e^{\sigma\lambda_{n_k}} - e^{\sigma\lambda_{n_k+p}}| + e^{\sigma\lambda_{n_k+p}} 2\overline{M}_u(\sigma) \\ &\leq 4\overline{M}_u(\sigma) e^{(k+\text{sgn}\sigma)\sigma} = 4e^{|\sigma|} \overline{M}_u(\sigma) e^{k\sigma}. \end{aligned}$$

By the definition of  $\overline{m}(\sigma)$ ,

$$(2) \quad \frac{\overline{m}(\sigma)}{4e^{|\sigma|}} \leq \overline{M}_u(\sigma).$$

Suppose  $n_k + p < n_{k+1}$ . Abel's transformation yields

$$\begin{aligned} \sum_{j=n_k}^{n_k+p} a_j e^{-(\sigma+it)\lambda_j} &= \sum_{j=n_k}^{n_k+p} a_j e^{-it\lambda_j} e^{-\sigma\lambda_j} \\ &= \sum_{j=n_k}^{n_k+p-1} \left( \sum_{q=n_k}^j a_q e^{-it\lambda_q} \right) (e^{-\sigma\lambda_j} - e^{-\sigma\lambda_{j+1}}) \\ &\quad + \left( \sum_{q=n_k}^{n_k+p} a_q e^{-it\lambda_q} \right) e^{-\sigma\lambda_{n_k+p}}. \end{aligned}$$

When  $\sigma > 0$ ,

$$\begin{aligned} \left| \sum_{j=n_k}^{n_k+p} a_j e^{-(\sigma+it)\lambda_j} \right| &\leq \bar{A}_k \sum_{j=n_k}^{n_k+p-1} (e^{-\sigma\lambda_j} - e^{-\sigma\lambda_{j+1}}) + \bar{A}_k e^{-\sigma\lambda_{n_k+p}} \\ &\leq \bar{A}_k e^{-\sigma\lambda_{n_k}} \leq \bar{A}_k e^{-\sigma k}. \end{aligned}$$

When  $\sigma \leq 0$ ,

$$\left| \sum_{j=n_k}^{n_k+p} a_j e^{-(\sigma+it)\lambda_j} \right| \leq \bar{A}_k \sum_{j=n_k}^{n_k+p-1} (e^{-\sigma\lambda_{j+1}} - e^{-\sigma\lambda_j}) + \bar{A}_k e^{-\sigma\lambda_{n_k+p}} \leq 2\bar{A}_k e^{-\sigma(k+1)}.$$

Then for any  $\sigma \in \mathbb{R}$  we have

$$\left| \sum_{j=n_k}^{n_k+p} a_j e^{-(\sigma+it)\lambda_j} \right| \leq 2\bar{A}_k e^{|\sigma|} e^{-k\sigma}.$$

Therefore for any  $\varepsilon > 0$

$$\begin{aligned} \left| \sum_{j=0}^{n_k+p} a_j e^{-(\sigma+it)\lambda_j} \right| &\leq 2e^{|\sigma|} \sum_{j=0}^k \bar{A}_j e^{-(\sigma-\varepsilon)j} e^{-j\varepsilon} \\ &\leq 2e^{|\sigma|} \bar{m}(\sigma - \varepsilon) \sum_{j=0}^{+\infty} e^{-j\varepsilon} = \frac{2e^{|\sigma|} \bar{m}(\sigma - \varepsilon)}{1 - e^{-\varepsilon}}. \end{aligned}$$

By the definition of  $\bar{M}_u(\sigma)$ ,

$$(3) \quad \bar{M}_u(\sigma) \leq \frac{2e^{|\sigma|} \bar{m}(\sigma - \varepsilon)}{1 - e^{-\varepsilon}}.$$

Combining (2) and (3), we prove (I).

Note that

$$\ln^+ \ln^+ \bar{m}(\sigma) \leq \ln^+ \ln^+ \frac{1}{4} e^{-|\sigma|} \bar{m}(\sigma) + \ln^+ \ln^+ 4e^{|\sigma|} + \ln 2, \text{ and } \overline{\lim}_{\sigma \rightarrow -\infty} \ln \bar{m}(\sigma) = +\infty.$$

Hence

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln \bar{m}(\sigma)}{-\sigma} = \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln^+ \ln^+ \bar{m}(\sigma)}{-\sigma} \leq \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln^+ \ln^+ \frac{1}{4} e^{-|\sigma|} \bar{m}(\sigma)}{-\sigma}.$$

Similarly,

$$\begin{aligned}
& \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln^+ \ln^+ 2e^{|\sigma| \overline{m}(\sigma - \varepsilon)} / (1 - e^{-\varepsilon})}{-\sigma} \\
& \leq \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln^+ \ln^+ \overline{m}(\sigma - \varepsilon)}{-\sigma} + \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln^+ \ln^+ 2e^{|\sigma|} / (1 - e^{-\varepsilon})}{-\sigma} \\
& = \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln^+ \ln^+ \overline{m}(\sigma)}{-\sigma}.
\end{aligned}$$

Thus (II) is proved.

**Lemma 2.** *Suppose that  $M_x < e^{-c\sigma} + x\sigma$  holds for any  $\sigma < \sigma_\varepsilon$  and any  $x \in \mathbb{R}$ ,  $x > 0$ , where  $c$  is a positive real number,  $\sigma_\varepsilon$  is a given real number,  $M_x$  is a real number depending on  $x$ . Then there exists  $x_\varepsilon \in \mathbb{R}$ ,  $x_\varepsilon > 0$  such that for any  $x > x_\varepsilon$ ,*

$$M_x < \min\{e^{-c\sigma} + x\sigma : \sigma \in \mathbb{R}\} = \frac{x}{c}(\ln ce - \ln x).$$

*Proof.* Let  $\psi(\sigma) = e^{-c\sigma} + x\sigma$  ( $-\infty < \sigma < +\infty$ ). Then there exists  $\sigma_x = (\ln c - \ln x)/c$  such that  $\min\{\psi(\sigma) : \sigma \in \mathbb{R}\} = \psi(\sigma_x) = xc^{-1}(\ln ce - \ln x)$ . Note that  $\sigma_x$  is a monotonic decreasing function of  $x$ , and  $\sigma_x \rightarrow -\infty \Leftrightarrow x \rightarrow +\infty$ . Then there exists  $x_\varepsilon \in \mathbb{R}$  such that  $\sigma_{x_\varepsilon} < \sigma_\varepsilon$ . Hence, for any  $x > x_\varepsilon$ , we have  $\sigma_x < \sigma_\varepsilon$ . Therefore  $M_x < \psi(\sigma_x) = xc^{-1}(\ln ce - \ln x)$ ,  $x > x_\varepsilon$ .

**Lemma 3.** *If  $f(s)$  is uniformly convergent on the whole plane, then*

$$(4) \quad \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln \overline{M}_u(\sigma)}{-\sigma} = \varrho_u \Leftrightarrow \overline{\lim}_{k \rightarrow +\infty} \frac{\ln \overline{A}_k}{k \ln k} = \begin{cases} -\infty, & \varrho = 0; \\ -\frac{1}{\varrho_u}, & 0 < \varrho < +\infty; \\ 0, & \varrho = +\infty. \end{cases}$$

*Proof.* We prove the necessity of the right hand side condition of (4). Consider the case  $0 < \varrho_u < \infty$ . Using Lemma 1(II), for  $\forall \varepsilon > 0$  we have, when  $-\sigma$  is large enough,

$$\overline{m}(\sigma) < \exp\{e^{-(\varrho_u + \varepsilon)\sigma}\}.$$

Then for each  $k \in \mathbb{N}$ ,

$$\ln \overline{A}_k < e^{-(\varrho_u + \varepsilon)\sigma} + k\sigma.$$

Note that

$$\min\{e^{-(\varrho_u + \varepsilon)\sigma} + k\sigma : \sigma \in \mathbb{R}\} = -\frac{k}{\varrho_u + \varepsilon} \ln \frac{k}{e(\varrho_u + \varepsilon)}.$$

By Lemma 2, when  $k$  is sufficiently large,

$$\ln \bar{A}_k < -\frac{k}{\varrho_u + \varepsilon} \ln \frac{k}{e(\varrho_u + \varepsilon)}.$$

As  $\varepsilon \rightarrow 0$ ,

$$\overline{\lim}_{k \rightarrow +\infty} \frac{\ln \bar{A}_k}{k \ln k} \leq -\frac{1}{\varrho_u}.$$

Suppose  $\overline{\lim}_{k \rightarrow +\infty} (\ln \bar{A}_k)/k \ln k < -1/\varrho_u$ . Then there exists  $\varrho'_u \in (0, \varrho_u)$ ,  $c > 0$  such that for each  $k \in \mathbb{N}$  and for any  $\sigma < 0$ ,

$$\bar{A}_k e^{-k\sigma} < \exp \left\{ -\frac{k \ln k}{\varrho'_u} - k\sigma + c \right\}.$$

Note that

$$\max \left\{ -\frac{k \ln k}{\varrho'_u} - k\sigma : k \in \mathbb{N} \right\} \leq \max \left\{ -\frac{k \ln k}{\varrho'_u} - k\sigma : k > 0, k \in \mathbb{R} \right\} = e^{-\varrho'_u \sigma - \ln(\varrho'_u) - 1}.$$

Then

$$\overline{m}(\sigma) < \exp \{ e^{-\varrho'_u \sigma - \ln(\varrho'_u) - 1} + c \}.$$

Applying Lemma 1(II), we conclude

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln \overline{M}_u(\sigma)}{-\sigma} \leq \varrho'_u < \varrho_u,$$

which contradicts the left-hand side of (4). Thus the necessity of the right-hand side of (4) is proved. By the above proof, we can easily prove the sufficiency of the right-hand side of (4).

Using the conclusion of the case  $0 < \varrho_u < \infty$ , we can easily prove the case  $\varrho_u = 0$ ,  $\varrho_u = +\infty$ . Thus the lemma is proved.

**Lemma 4.** *If  $f(s)$  is uniformly convergent on the whole plane and  $\Delta = 0$ , then*

$$(5) \quad \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma)}{-\sigma} = \varrho \Leftrightarrow \overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n \ln \lambda_n} = \begin{cases} -\infty, & \varrho = 0; \\ -\frac{1}{\varrho}, & 0 < \varrho < +\infty; \\ 0, & \varrho = +\infty. \end{cases}$$

**Proof.** For any  $\varepsilon > 0$ , when  $-\sigma$  is large enough,

$$M(\sigma) < \exp \{ e^{-(\varrho+\varepsilon)\sigma} \}.$$



Taking into account Hadamard's theorem (see [8]),  $a_n e^{-\lambda_n \sigma} \leq M(\sigma)$ ,

$$|a_n| < \exp\{e^{-(\varrho+\varepsilon)\sigma} + \lambda_n \sigma\}, \quad \forall n \in \mathbb{N}.$$

Note that

$$\min\{e^{-(\varrho+\varepsilon)\sigma} + \lambda_n \sigma : \sigma \in \mathbb{R}\} = -\frac{\lambda_n}{\varrho + \varepsilon} \ln \frac{\lambda_n}{e(\varrho + \varepsilon)}.$$

By Lemma 2, when  $\lambda_n$  is sufficiently large,

$$|a_n| < \exp\left\{-\frac{\lambda_n}{\varrho + \varepsilon} \ln \frac{\lambda_n}{e(\varrho + \varepsilon)}\right\}.$$

Let  $\varepsilon \rightarrow 0$ , then

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n \ln \lambda_n} \leq -\frac{1}{\varrho}.$$

Assume that  $\overline{\lim}_{n \rightarrow +\infty} (\ln |a_n|)/\lambda_n \ln \lambda_n < -1/\varrho$ . Then there exists  $0 < \varrho' < \varrho$  such that for sufficiently large  $n \in \mathbb{N}$

$$|a_n| < \exp\left\{-\frac{1}{\varrho'} \lambda_n \ln \lambda_n\right\}.$$

By the definition of  $\overline{A}_k$ , for sufficiently large  $k$ ,

$$\overline{A}_k < \sum_{j=n_k}^{n_k+p_k} \exp\left\{-\frac{\lambda_j}{\varrho'} \ln \lambda_j\right\} \leq \exp\left\{-\frac{k}{\varrho'} \ln k + \ln(p_k + 1)\right\}.$$

Note that  $\Delta = 0$ . Then

$$\overline{\lim}_{k \rightarrow +\infty} \frac{\ln \overline{A}_k}{k \ln k} \leq -\frac{1}{\varrho'} < -\frac{1}{\varrho}.$$

By Lemma 3,

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln \overline{M}(\sigma)}{-\sigma} < \varrho.$$

Note  $M(\sigma) \leq \overline{M}(\sigma)$ . Then

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma)}{-\sigma} < \varrho,$$

which contradicts the left-hand side of (5). Thus the necessity of the right-hand side of (5) is proved. By the above proof, we can easily prove the sufficiency of the right-hand side of (5). Thus we have proved the case  $0 < \varrho < +\infty$ .

By the case  $0 < \varrho < +\infty$ , we can easily prove the case  $\varrho = 0$ ,  $\varrho = +\infty$ . Thus the lemma is proved.

**Lemma 5.** *If  $f(s, \omega)$  satisfies  $\sigma_u(\omega) = -\infty$  a.s. and  $\Delta = 0$ , then*

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma, \omega)}{-\sigma} = \varrho \text{ a.s.} \Leftrightarrow \overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n \ln \lambda_n} = \begin{cases} -\infty, & \varrho = 0; \\ -\frac{1}{\varrho}, & 0 < \varrho < +\infty; \\ 0, & \varrho = +\infty. \end{cases}$$

*Proof.* Note that  $|a_n X_n(\omega)| = |a_n|$  a.s. Then

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n X_n(\omega)|}{\lambda_n \ln \lambda_n} = \overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n \ln \lambda_n} \text{ a.s.}$$

By Lemma 4, the lemma is proved.

**Lemma P.Z** (see [2]). *For every  $E \subseteq \Omega$  satisfying  $P(E) > 0$ , there exists a positive integer  $N = N(E)$  such that*

$$\int_E \left| \sum_{n=N}^{N'} c_n \varepsilon_n(\omega) \right|^2 d\omega \geq \frac{1}{2} P(E) \sum_{n=N}^{N'} |c_n|^2,$$

$$\int_E \left| \sum_{n=N}^{N'} c_n \gamma_n(\omega) \right|^2 d\omega \geq \frac{1}{2} P(E) \sum_{n=N}^{N'} |c_n|^2,$$

where  $\{c_n\}_{n=1}^{+\infty}$  is an arbitrary sequence of complex numbers.

### 3. PROOFS OF MAIN RESULT

*Proof of Theorem 1.* By Lemma 5, there exists  $\varrho \geq 0$  such that

$$(6) \quad \varrho = \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma, \omega)}{-\sigma} \text{ a.s.}$$

Case I:  $\varrho = 0$ . The theorem holds obviously.

Case II:  $0 < \varrho < \infty$ . Evidently,

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma; \alpha; \beta; \omega)}{-\sigma} \leq \varrho \text{ a.s.}$$

Assume that  $\overline{\lim}_{\sigma \rightarrow -\infty} -\sigma^{-1} \ln \ln M(\sigma; \alpha; \beta; \omega) < \varrho$  a.s. Then there exists  $E \subset \Omega$  satisfying  $P(E) > 0$  and  $0 < \varrho' < \varrho$  and  $\alpha_0, \beta_0, \alpha_0 < \beta_0$  such that

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma; \alpha_0, \beta_0; \omega)}{-\sigma} < \varrho', \quad \omega \in E.$$

Then, when  $-\sigma$  is large enough,  $\alpha_0 \leq t \leq \beta_0$ , we have

$$\left| \sum_{n=0}^{+\infty} a_n X_n(\omega) e^{-\lambda_n(\sigma+it)} \right| < \exp\{e^{-\varrho'\sigma}\}, \quad \omega \in E,$$

where  $N = N(E)$  is determined by Lemma P.Z for  $E$ . By Lemma P.Z, when  $-\sigma$  is large enough,

$$\frac{1}{2}P(E) \sum_{n=N}^{N'} |a_n|^2 e^{-2\lambda_n\sigma} \leq \int_E \left| \sum_{n=N}^{N'} a_n X_n(\omega) e^{-\lambda_n(\sigma+it)} \right|^2 d\omega, \quad N' > N, \omega \in E.$$

Since

$$\left| \sum_{n=N}^{N'} a_n X_n(\omega) e^{-\lambda_n(\sigma+it)} \right| \leq \sum_{n=0}^{+\infty} |a_n| e^{-\lambda_n\sigma} < +\infty, \quad -\infty < \sigma < 0, \omega \in E,$$

we have

$$\sum_{n=N}^{\infty} |a_n|^2 e^{-2\lambda_n\sigma} \leq \frac{2}{P(E)} \int_E \left| \sum_{n=N}^{+\infty} a_n X_n(\omega) e^{-\lambda_n(\sigma+it)} \right|^2 d\omega \leq 8 \exp\{2e^{-\varrho'\sigma}\}.$$

Therefore, when  $-\sigma$  is large enough,  $n \geq N$ , then

$$|a_n| e^{-\lambda_n\sigma} < 3 \exp\{e^{-\varrho'\sigma}\}.$$

Note that

$$\min\{e^{-\varrho'\sigma} + \lambda_n\sigma : \sigma \in \mathbb{R}\} = -\frac{\lambda_n}{\varrho'} \ln \frac{\lambda_n}{e(\varrho')}.$$

By Lemma 2, when  $n$  is sufficiently large,

$$|a_n| < 3 \exp\left\{-\frac{\lambda_n}{\varrho'} \ln \frac{\lambda_n}{e(\varrho')}\right\}.$$

Then

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n \ln \lambda_n} \leq -\frac{1}{\varrho'} < -\frac{1}{\varrho}.$$

By Lemma 5,

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma, \omega)}{-\sigma} < \varrho \quad \text{a.s.},$$

which contradicts (6).

Case III:  $\varrho = +\infty$ . We must have

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma; \alpha, \beta; \omega)}{-\sigma} = +\infty \quad \text{a.s.}$$

Otherwise, assume that  $\overline{\lim}_{\sigma \rightarrow -\infty} -\sigma^{-1} \ln \ln M(\sigma; \alpha, \beta; \omega) < +\infty$  a.s. Then by the proof of cases I and II,

$$\overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma, \omega)}{-\sigma} < +\infty \quad \text{a.s.}$$

which contradicts the given condition  $\overline{\lim}_{\sigma \rightarrow -\infty} -\sigma^{-1} \ln \ln M(\sigma, \omega) = +\infty$  a.s.  $\square$

#### 4. COROLLARY AND EXAMPLE

**Corollary 1.** Consider  $f(s, \omega)$  as in the introduction. If

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n} = -\infty \quad \text{and} \quad \overline{\lim}_{n \rightarrow +\infty} \frac{\ln n}{\lambda_n} = D < +\infty,$$

then  $f(s, \omega)$  a.s. (almost surely) satisfies

$$(7) \quad \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma, \omega)}{-\sigma} = \overline{\lim}_{\sigma \rightarrow -\infty} \frac{\ln \ln M(\sigma; \alpha, \beta; \omega)}{-\sigma}.$$

*Proof.* Note that  $\overline{\lim}_{n \rightarrow +\infty} \lambda_n^{-1} \ln |a_n X_n(\omega)| = \overline{\lim}_{n \rightarrow +\infty} \lambda_n^{-1} \ln |a_n|$  a.s. Then, by G. Valiron's formula (see [6], [9]),

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n} \leq \sigma_c(\omega) \leq \sigma_u(\omega) \leq \sigma_a(\omega) \leq \overline{\lim}_{n \rightarrow +\infty} \frac{\ln |a_n|}{\lambda_n} + \overline{\lim}_{n \rightarrow +\infty} \frac{\ln n}{\lambda_n} = -\infty \quad \text{a.s.},$$

which gives  $\sigma_u(\omega) = -\infty$  a.s.

By virtue of  $\overline{\lim}_{n \rightarrow +\infty} \lambda_n^{-1} \ln n = D < +\infty$ , for any  $\varepsilon > 0$ , when  $n \in \mathbb{N}$  is large enough,

$$n < e^{\lambda_n(D+\varepsilon)}.$$

Hence, when  $k$  is large enough,

$$p_k + 1 \leq n_{k+1} < e^{\lambda_{n_{k+1}}(D+\varepsilon)} \leq e^{(k+2)(D+\varepsilon)}.$$

Therefore

$$0 \leq \overline{\lim}_{k \rightarrow +\infty} \frac{\ln(p_k + 1)}{k \ln k} \leq \overline{\lim}_{k \rightarrow +\infty} \frac{(k+2)(D+\varepsilon)}{k \ln k} = 0.$$

Thus  $\Delta = 0$ . By Theorem 1, (7) holds.

Now we give an example to show that  $\Delta = 0$  is less restricting than

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln n}{\lambda_n} < +\infty.$$

**Example 1.** Consider  $f(s, \omega)$ . For any  $i \in \mathbb{N}$ , put  $[\ln i] = \max\{z: z \in \mathbb{N}, z \leq \ln i\}$ . For any  $k \in \mathbb{N}$ , take  $n_k = 1 + \sum_{i=1}^{k-1} ([\ln i])^i$ ,  $p_k = ([\ln k])^k - 1$ ,  $0 \leq p \leq p_k$ ,  $\lambda_{n_k+p} = k + p/p_k$ ,  $a_{n_k+p} = ([\ln k])^{-2k}$ . Prove that  $\sigma_u(\omega) = -\infty$  a.s.,  $\Delta = 0$ ,  $\overline{\lim}_{n \rightarrow +\infty} \lambda_n^{-1} \ln n = +\infty$ .

**Proof.** By the definition of  $\overline{A}_k(\omega)$ ,

$$\begin{aligned} \overline{A}_k(\omega) &= \sup_{0 \leq p \leq p_k, t \in \mathbb{R}} \left| \sum_{j=0}^p a_{n_k+j} X_{n_k+j}(\omega) e^{-it\lambda_{n_k+j}} \right| \\ &\leq \sum_{j=0}^{p_k} |a_{n_k+j}| = ([\ln k])^k ([\ln k])^{-2k} = ([\ln k])^{-k}, \text{ a.s.} \end{aligned}$$

Then by the formulas for  $\sigma_u(\omega)$  in terms of  $\overline{A}_k(\omega)$  in Section 2,

$$\sigma_u(\omega) = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln \overline{A}_k(\omega)}{k} \leq \overline{\lim}_{k \rightarrow +\infty} \frac{\ln([\ln k])^{-k}}{k} \leq \overline{\lim}_{k \rightarrow +\infty} \frac{\ln(\ln k - 1)^{-k}}{k} = -\infty \text{ a.s.}$$

By the definition of  $\Delta$ ,

$$\Delta = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln(p_k + 1)}{k \ln k} = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln([\ln k])^k}{k \ln k} \leq \overline{\lim}_{k \rightarrow +\infty} \frac{\ln(\ln k)^k}{k \ln k} = 0$$

but

$$\overline{\lim}_{n \rightarrow +\infty} \frac{\ln n}{\lambda_n} \geq \overline{\lim}_{n \rightarrow +\infty} \frac{\ln(p_k + 1)}{k+1} = \overline{\lim}_{k \rightarrow +\infty} \frac{\ln([\ln k])^k}{k+1} \geq \overline{\lim}_{k \rightarrow +\infty} \frac{\ln(\ln k - 1)^k}{k+1} = +\infty.$$

### References

- [1] *T. Kawata*: Fourier Analysis in Probability Theory. New York, Acad. Press, 1972.
- [2] *Paley Reac, A. Zygmund*: On some series of functions (1), (2), (3). Proc. Camb. Phil. Soc. 26 (1930), 337–357, 458–474; 28 (1932), 190–205.
- [3] *H. Bohr*: Collected Mathematical Works. Dansk Matematisk Forening Copenhagen, 1952.
- [4] *Yu Jiarong*: Some properties of random Dirichlet series. Acta Math. Sin. 21 (1978), 97–118. (In Chinese.)
- [5] *K. Knopp*: Über de Konvergenzabszisse des Laplace-Integrals. Math. Z. 54 (1951), 291–296. (In German.)
- [6] *G. Valiron*: Sur l’abscisse de convergence des séries de Dirichlet. Bulletin de la Société Mathématique de France 52 (1924), 166–174. (In French.)
- [7] *C. Tanaka*: Note on Dirichlet series (V), On the integral functions defined by Dirichlet series (I). Tôhoku Math. J., II. Ser. 5 (1953), 67–78.
- [8] *G. Valiron*: Théorie générale des séries de Dirichlet. Mémorial des sciences mathématiques, Fasc. 17 (1926). (In French.)
- [9] *G. Valiron*: Entire functions and Borel’s directions. Proc. Nat. Acad. Sci. USA 20 (1934), 211–215.

*Authors’ addresses*: *Zhendong Gu*, School of Mathematical Sciences, South China Normal University, Guangzhou, 510631, People’s Republic of China, e-mail: [guzhd@qq.com](mailto:guzhd@qq.com); *Daochun Sun*, School of Mathematical Sciences, South China Normal University, Guangzhou, 510631, P. R. China, e-mail: [1457330943@qq.com](mailto:1457330943@qq.com), [sundch@scnu.edu.cn](mailto:sundch@scnu.edu.cn).