

Emma Leppälä; Markku Niemenmaa

A note on loops of square-free order

Commentationes Mathematicae Universitatis Carolinae, Vol. 54 (2013), No. 1, 1--3

Persistent URL: <http://dml.cz/dmlcz/143146>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A note on loops of square-free order

EMMA LEPPÄLÄ, MARKKU NIEMENMAA

Abstract. Let Q be a loop such that $|Q|$ is square-free and the inner mapping group $I(Q)$ is nilpotent. We show that Q is centrally nilpotent of class at most two.

Keywords: loop, inner mapping group, centrally nilpotent loop

Classification: 20N05, 20D15

1. Introduction

A positive integer n is said to be square-free if n is divisible by no (perfect) square except 1. If n is square-free, then in the prime factorization of n , no prime occurs more than once. In this short note we show that if a loop Q has square-free order and the inner mapping group $I(Q)$ is nilpotent, then Q is centrally nilpotent of class at most two. Our notation in loop theory and group theory is standard. If $H \leq G$, then by H_G we denote the core of H in G (the largest normal subgroup of G contained in H). By $H^G = \langle H^g \mid g \in G \rangle$ we denote the normal closure of H in G .

2. Loops and connected transversals

If Q is a loop, then the two mappings $L_a(x) = ax$ and $R_a(x) = xa$ are permutations on Q for every $a \in Q$. We write $M(Q) = \langle L_a, R_a \mid a \in Q \rangle$ and say that $M(Q)$ is the multiplication group of Q . The stabilizer of the neutral element $e \in Q$ is denoted by $I(Q)$ and it is said to be the inner mapping group of Q . If we write $A = \{L_a \mid a \in Q\}$ and $B = \{R_a \mid a \in Q\}$, then the commutator subgroup $[A, B] \leq I(Q)$ and A, B are left transversals to $I(Q)$ in $M(Q)$. Furthermore, $I(Q)$ is core-free in $M(Q)$. A loop Q is centrally nilpotent of class at most two if and only if $M(Q)' \leq N_{M(Q)}(I(Q))$ (for the details, see [1, p. 281]).

If we replace $M(Q)$ by G and $I(Q)$ by H , then we have the following situation: $H \leq G$ and A and B are two left transversals to H in G . As $[A, B] \leq H$, we say that A and B are H -connected transversals in G . If $G' \leq N_G(H)$, then Q is centrally nilpotent of class at most two. In 1990 Kepka and Niemenmaa proved the following [3, Theorem 4.1]

Theorem 2.1. *A group G is isomorphic to the multiplication group of a loop if and only if there exists a subgroup H which is core-free in G and H -connected transversals A and B such that $G = \langle A, B \rangle$.*

In the following two lemmas, we assume that A and B are H -connected transversals in G .

Lemma 2.2. *Let N be a normal subgroup of G and let L be the core of HN in G . Then AL/L and BL/L are HL/L -connected transversals in G/L .*

Lemma 2.3. *If G is a finite group, H is nilpotent and $G = \langle A, B \rangle$, then H is subnormal in G .*

For the proofs, see [3, Lemma 2.8] and [4, Theorem 2.8].

Lemma 2.4. *If $H \leq G$ is nilpotent and subnormal in G , then the normal closure H^G is nilpotent.*

For the proof, see [2, Theorem 8.8, p. 29].

3. Main theorems

Theorem 3.1. *Let H be a nilpotent proper subgroup of a finite group G and let A and B be H -connected transversals in G such that $G = \langle A, B \rangle$. If $[G : H]$ is square-free, then $G' \leq N_G(H)$.*

PROOF: Clearly, we can assume that $H \neq 1$. Let G be a minimal counterexample. By Lemma 2.3, H is subnormal in G and then, by Lemma 2.4, H^G is nilpotent. Let $H^G = P \times Q$, where $P > 1$ is a Sylow p -subgroup of H^G and Q is a Hall π -subgroup of H^G with $p \notin \pi$ (here π is a — possibly empty — set of prime numbers).

Let us first assume that $Q > 1$. Then P and Q are both nontrivial characteristic subgroups of H^G , hence they are nontrivial normal subgroups of G . Let K be the core of HP in G . By Lemma 2.2, AK/K and BK/K are HP/K -connected transversals in G/K . As $[G/K : HP/K]$ is square-free, we may conclude that $(G/K)' \leq N_{G/K}(HP/K)$, which means that $G' \leq N_G(HP)$. In a similar manner we can show that $G' \leq N_G(HQ)$. Thus $G' \leq N_G(HP) \cap N_G(HQ) \leq N_G(HP \cap HQ) = N_G(H)$.

Then assume that $Q = 1$. It follows that $H \leq H^G = P$. If $x, y \in G$, then $x = ah$ and $y = bk$, where $a \in A$, $b \in B$ and $h, k \in H$. Since A and B are H -connected transversals in G , it follows that $[xP, yP] = [aP, bP] = [a, b]P = P$. We conclude that G/P is an abelian group and thus $G' \leq P$. As $[G : H]$ is square-free, we see that $[P : H] = p$ or $P = H$. But then $N_G(H) \geq P$, whence $G' \leq N_G(H)$. $\square$

We can slightly loosen the condition on $[G : H]$ if we assume that H is a p -group.

Theorem 3.2. *Let $H < G$ be a p -group and let A and B be H -connected transversals in G such that $G = \langle A, B \rangle$. If $[G : H] = pk$ and $p \nmid k$, then $G' \leq N_G(H)$.*

The proof is similar to the proof of Theorem 3.1. By using Theorems 3.1, 3.2 and 2.1, we get

Corollary 3.3. *If Q is a loop of square-free order and $I(Q)$ is nilpotent, then Q is centrally nilpotent of class at most two.*

Corollary 3.4. *Let Q be a loop and $|Q| = pk$, where p is a prime number and $p \nmid k$. If $I(Q)$ is a p -group, then Q is centrally nilpotent of class at most two.*

Remark. If G is a finite group and $|G|$ is square-free, then G is solvable (this is an easy consequence of the Burnside normal complement theorem). If $|G|$ is square-free and $I(G) = \text{Inn}(G)$ (the group of inner automorphisms of G) is nilpotent, then G is cyclic.

It is easy to construct loops of order six which are not solvable. Now, consider the following loop Q of order six:

1	2	3	4	5	6
2	1	4	3	6	5
3	4	5	6	2	1
4	3	6	5	1	2
5	6	1	2	3	4
6	5	2	1	4	3

Here $|Q|$ is square-free, $M(Q)$ has order 24, $I(Q)$ is isomorphic to Klein's four group and Q is centrally nilpotent of class two.

REFERENCES

- [1] Bruck R.H., *Contributions to the theory of loops*, Trans. Amer. Math. Soc. **60** (1946), 245–354.
- [2] Doerk K., Hawkes T., *Finite Soluble Groups*, de Gruyter, Berlin, 1992.
- [3] Kepka T., Niemenmaa M., *On multiplication groups of loops*, J. Algebra **135** (1990), 112–122.
- [4] Niemenmaa M., *Finite loops with nilpotent inner mapping groups are centrally nilpotent*, Bull. Aust. Math. Soc. **79** (2009), no. 1, 109–114.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF OULU, PL 3000, 90014 OULU, FINLAND

E-mail: emma.leppala@oulu.fi,
markku.niemenmaa@oulu.fi

(Received June 6, 2012, revised November 29, 2012)