

Norbert Mahoungou Moukala; Basile Guy Richard Bossoto
Prolongation of Poisson 2-form on Weil bundles

Archivum Mathematicum, Vol. 52 (2016), No. 2, 91–111

Persistent URL: <http://dml.cz/dmlcz/145748>

Terms of use:

© Masaryk University, 2016

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

PROLONGATION OF POISSON 2-FORM ON WEIL BUNDLES

NORBERT MAHOUNGOU MOUKALA AND BASILE GUY RICHARD BOSSOTO

ABSTRACT. In this paper, M denotes a smooth manifold of dimension n , A a Weil algebra and M^A the associated Weil bundle. When (M, ω_M) is a Poisson manifold with 2-form ω_M , we construct the 2-Poisson form $\omega_{M^A}^A$, prolongation on M^A of the 2-Poisson form ω_M . We give a necessary and sufficient condition for that M^A be an A -Poisson manifold.

1. INTRODUCTION

1.1. Weil algebra and Weil bundle. In what follows, all structures are assumed to be of class C^∞ . We denote by M a smooth differential manifold, $C^\infty(M)$ the algebra of differentiable functions on M and by $\mathfrak{X}(M)$, the $C^\infty(M)$ -module of vectors field on M .

A Weil algebra is a real, unitary, commutative algebra of finite dimension with a unique maximal ideal of codimension 1 on \mathbb{R} , [15]. Let A be a Weil algebra and \mathfrak{m} be its maximal ideal. We have $A = \mathbb{R} \oplus \mathfrak{m}$ and the first projection

$$A = \mathbb{R} \oplus \mathfrak{m} \rightarrow \mathbb{R}$$

is a homomorphism of algebras which is surjective, called augmentation and the unique non zero integer $h \in \mathbb{N}$ such that $\mathfrak{m}^h \neq (0)$ and $\mathfrak{m}^{h+1} = (0)$ is the height of A , [15].

If M is a smooth manifold, and A a Weil algebra of maximal ideal \mathfrak{m} , an infinitely near point to $x \in M$ of kind A is a homomorphism of algebras

$$\xi: C^\infty(M) \rightarrow A$$

such that $[\xi(f) - f(x)] \in \mathfrak{m}$ for any $f \in C^\infty(M)$. i.e. the real part of $\xi(f)$ is exactly $f(x)$, [15].

We denote by M_x^A the set of all infinitely near points to $x \in M$ of kind A and $M^A = \bigcup_{x \in M} M_x^A$ the manifold of infinitely near points of kind A . We have $\dim M^A = \dim M \times \dim A$, [6].

2010 *Mathematics Subject Classification:* primary 58A20; secondary 58A32, 17D63, 53D17, 53D05.

Key words and phrases: Weil bundle, Weil algebra, Poisson manifold, Lie derivative, Poisson 2-form.

Received September 24, 2015. Editor I. Kolář.

DOI: 10.5817/AM2016-2-91

When both M and N are smooth manifolds and when $h: M \rightarrow N$ is a differentiable application, then the map

$$h^A: M^A \rightarrow N^A, \quad \xi \mapsto h^A(\xi),$$

such that, for any $g \in C^\infty(N)$,

$$[h^A(\xi)](g) = \xi(g \circ h)$$

is also differentiable. When h is a diffeomorphism, it is the same for h^A , [1].

Moreover, if $\varphi: A \rightarrow B$ is a homomorphism of Weil algebras, for any smooth manifold M , the map

$$\varphi_M: M^A \rightarrow M^B, \quad \xi \mapsto \varphi \circ \xi$$

is differentiable. In particular, the augmentation

$$A \rightarrow \mathbb{R}$$

defines for any smooth manifold M , the projection

$$\pi_M: M^A \rightarrow M,$$

which assigns every infinitely near point to $x \in M$ to its origin x . Thus (M^A, π_M, M) defines the bundle of infinitely near points or simply Weil bundle (see [3], [6], [15], [9]).

If (U, φ) is a local chart of M with coordinate functions (x_1, x_2, \dots, x_n) , the application

$$U^A \rightarrow A^n, \quad \xi \mapsto (\xi(x_1), \xi(x_2), \dots, \xi(x_n)),$$

is a bijection from U^A into an open of A^n . The manifold M^A is a smooth manifold modeled over A^n , that is to say an A -manifold of dimension n , [2], [13].

The set, $C^\infty(M^A, A)$ of differentiable functions on M^A with values in A is a commutative, unitary algebra over A . When one identifies \mathbb{R}^A with A , for $f \in C^\infty(M)$, the map

$$f^A: M^A \rightarrow A, \quad \xi \mapsto \xi(f)$$

is differentiable and the map

$$C^\infty(M) \rightarrow C^\infty(M^A, A), \quad f \mapsto f^A,$$

is an injective homomorphism of algebras and we have:

$$(f + g)^A = f^A + g^A; \quad (\lambda \cdot f)^A = \lambda \cdot f^A; \quad (f \cdot g)^A = f^A \cdot g^A$$

for $\lambda \in \mathbb{R}$, $f, g \in C^\infty(M)$.

We denote $\mathfrak{X}(M^A)$, the set of all vector fields on M^A . According to [2], [8] we have the following equivalent assertions:

Theorem 1. *The following assertions are equivalent:*

- (1) *A vector field on M^A is a differentiable section of the tangent bundle (TM^A, π_{M^A}, M^A) .*
- (2) *A vector field on M^A is a derivation of $C^\infty(M^A)$.*
- (3) *A vector field on M^A is a derivation of $C^\infty(M^A, A)$ which is A -linear.*

- (4) A vector field on M^A is a linear map $X: C^\infty(M) \rightarrow C^\infty(M^A, A)$ such that

$$X(f \cdot g) = X(f) \cdot g^A + f^A \cdot X(g), \quad \text{for any } f, g \in C^\infty(M).$$

Consequently [8],

Theorem 2. *The map*

$$\mathfrak{X}(M^A) \times \mathfrak{X}(M^A) \longrightarrow \mathfrak{X}(M^A), (X, Y) \longmapsto [X, Y] = X \circ Y - Y \circ X$$

is skew-symmetric A -bilinear and defines a structure of A -Lie algebra over $\mathfrak{X}(M^A)$.

Thus, if $\text{Der}_A[C^\infty(M^A, A)]$ denotes the $C^\infty(M^A, A)$ -module of derivations of $C^\infty(M^A, A)$ which are A -linear, a vector field on M^A is a derivation of $C^\infty(M^A, A)$ which is A -linear i.e. a A -linear map

$$X: C^\infty(M^A, A) \rightarrow C^\infty(M^A, A)$$

such that

$$X(\varphi \cdot \psi) = X(\varphi) \cdot \psi + \varphi \cdot X(\psi), \quad \text{for any } \varphi, \psi \in C^\infty(M^A, A).$$

Thus, we have

$$\mathfrak{X}(M^A) = \text{Der}_A [C^\infty(M^A, A)].$$

Proposition 1 ([2], [8]). *If $\theta: C^\infty(M) \rightarrow C^\infty(M)$ is a vector field on M , then there exists one and only one A -linear derivation*

$$\theta^A: C^\infty(M^A, A) \rightarrow C^\infty(M^A, A)$$

such that

$$\theta^A(f^A) = [\theta(f)]^A$$

for any $f \in C^\infty(M)$.

Proposition 2 ([2], [8]). *If θ, θ_1 and θ_2 are vector fields on M and if $f \in C^\infty(M)$, then we have:*

- (1) $(\theta_1 + \theta_2)^A = \theta_1^A + \theta_2^A;$
- (2) $(f \cdot \theta)^A = f^A \cdot \theta^A;$
- (3) $[\theta_1, \theta_2]^A = [\theta_1^A, \theta_2^A].$

Corollary 1. *The map*

$$\mathfrak{X}(M) \rightarrow \text{Der}_A [C^\infty(M^A, A)], \theta \mapsto \theta^A$$

is an injective homomorphism of \mathbb{R} -Lie algebras. If $\mu: A \rightarrow A$, is a \mathbb{R} -endomorphism, and $\theta: C^\infty(M) \rightarrow C^\infty(M)$ a vector field on M , then

$$\theta^A(\mu \circ f^A) = \mu \circ [\theta(f)]^A \quad \text{for any } f \in C^\infty(M).$$

1.2. Poisson manifold. We recall that a Poisson structure on a smooth manifold M is due to the existence of a bracket $\{ , \}_M$ on $C^\infty(M)$ such that the pair $(C^\infty(M), \{ , \}_M)$ is a real Lie algebra such that, for any $f \in C^\infty(M)$ the map

$$\text{ad}(f) : C^\infty(M) \rightarrow C^\infty(M), \quad g \mapsto \{f, g\}_M$$

is a derivation of commutative algebra i.e.

$$\{f, g \cdot h\}_M = \{f, g\}_M \cdot h + g \cdot \{f, h\}_M$$

for $f, g, h \in C^\infty(M)$. In this case we say that $C^\infty(M)$ is a Poisson algebra and M is a Poisson manifold ([5], [14], [11]).

Let $\Omega_{\mathbb{R}}[C^\infty(M)]$ be the $C^\infty(M)$ -module of Kähler differentials of $C^\infty(M)$ and

$$\delta_M : C^\infty(M) \rightarrow \Omega_{\mathbb{R}}[C^\infty(M)], \quad f \mapsto \overline{f \otimes 1_{C^\infty(M)} - 1_{C^\infty(M)} \otimes f}$$

the canonical derivation which the image of δ_M generates the $C^\infty(M)$ -module $\Omega_{\mathbb{R}}[C^\infty(M)]$ i.e. for $x \in \Omega_{\mathbb{R}}[C^\infty(M)]$,

$$x = \sum_{i \in I: \text{ finite}} f_i \cdot \delta_M(g_i),$$

with $f_i, g_i \in C^\infty(M)$ for any $i \in I$ ([4], [11], [10]).

The manifold M is a Poisson manifold if and only if there exists a skew-symmetric 2-form

$$\omega_M : \Omega_{\mathbb{R}}[C^\infty(M)] \times \Omega_{\mathbb{R}}[C^\infty(M)] \rightarrow C^\infty(M)$$

such that for any f and g in $C^\infty(M)$,

$$\{f, g\}_M = -\omega_M[\delta_M(f), \delta_M(g)]$$

defines a structure of Lie algebra over $C^\infty(M)$, [11], [10]. In this case, we say that ω_M is the Poisson 2-form of the Poisson manifold M and we denote (M, ω_M) the Poisson manifold of Poisson 2-form ω_M .

The main goal of this paper is to study the prolongation of the Poisson 2-form ω_M of Poisson manifold on Weil bundles.

2. THE ALGEBRA OF KÄHLER FORMS ON $C^\infty(M^A, A)$

Definition 1. The $C^\infty(M^A)$ -module of Kähler differentials of $C^\infty(M^A)$ is the set

$$\Omega_{\mathbb{R}}[C^\infty(M^A)] = \frac{J}{J^2}$$

where J is the $C^\infty(M^A)$ -submodule of $C^\infty(M^A) \bigotimes_{\mathbb{R}} C^\infty(M^A)$ generated by the elements of the form $F \otimes 1_{C^\infty(M^A)} - 1_{C^\infty(M^A)} \otimes F$ with $F \in C^\infty(M^A)$. Thus, the map

$$d_{M^A} : C^\infty(M^A) \rightarrow \Omega_{\mathbb{R}}[C^\infty(M^A)], F \mapsto \overline{F \otimes 1_{C^\infty(M^A)} - 1_{C^\infty(M^A)} \otimes F}$$

is a derivation and the image of d_{M^A} generates $\Omega_{\mathbb{R}}[C^\infty(M^A)]$.

The A -algebra $C^\infty(M^A, A) \otimes_A C^\infty(M^A, A)$ admits a structure of $C^\infty(M^A, A)$ -module defined by the homomorphism of A -algebras

$$C^\infty(M^A, A) \rightarrow C^\infty(M^A, A) \otimes_A C^\infty(M^A, A), \varphi \mapsto \varphi \otimes 1_{C^\infty(M^A, A)}.$$

In this case, we say that $C^\infty(M^A, A) \otimes_A C^\infty(M^A, A)$ admits a structure of $C^\infty(M^A, A)$ -module defined by the first factor. The second factor is defined by

$$C^\infty(M^A, A) \rightarrow C^\infty(M^A, A) \otimes_A C^\infty(M^A, A), \varphi \mapsto 1_{C^\infty(M^A, A)} \otimes \varphi.$$

The map

$$C^\infty(M^A, A) \times C^\infty(M^A, A) \rightarrow C^\infty(M^A, A), (\varphi, \psi) \mapsto \varphi \cdot \psi$$

being A -bilinear, then there exists a unique A -linear map

$$m: C^\infty(M^A, A) \otimes_A C^\infty(M^A, A) \rightarrow C^\infty(M^A, A)$$

such that

$$m(\varphi \otimes \psi) = \varphi \cdot \psi.$$

The kernel of m is the $C^\infty(M^A, A)$ -submodule of $C^\infty(M^A, A) \otimes C^\infty(M^A, A)$ generated by the elements of the form $\varphi \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes \varphi$ with $\varphi \in C^\infty(M^A, A)$.

We denote $\Omega_A[C^\infty(M^A, A)]$, the $C^\infty(M^A, A)$ -module of Kähler differentials of $C^\infty(M^A, A)$ which are A -linears. In this case, for $\varphi \in C^\infty(M^A, A)$, we denote $\overline{\varphi \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes \varphi}$, the class of $\varphi \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes \varphi$ in $C^\infty(M^A, A)$.

The map

$$C^\infty(M) \rightarrow \Omega_A[C^\infty(M^A, A)], f \mapsto \overline{f^A \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes f^A}$$

is a derivation.

Thus,

Proposition 3. *There exists a unique A -linear derivation*

$$\delta_{M^A}^A: C^\infty(M^A, A) \rightarrow \Omega_A[C^\infty(M^A, A)]$$

such that

$$\delta_{M^A}^A(f^A) = [\delta_M(f)]^A$$

for any $f \in C^\infty(M)$.

Proof. Let

$$\begin{aligned} \delta_{M^A}^A: C^\infty(M^A, A) &\xrightarrow{\sigma^{-1}} A \otimes C^\infty(M^A) \xrightarrow{\text{id}_A \otimes d_{M^A}} A \\ &\otimes \Omega_{\mathbb{R}}[C^\infty(M^A)] \xrightarrow{\varpi} \Omega_A[C^\infty(M^A, A)] \end{aligned}$$

be that map, where

$$\sigma^{-1}: \varphi = \sum_{\alpha=1}^{\dim A} (a_\alpha^* \circ \varphi) \cdot a_\alpha \longmapsto \sum_{\alpha=1}^{\dim A} a_\alpha \otimes (a_\alpha^* \circ \varphi)$$

with $(a_\alpha)_{\alpha=1,\dots,\dim A}$ a basis of A and $(a_\alpha^*)_{\alpha=1,\dots,\dim A}$ the dual basis of the basis $(a_\alpha)_{\alpha=1,\dots,\dim A}$,

$$\begin{aligned} \text{id}_A \otimes d_{M^A} : \sum_{\alpha=1}^{\dim A} a_\alpha \otimes (a_\alpha^* \circ \varphi) &\mapsto \sum_{\alpha=1}^{\dim A} a_\alpha \otimes d_{M^A}(a_\alpha^* \circ \varphi) \\ &= \sum_{\alpha=1}^{\dim A} a_\alpha \otimes \left[\overline{(a_\alpha^* \circ \varphi) \otimes 1_{C^\infty(M^A)} - 1_{C^\infty(M^A)} \otimes (a_\alpha^* \circ \varphi)} \right], \\ \varpi : \sum_{\alpha=1}^{\dim A} a_\alpha \otimes d_{M^A}(a_\alpha^* \circ \varphi) &\mapsto \sum_{\alpha=1}^{\dim A} \left[\overline{(a_\alpha^* \circ \varphi)a_\alpha \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes (a_\alpha^* \circ \varphi)a_\alpha} \right]. \end{aligned}$$

Thus,

$$\begin{aligned} \delta_{M^A}^A(\varphi) &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\varphi) \\ &= \sum_{\alpha=1}^{\dim A} \overline{[(a_\alpha^* \circ \varphi)a_\alpha \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes (a_\alpha^* \circ \varphi)a_\alpha]}. \end{aligned}$$

For any $\varphi, \psi \in C^\infty(M^A, A)$, we have

$$\begin{aligned} \delta_{M^A}^A(\varphi + \psi) &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\varphi + \psi) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\varphi) + \sigma^{-1}(\psi)) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\varphi)) + [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\psi)) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\varphi) + [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\psi) \\ &= \delta_{M^A}^A(\varphi) + \delta_{M^A}^A(\psi). \end{aligned}$$

For any $\varphi \in C^\infty(M^A, A)$ and $a \in A$, we have

$$\begin{aligned} \delta_{M^A}^A(a \cdot \varphi) &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](a \cdot \varphi) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(a \cdot \varphi)) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A})](a \cdot \sigma^{-1}(\varphi)) \\ &= a \cdot [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\varphi)) \\ &= a \cdot \delta_{M^A}^A(\varphi). \end{aligned}$$

For any $\varphi, \psi \in C^\infty(M^A, A)$, we have

$$\begin{aligned}\delta_{M^A}^A(\varphi \cdot \psi) &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\varphi \cdot \psi) \\ &= [\varpi \circ (id_A \otimes d_{M^A})](\sigma^{-1}(\varphi \cdot \psi)) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\varphi) \cdot \sigma^{-1}(\psi)) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\varphi)) \cdot \psi + \varphi \cdot [\varpi \circ (\text{id}_A \otimes d_{M^A})](\sigma^{-1}(\psi)) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\varphi) \cdot \psi + \varphi \cdot [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\psi) \\ &= [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\varphi) + [\varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}](\psi) \\ &= \delta_{M^A}^A(\varphi) \cdot \psi + \varphi \cdot \delta_{M^A}^A(\psi).\end{aligned}$$

As

$$\delta_M: C^\infty(M) \rightarrow \Omega_{\mathbb{R}}[C^\infty(M)]$$

is a derivation, then the map

$$C^\infty(M) \rightarrow \Omega_A[C^\infty(M^A, A)], \quad f \mapsto [\delta_M(f)]^A$$

is a derivation. Thus, for any $f \in C^\infty(M)$

$$\begin{aligned}\delta_{M^A}^A(f^A) &= \varpi \circ (\text{id}_A \otimes d_{M^A}) \circ \sigma^{-1}(f^A) \\ &= \overline{\sum_{\alpha=1}^{\dim A} [(a_\alpha^* \circ f^A)a_\alpha \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes (a_\alpha^* \circ f^A)a_\alpha]} \\ &= \overline{\sum_{\alpha=1}^{\dim A} (a_\alpha^* \circ f^A)a_\alpha \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes \sum_{\alpha=1}^{\dim A} (a_\alpha^* \circ f^A)a_\alpha} \\ &= \overline{f^A \otimes 1_{C^\infty(M^A, A)} - 1_{C^\infty(M^A, A)} \otimes f^A} \\ &= \overline{[f \otimes 1_{C^\infty(M)} - 1_{C^\infty(M)} \otimes f]}^A\end{aligned}$$

i.e.

$$\delta_{M^A}^A(f^A) = [\delta_M(f)]^A.$$

□

Proposition 4. *The map*

$$\Omega_{\mathbb{R}}[C^\infty(M)] \rightarrow \Omega_A[C^\infty(M^A, A)], \quad x \mapsto x^A$$

is an injective homomorphism of \mathbb{R} -modules.

Proof. Let

$$\Psi: \Omega_{\mathbb{R}}[C^\infty(M)] \rightarrow \Omega_A[C^\infty(M^A, A)], \quad x \mapsto x^A$$

be that map. □

For any $x, y \in \Omega_{\mathbb{R}}[C^\infty(M)]$,

$$\begin{aligned}\Psi(x + y) &= (x + y)^A \\ &= \left(\sum_{i \in I: \text{ finite}} f_i \cdot \delta_M(f'_i) + \sum_{j \in I: \text{ finite}} g_j \cdot \delta_M(g'_j) \right)^A \\ &= \left(\sum_{i \in I: \text{ finite}} f_i \cdot \delta_M(f'_i) \right)^A + \left(\sum_{j \in I: \text{ finite}} g_j \cdot \delta_M(g'_j) \right)^A \\ &= x^A + y^A.\end{aligned}$$

For any $x \in \Omega_{\mathbb{R}}[C^\infty(M)]$ and for $\lambda \in \mathbb{R}$,

$$\begin{aligned}\Psi(\lambda \cdot x) &= (\lambda \cdot x)^A \\ &= \left(\lambda \cdot \sum_{i \in I: \text{ finite}} f_i \cdot \delta_M(f'_i) \right)^A \\ &= \lambda \cdot \left(\sum_{i \in I: \text{ finite}} f_i \cdot \delta_M(f'_i) \right)^A \\ &= \lambda \cdot x^A.\end{aligned}$$

The pair $(\Omega_A[C^\infty(M^A, A)], \delta_{M^A}^A)$ satisfies the following universal property: for every $C^\infty(M^A, A)$ -module E and every A -derivation

$$\Phi: C^\infty(M^A, A) \rightarrow E,$$

there exists a unique $C^\infty(M^A, A)$ -linear map

$$\tilde{\Phi}: \Omega_A[C^\infty(M^A, A)] \rightarrow E$$

such that

$$\tilde{\Phi} \circ \delta_{M^A}^A = \Phi.$$

In other words, there exists a unique $\tilde{\Phi}$ which makes the following diagram commutative

$$\begin{array}{ccc} \Omega_A[C^\infty(M^A, A)] & & \\ \delta_{M^A}^A \uparrow & \searrow \tilde{\Phi} & \\ C^\infty(M^A, A) & \xrightarrow[\Phi]{} & E. \end{array}$$

This fact implies the existence of a natural isomorphism of $C^\infty(M^A, A)$ -modules $\text{Hom}_{C^\infty(M^A, A)}(\Omega_A[C^\infty(M^A, A)], E) \rightarrow \text{Der}_A[C^\infty(M^A, A)], E]$, $\psi \mapsto \psi \circ \delta_{M^A}^A$.

In particular, if $E = C^\infty(M^A, A)$, we have

$$\begin{aligned}\Omega_A[C^\infty(M^A, A)]^* &\simeq \text{Der}_A[C^\infty(M^A, A)] \\ &= \mathfrak{X}(M^A).\end{aligned}$$

For any $p \in \mathbb{N}$, $\Lambda^p(\Omega_A[C^\infty(M^A, A)]) = \mathfrak{L}_{sks}^p(\Omega_A[C^\infty(M^A, A)], C^\infty(M^A, A))$ denotes the $C^\infty(M^A, A)$ -module of skew-symmetric multilinear forms of degree p from $\Omega_A[C^\infty(M^A, A)]$ into $C^\infty(M^A, A)$ and

$$\Lambda(\Omega_A[C^\infty(M^A, A)]) = \bigoplus_{p \in \mathbb{N}} \Lambda^p(\Omega_A[C^\infty(M^A, A)])$$

the exterior $C^\infty(M^A, A)$ -algebra of $\Omega_A[C^\infty(M^A, A)]$.

$$\Lambda^0(\Omega_A[C^\infty(M^A, A)]) = C^\infty(M^A, A),$$

$$\Lambda^1(\Omega_A[C^\infty(M^A, A)]) = \Omega_A[C^\infty(M^A, A)]^*.$$

We denote,

$$\delta_{M^A}^A = \delta_{M^A} : \Lambda(\Omega_A[C^\infty(M^A, A)]) \rightarrow \Lambda(\Omega_A[C^\infty(M^A, A)])$$

a unique derivation, of degree +1, which extends the canonical derivation

$$\delta_{M^A}^0 : C^\infty(M^A, A) \rightarrow \Omega_A[C^\infty(M^A, A)].$$

For any $\varphi, \psi, \psi_1, \psi_2, \dots, \psi_p \in C^\infty(M^A, A)$ and $\omega \in \Omega_A[C^\infty(M^A, A)]^*$, we get

1. $\delta_{M^A}(\varphi \cdot \delta_{M^A}(\psi_1) \wedge \cdots \wedge \delta_{M^A}(\psi_p)) = \delta_{M^A}(\varphi) \wedge \delta_{M^A}(\psi_1) \wedge \cdots \wedge \delta_{M^A}(\psi_p).$
2. $\delta_{M^A}^1[\psi \cdot \delta_{M^A}^0(\varphi)] = \delta_{M^A}^0(\psi) \wedge \delta_{M^A}^0(\varphi).$
3. $\delta_{M^A}^1(\varphi \cdot \omega) = \delta_{M^A}^0(\varphi) \wedge \omega + \varphi \cdot \delta_{M^A}^1(\omega).$

Proposition 5. *If $\eta \in \Lambda^p(\Omega_\mathbb{R}[C^\infty(M)])$, then $\eta^A \in \Lambda^p(\Omega_A[C^\infty(M^A, A)])$.*

Proof. Indeed, for any $\eta \in \Lambda^p(\Omega_\mathbb{R}[C^\infty(M)])$, η is of the form $\delta_M(f_1) \wedge \cdots \wedge \delta_M(f_p)$ with $f_1, f_2, \dots, f_p \in C^\infty(M)$.

$$\begin{aligned} \eta^A &= [\delta_M(f_1) \wedge \cdots \wedge \delta_M(f_p)]^A \\ &= [\delta_M(f_1)]^A \wedge \cdots \wedge [\delta_M(f_p)]^A \\ &= \delta_{M^A}^0(f_1^A) \wedge \cdots \wedge \delta_{M^A}^0(f_p^A). \end{aligned}$$

Thus, the $C^\infty(M^A, A)$ -module $\Lambda^p(\Omega_A[C^\infty(M^A, A)])$ is generated by elements of the form

$$\eta^A = \delta_{M^A}^0(\varphi_1) \wedge \cdots \wedge \delta_{M^A}^0(\varphi_p)$$

with $\varphi_1 = f_1^A, \dots, \varphi_p = f_p^A \in C^\infty(M^A, A)$. \square

The algebra

$$\Lambda(\Omega_A[C^\infty(M^A, A)]) = \bigoplus_{p \in \mathbb{N}} \Lambda^p(\Omega_A[C^\infty(M^A, A)])$$

is the algebra of Kähler forms on $C^\infty(M^A, A)$.

The pair $(\Lambda(\Omega_A[C^\infty(M^A, A)]), \delta_{M^A}^A)$ is a differential complex and the map

$$A \times \Omega_\mathbb{R}[C^\infty(M)] \mapsto \Omega_A[C^\infty(M^A, A)], (a, x) \mapsto a \cdot x^A$$

induces the morphism of the differential complex $(A \otimes \Lambda(\Omega_{\mathbb{R}}[C^\infty(M)]), \text{id}_A \otimes \delta_M)$ into the differential complex $(\Lambda(\Omega_A[C^\infty(M^A, A)]), \delta_{M^A}^A)$.

3. LIE DERIVATIVE WITH RESPECT TO A DERIVATION ON M^A

Let

$$\theta: C^\infty(M) \rightarrow C^\infty(M)$$

be a derivation and

$$\sigma_\theta: [\Omega_{\mathbb{R}}[C^\infty(M)]]^p \longrightarrow \Lambda^p(\Omega_{\mathbb{R}}[C^\infty(M)]) ,$$

be the $C^\infty(M)$ -skew-symmetric multilinear map such that for any $x_1, x_2, \dots, x_p \in \Omega_{\mathbb{R}}[C^\infty(M)]$,

$$\sigma_\theta(x_1, x_2, \dots, x_p) = \sum_{i=1}^p (-1)^{i-1} \tilde{\theta}(x_i) \cdot x_1 \wedge \cdots \wedge \hat{x}_i \wedge \cdots \wedge x_p ,$$

where

$$\tilde{\theta}: \Omega_{\mathbb{R}}[C^\infty(M)] \rightarrow C^\infty(M)$$

is a unique $C^\infty(M)$ -linear map such that $\tilde{\theta} \circ \delta_M = \theta$. Then,

$$\sigma_{\theta^A}^A: [\Omega_A[C^\infty(M^A, A)]]^p \rightarrow \Lambda^p(\Omega_A[C^\infty(M^A, A)])$$

is a unique $C^\infty(M^A, A)$ -skew-symmetric multilinear map such that

$$\sigma_{\theta^A}^A(x_1^A, x_2^A, \dots, x_p^A) = [\sigma_\theta(x_1, x_2, \dots, x_p)]^A .$$

We denote

$$\widetilde{\sigma_{\theta^A}^A}: \Lambda^p(\Omega_A[C^\infty(M^A, A)]) \rightarrow \Lambda^{p-1}(\Omega_A[C^\infty(M^A, A)]) ,$$

the unique $C^\infty(M^A, A)$ -skew-symmetric multilinear map such that

$$\widetilde{\sigma_{\theta^A}^A}(x_1^A \wedge x_2^A \wedge \cdots \wedge x_p^A) = \sigma_{\theta^A}^A(x_1^A, x_2^A, \dots, x_p^A)$$

i.e. $\sigma_{\theta^A}^A$ induces a derivation

$$i_{\theta^A} = \widetilde{\sigma_{\theta^A}^A}: \Lambda(\Omega_A[C^\infty(M^A, A)]) \rightarrow \Lambda(\Omega_A[C^\infty(M^A, A)])$$

of degree -1 .

Proposition 6. *For any $\theta \in \text{Der}_{\mathbb{R}}[C^\infty(M)]$ and for any $\eta \in \Lambda^p(\Omega_{\mathbb{R}}[C^\infty(M)])$, we have*

$$i_{\theta^A}(\eta^A) = [i_\theta(\eta)]^A .$$

Proof. If $\eta \in \Lambda^p(\Omega_{\mathbb{R}}[C^\infty(M)])$, then there exists $f_1, f_2, \dots, f_p \in C^\infty(M)$, such that $\eta = \delta_M(f_1) \wedge \cdots \wedge \delta_M(f_p)$. Thus,

$$\begin{aligned} i_{\theta^A}(\eta^A) &= i_{\theta^A}([\delta_M(f_1) \wedge \cdots \wedge \delta_M(f_p)]^A) \\ &= i_{\theta^A}([\delta_M(f_1)]^A \wedge \cdots \wedge [\delta_M(f_p)]^A) \\ &= \sigma_{\theta^A}^A([\delta_M(f_1)]^A, \dots, [\delta_M(f_p)]^A) \end{aligned}$$

$$\begin{aligned}
&= [\sigma_\theta(\delta_M(f_1), \dots, \delta_M(f_p))]^A \\
&= [i_\theta(\delta_M(f_1) \wedge \dots \wedge \delta_M(f_p))]^A \\
&= [i_\theta(\eta)]^A.
\end{aligned}$$

For $p = 1$, we have

$$\begin{aligned}
i_{\theta^A} &= \widetilde{\sigma_{\theta^A}} : \Lambda^1(\Omega_A[C^\infty(M^A, A)]) \\
&= \Omega_A[C^\infty(M^A, A)]^* \rightarrow \Lambda^0(\Omega_A[C^\infty(M^A, A)]) = C^\infty(M^A, A),
\end{aligned}$$

and for any $y \in \Omega_{\mathbb{R}}[C^\infty(M)]$,

$$i_{\theta^A}(y^A) = \widetilde{\theta^A}(y^A).$$

For $p = 2$, we have

$$\sigma_{\theta^A}^A : \Omega_A[C^\infty(M^A, A)] \times \Omega_A[C^\infty(M^A, A)] \rightarrow C^\infty(M^A, A)$$

and for any $x, y \in \Omega_{\mathbb{R}}[C^\infty(M)]$,

$$\sigma_{\theta^A}^A(x^A, y^A) = \widetilde{\theta^A}(x^A) \cdot y^A - \widetilde{\theta^A}(y^A) \cdot x^A.$$

Thus, the map

$$i_{\theta^A} : \Lambda^2(\Omega_A[C^\infty(M^A, A)]) \rightarrow \Omega_A[C^\infty(M^A, A)]^*$$

is the unique $C^\infty(M^A, A)$ -linear map such that

$$i_{\theta^A}(x^A \wedge y^A) = \sigma_{\theta^A}^A(x^A, y^A) = \widetilde{\theta^A}(x^A) \cdot y^A - \widetilde{\theta^A}(y^A) \cdot x^A.$$

□

Definition 2. The Lie derivative with respect to $D \in \text{Der}_A[C^\infty(M^A, A)]$ is the derivation of degree 0

$$\mathcal{L}_D = i_D \circ \delta_{M^A}^A + \delta_{M^A}^A \circ i_D : \Lambda(\Omega_A[C^\infty(M^A, A)]) \rightarrow \Lambda(\Omega_A[C^\infty(M^A, A)]).$$

Proposition 7. For any $\theta \in \mathfrak{X}(M)$, the map

$$\mathcal{L}_{\theta^A} : \Lambda(\Omega_A[C^\infty(M^A, A)]) \rightarrow \Lambda(\Omega_A[C^\infty(M^A, A)])$$

is a unique A -linear derivation such that

$$\mathcal{L}_{\theta^A}(\eta^A) = [\mathcal{L}_\theta(\eta)]^A,$$

for any $\eta \in \Lambda(\Omega_{\mathbb{R}}[C^\infty(M)])$.

Proof. For any $\eta \in \Lambda(\Omega_{\mathbb{R}}[C^\infty(M)])$, we have

$$\begin{aligned}\mathfrak{L}_{\theta^A}(\eta^A) &= i_{\theta^A}[\delta_{MA}^A(\eta^A)] + \delta_{MA}^A[i_{\theta^A}(\eta^A)] \\ &= i_{\theta^A}([\delta_M(\eta)]^A) + \delta_{MA}^A([i_\theta(\eta)]^A) \\ &= (i_\theta[\delta_M(\eta)])^A + (\delta_M[i_\theta(\eta)])^A \\ &= (i_\theta[\delta_M(\eta)] + \delta_M[i_\theta(\eta)])^A \\ &= [\mathfrak{L}_\theta(\eta)]^A.\end{aligned}$$

□

Proposition 8. For any $\theta \in \mathfrak{X}(M)$, for any $x \in \Omega_{\mathbb{R}}[C^\infty(M)]$ and for any $f \in C^\infty(M)$, we have

$$(1) \quad \mathfrak{L}_{f^A \cdot \theta^A}(x^A) = [\mathfrak{L}_{f \cdot \theta}(x)]^A.$$

$$(2) \quad \mathfrak{L}_{\theta^A}(f^A \cdot x^A) = [\mathfrak{L}_\theta(f \cdot x)]^A.$$

$$(3) \quad \mathfrak{L}_{\theta^A}[\delta_{MA}^A(f^A)] = [\mathfrak{L}_\theta(\delta_M(f))]^A.$$

Proof. For any $\theta \in \mathfrak{X}(M)$, for any $x \in \Omega_{\mathbb{R}}[C^\infty(M)]$ and for any $f \in C^\infty(M)$, we have

$$\begin{aligned}1. \quad \mathfrak{L}_{f^A \cdot \theta^A}(x^A) &= i_{f^A \cdot \theta^A}[\delta_{MA}^A(x^A)] + \delta_{MA}^A[i_{f^A \cdot \theta^A}(x^A)] \\ &= f^A \cdot i_{\theta^A}[\delta_{MA}^A(x^A)] + \delta_{MA}^A[f^A \cdot i_{\theta^A}(x^A)] \\ &= f^A \cdot i_{\theta^A}([\delta_M(x)]^A) + \delta_{MA}^A(f^A \cdot [i_\theta(x)]^A) \\ &= f^A \cdot i_{\theta^A}([\delta_M(x)]^A) + i_{\theta^A}(x^A) \cdot \delta_{MA}^A(f^A) + f^A \cdot \delta_{MA}^A[i_{\theta^A}(x^A)] \\ &= f^A \cdot (i_\theta[\delta_M(x)])^A + [i_\theta(x)]^A \cdot [\delta_M(f)]^A + f^A \cdot (\delta_M[i_\theta(x)])^A \\ &= (f \cdot i_\theta[\delta_M(x)] + i_\theta(x) \cdot \delta_M(f) + f \cdot \delta_M[i_\theta(x)])^A \\ &= (f \cdot i_\theta[\delta_M(x)] + \delta_M(f \cdot [i_\theta(x)]))^A \\ &= (i_{f \cdot \theta}[\delta_M(x)] + \delta_M[i_{f \cdot \theta}(x)])^A \\ &= [\mathfrak{L}_{f \cdot \theta}(x)]^A.\end{aligned}$$

Thus,

$$\mathfrak{L}_{f^A \cdot \theta^A}(x^A) = [\mathfrak{L}_{f \cdot \theta}(x)]^A.$$

$$\begin{aligned}2. \quad \mathfrak{L}_{\theta^A}(f^A \cdot x^A) &= i_{\theta^A}[\delta_{MA}^A(f^A \cdot x^A)] + \delta_{MA}^A[i_{\theta^A}(f^A \cdot x^A)] \\ &= i_{\theta^A}[\delta_{MA}^A(f^A)\Lambda x^A + f^A \cdot \delta_{MA}^A(x^A)] + \delta_{MA}^A[f^A \cdot i_{\theta^A}(x^A)] \\ &= \theta^A(f^A) \cdot x^A - \delta_{MA}^A(f^A) \cdot \widetilde{\theta^A}(x^A) + f^A \cdot \theta^A(x^A) \\ &\quad + \delta_{MA}^A(f^A) \cdot \widetilde{\theta^A}(x^A) + f^A \cdot \delta_{MA}^A[i_{\theta^A}(x^A)]\end{aligned}$$

$$\begin{aligned}
&= (\theta(f) \cdot x - \delta_M(f) \cdot \tilde{\theta}(x) + f \cdot \theta(x) \\
&\quad + \delta_M(f) \cdot \tilde{\theta}(x) + f \cdot \delta_M[i_\theta(x)])^A \\
&= (i_\theta[\delta_M(f)\Lambda x + f \cdot \delta_M(x)] + \delta_M[f \cdot i_\theta(x)])^A \\
&= (i_\theta[\delta_M(f \cdot x)] + \delta_M[i_\theta(f \cdot x)])^A \\
&= [\mathcal{L}_\theta(f \cdot x)]^A.
\end{aligned}$$

$$\begin{aligned}
3. \quad \mathcal{L}_{\theta^A}[\delta_{MA}^A(f^A)] &= i_{\theta^A}[\delta_{MA}^A(\delta_{MA}^A(f^A))] + \delta_{MA}^A[i_{\theta^A}(\delta_{MA}^A(f^A))] \\
&= 0 + \delta_{MA}^A[\theta^A(f^A)] \\
&= \delta_{MA}^A([\theta(f)]^A) \\
&= (\delta_M[\theta(f)])^A \\
&= (0 + \delta_M[\theta(f)])^A \\
&= (i_\theta[\delta_M(\delta_M(f))] + \delta_M[i_\theta(\delta_M(f))])^A \\
&= [\mathcal{L}_\theta(\delta_M(f))]^A.
\end{aligned}$$

□

Proposition 9. For any $D \in \text{Der}_A[C^\infty(M^A, A)]$, $X \in \Omega_A[C^\infty(M^A, A)]$, and $\varphi \in C^\infty(M^A, A)$, we have

$$(1) \quad \mathcal{L}_{\varphi \cdot D}(X) = \varphi \cdot \mathcal{L}_D(X) + \tilde{D}(X) \cdot \delta_{MA}^A(\varphi);$$

$$(2) \quad \mathcal{L}_D(\varphi \cdot X) = D(\varphi) \cdot X + \varphi \cdot \mathcal{L}_D(X);$$

$$(3) \quad \mathcal{L}_D[\delta_{MA}^A(\varphi)] = \delta_{MA}^A[D(\varphi)].$$

Proof. For any $D \in \text{Der}_A[C^\infty(M^A, A)]$, $X \in \Omega_A[C^\infty(M^A, A)]$, and $\varphi \in C^\infty(M^A, A)$, we have

$$\begin{aligned}
1. \quad \mathcal{L}_{\varphi \cdot D}(X) &= i_{\varphi \cdot D}[\delta_{MA}^A(X)] + \delta_{MA}^A[i_{\varphi \cdot D}(X)] \\
&= \varphi \cdot i_D[\delta_{MA}^A(X)] + \delta_{MA}^A[\varphi \cdot i_D(X)] \\
&= \varphi \cdot i_D[\delta_{MA}^A(X)] + i_D(X) \cdot \delta_{MA}^A(\varphi) + \varphi \cdot \delta_{MA}^A[i_D(X)] \\
&= \varphi \cdot \mathcal{L}_D(X) + \tilde{D}(X) \cdot \delta_{MA}^A(\varphi).
\end{aligned}$$

$$\begin{aligned}
2. \quad \mathfrak{L}_D(\varphi \cdot X) &= i_D[\delta_{M^A}^A(\varphi \cdot X)] + \delta_{M^A}^A[i_D(\varphi \cdot X)] \\
&= i_D[\delta_{M^A}^A(\varphi)\Lambda X + \varphi \cdot \delta_{M^A}^A(X)] + \delta_{M^A}^A[\varphi \cdot i_D(X)] \\
&= \tilde{D}[\delta_{M^A}^A(\varphi)] \cdot X - \delta_{M^A}^A(\varphi) \cdot \tilde{D}(X) + \varphi \cdot i_D[\delta_{M^A}^A(X)] \\
&\quad + \delta_{M^A}^A(\varphi) \cdot \tilde{D}(X) + \varphi \cdot \delta_{M^A}^A[i_D(X)] \\
&= D(\varphi) \cdot X + \varphi \cdot i_D[\delta_{M^A}^A(X)] + \varphi \cdot \delta_{M^A}^A[i_D(X)] \\
&= D(\varphi) \cdot X + \varphi \cdot (i_D[\delta_{M^A}^A(X)] + \delta_{M^A}^A[i_D(X)]) \\
&= D(\varphi) \cdot X + \varphi \cdot \mathfrak{L}_D(X).
\end{aligned}$$

$$\begin{aligned}
3. \quad \mathfrak{L}_D[\delta_{M^A}^A(\varphi)] &= i_D[\delta_{M^A}^A(\delta_{M^A}^A(\varphi))] + \delta_{M^A}^A[i_D(\delta_{M^A}^A(\varphi))] \\
&= 0 + \delta_{M^A}^A[\tilde{D} \circ \delta_{M^A}^A(\varphi)] \\
&= \delta_{M^A}^A[D(\varphi)].
\end{aligned}$$

□

4. THE POISSON 2-FORM ON WEIL BUNDLES

We recall that, when M is a smooth manifold, A a Weil algebra and M^A the associated Weil bundle, the A -algebra $C^\infty(M^A, A)$ is a Poisson algebra over A if there exists a bracket $\{\cdot, \cdot\}$ on $C^\infty(M^A, A)$ such that the pair $(C^\infty(M^A, A), \{\cdot, \cdot\})$ is a Lie algebra over A satisfying

$$\{\varphi, \psi_1 \cdot \psi_2\} = \{\varphi, \psi_1\} \cdot \psi_2 + \psi_1 \cdot \{\varphi, \psi_2\}$$

for any $\varphi, \psi_1, \psi_2 \in C^\infty(M^A, A)$. When $C^\infty(M^A, A)$ is a Poisson A -algebra, we will say that the manifold M^A is a A -Poisson manifold [1], [7].

When $(M, \{\cdot, \cdot\})$ is a Poisson manifold, the map

$$\text{ad}: C^\infty(M) \rightarrow \text{Der}_{\mathbb{R}}[C^\infty(M)], \quad f \mapsto \text{ad}(f)$$

such that $[\text{ad}(f)](g) = \{f, g\}$ for any $g \in C^\infty(M)$, is a derivation. Thus:

Proposition 10. *There exists a derivation*

$$\text{ad}^A: C^\infty(M^A, A) \rightarrow \text{Der}_A[C^\infty(M^A, A)]$$

such that

$$\text{ad}^A(f^A) = [\text{ad}(f)]^A.$$

Let's consider the following diagram commutative:

$$\begin{array}{ccc}
C^\infty(M^A, A) & \xrightarrow{\tilde{\tau}} & \text{Der}_A[C^\infty(M^A, A)] \\
\gamma_M \uparrow & & \uparrow \Phi \\
C^\infty(M) & \xrightarrow{\text{ad}} & \text{Der}_{\mathbb{R}}[C^\infty(M)]
\end{array}$$

i.e.

$$\tilde{\tau} \circ \gamma_M = \Phi \circ \text{ad},$$

where

$$\gamma_M: C^\infty(M) \rightarrow C^\infty(M^A, A), \quad f \mapsto f^A$$

and

$$\Phi: \text{Der}_{\mathbb{R}}[C^\infty(M)] \rightarrow \text{Der}_A[C^\infty(M^A, A)], \quad \theta \mapsto \theta^A.$$

For any $f \in C^\infty(M)$, we have

$$\tilde{\tau} \circ \gamma_M(f) = \tilde{\tau}(f^A)$$

and

$$\Phi \circ \text{ad}(f) = \Phi[\text{ad}(f)] = [\text{ad}(f)]^A.$$

Thus, there exists $\text{ad}^A = \tilde{\tau}$ such that

$$\text{ad}^A(f^A) = [\text{ad}(f)]^A.$$

As

$$\text{ad}^A: C^\infty(M^A, A) \rightarrow \text{Der}_A[C^\infty(M^A, A)]$$

is a derivation, then there exists a unique $C^\infty(M^A, A)$ -linear map

$$\widetilde{\text{ad}}^A: \Omega_A[C^\infty(M^A, A)] \rightarrow \text{Der}_A[C^\infty(M^A, A)]$$

such that

$$\widetilde{\text{ad}}^A \circ \delta_{M^A}^A = \text{ad}^A.$$

Let's consider the canonical isomorphism

$$\sigma_{M^A}: \Omega_A[C^\infty(M^A, A)]^* \longrightarrow \text{Der}_A[C^\infty(M^A, A)], \quad \Psi \longmapsto \Psi \circ \delta_{M^A}^A$$

and let

$$\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A: \Omega_A[C^\infty(M^A, A)] \xrightarrow{\widetilde{\text{ad}}^A} \text{Der}_A[C^\infty(M^A, A)] \xrightarrow{\sigma_{M^A}^{-1}} \Omega_A[C^\infty(M^A, A)]^*$$

be the map.

Proposition 11. *If (M, ω_M) is a Poisson manifold, then the map,*

$$\omega_{M^A}^A: \Omega_A[C^\infty(M^A, A)] \times \Omega_A[C^\infty(M^A, A)] \rightarrow C^\infty(M^A, A)$$

such that for any $X, Y \in \Omega_A[C^\infty(M^A, A)]$

$$\omega_{M^A}^A(X, Y) = -[\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(X)](Y)$$

is a skew-symmetric 2-form on $\Omega_A[C^\infty(M^A, A)]$ such that

$$\omega_{M^A}^A(x^A, y^A) = [\omega_M(x, y)]^A,$$

for any x and y in $\Omega_{\mathbb{R}}[C^\infty(M)]$.

Proof. For any $X \in \Omega_A[C^\infty(M^A, A)]$, we have $X = \sum_{i \in I: \text{fini}} \varphi_i \cdot \delta_{M^A}^A(\psi_i)$, with $\varphi_i \in C^\infty(M^A, A)$, $\psi_i \in C^\infty(M^A, A)$.

$$\begin{aligned}
\omega_{M^A}(X, X) &= -[\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(X)](X) \\
&= -\sum_{j \in I: \text{finite}} \varphi_j \cdot [\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(X)] \delta_{M^A}^A(\psi_j) \\
&= -\sum_{j \in I: \text{finite}} \varphi_j \cdot [\widetilde{\text{ad}}^A(X)](\psi_j) \\
&= -\sum_{j, k \in I: \text{finite}} \varphi_j \cdot \varphi_k \cdot [\widetilde{\text{ad}}^A(\delta_{M^A}^A(\psi_k))](\psi_j) \\
&= -\sum_{j, k \in I: \text{finite}} \varphi_j \cdot \varphi_k \cdot [\text{ad}^A(\psi_k)](\psi_j) \\
&= -\sum_{j, k \in I: \text{finite}} \varphi_j \cdot \varphi_k \cdot \{\psi_k, \psi_j\} \\
&= 0.
\end{aligned}$$

For any X_1 , X_2 and $Y \in \Omega_A[C^\infty(M^A, A)]$ and for any $\varphi \in C^\infty(M^A, A)$, we have

$$\begin{aligned}
\omega_{M^A}[(\varphi \cdot X_1 + X_2), Y] &= -[\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(\varphi \cdot X_1 + X_2)](Y) \\
&= -(\sigma_{M^A}^{-1}[\varphi \cdot \widetilde{\text{ad}}^A(X_1) + \widetilde{\text{ad}}^A(X_2)])(Y) \\
&= -\varphi \cdot (\sigma_{M^A}^{-1}[\widetilde{\text{ad}}^A(X_1)](Y) + (\sigma_{M^A}^{-1}[\widetilde{\text{ad}}^A(X_2)])(Y)) \\
&= \varphi \cdot \omega_{M^A}(X_1, Y) + \omega_{M^A}(X_2, Y).
\end{aligned}$$

For any x and y in $\Omega_{\mathbb{R}}[C^\infty(M)]$,

$$x^A = \sum_{i \in I: \text{fini}} f_i^A \cdot \delta_{M^A}^A(f_i'^A) \quad \text{and} \quad y^A = \sum_{j \in I: \text{fini}} g_j^A \cdot \delta_{M^A}^A(g_j'^A).$$

Thus,

$$\begin{aligned}
\omega_{M^A}^A(x^A, y^A) &= -[\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(x^A)](y^A) \\
&= -\left[\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A \left(\sum_{i \in I: \text{finite}} f_i^A \cdot \delta_{M^A}^A(f_i'^A) \right) \right] \left(\sum_{j \in I: \text{finite}} g_j^A \cdot \delta_{M^A}^A(g_j'^A) \right) \\
&= -\sum_{i, j \in I: \text{finite}} f_i^A \cdot g_j^A \cdot [\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(\delta_{M^A}^A(f_i'^A))] (\delta_{M^A}^A(g_j'^A))
\end{aligned}$$

$$\begin{aligned}
&= - \sum_{i,j \in I: \text{ finite}} f_i^A \cdot g_j^A \cdot [\sigma_{M^A}^{-1} \circ \widetilde{\text{ad}}^A(\delta_{M^A}^A(f_i'^A))] (\delta_{M^A}^A(g_j'^A)) \\
&= - \sum_{i,j \in I: \text{ finite}} f_i^A \cdot g_j^A \cdot [\sigma_{M^A}^{-1}(\widetilde{\text{ad}}^A \circ \delta_{M^A}^A(f_i'^A))] (\delta_{M^A}^A(g_j'^A)) \\
&= - \sum_{i,j \in I: \text{ finite}} f_i^A \cdot g_j^A \cdot [\sigma_{M^A}^{-1}(\text{ad}^A(f_i'^A))] (\delta_{M^A}^A(g_j'^A)) \\
&= - \sum_{i,j \in I: \text{ finite}} f_i^A \cdot g_j^A \cdot [\sigma_{M^A}^{-1}(\text{ad}(f_i')^A)] (\delta_M(g_j'))^A \\
&= - \left[\sum_{i,j \in I: \text{ finite}} f_i \cdot g_j \cdot [\sigma_M^{-1}(\text{ad}(f_i'))] (\delta_M(g_j')) \right]^A \\
&= - \left[\sum_{i,j \in I: \text{ finite}} f_i \cdot g_j \cdot [\sigma_M^{-1}(\widetilde{\text{ad}} \circ \delta_M(f_i'))] (\delta_M(g_j')) \right]^A \\
&= - \left[\sigma_M^{-1} \circ \widetilde{\text{ad}} \left(\sum_{i \in I: \text{ finite}} f_i \cdot \delta_M(f_i') \right) \left(\sum_{j \in I: \text{ finite}} g_j \cdot \delta_M(g_j') \right) \right]^A \\
&= [\omega_M(x, y)]^A.
\end{aligned}$$

□

Proposition 12. *When (M, ω_M) is a Poisson manifold of Poisson 2-form ω_M , then $(M^A, \omega_{M^A}^A)$ is a Poisson manifold.*

Proof. For any f and g in $C^\infty(M)$,

$$\begin{aligned}
\omega_{M^A}^A(\delta_{M^A}^A(f^A), \delta_{M^A}^A(g^A)) &= \omega_{M^A}^A([\delta_M(f)]^A, [\delta_M(g)]^A) \\
&= [\omega_M(\delta_M(f), \delta_M(g))]^A \\
&= -\{f, g\}_M^A,
\end{aligned}$$

and

$$\omega_{M^A}^A(x^A, y^A) = [\omega_M(x, y)]^A,$$

for any $x, y \in \Omega_{\mathbb{R}}[C^\infty(M)]$. We deduce that $(M^A, \omega_{M^A}^A)$ is a Poisson manifold. □

Theorem 3. *The manifold M^A is a Poisson manifold if and only if there exists a skew-symmetric 2-form*

$$\omega_{M^A}^A: \Omega_A[C^\infty(M^A, A)] \times \Omega_A[C^\infty(M^A, A)] \rightarrow C^\infty(M^A, A)$$

such that for any φ and ψ in $C^\infty(M^A, A)$,

$$\{\varphi, \psi\}_{M^A} = -\omega_{M^A}^A(\delta_{M^A}^A(\varphi), \delta_{M^A}^A(\psi))$$

defines a structure of A -Lie algebra over $C^\infty(M^A, A)$. Moreover, for any f and g in $C^\infty(M)$,

$$\{f^A, g^A\}_{M^A} = \{f, g\}_M^A.$$

Proof. Indeed, according to the previous proposition, the bracket

$$\{\varphi, \psi\}_{M^A} = -\omega_{M^A}^A(\delta_{M^A}^A(\varphi), \delta_{M^A}^A(\psi))$$

defines a structure of A -Lie algebra over $C^\infty(M^A, A)$. For any f and g in $C^\infty(M)$,

$$\{f^A, g^A\}_{M^A} = -\omega_{M^A}^A(\delta_{M^A}^A(f^A), \delta_{M^A}^A(g^A)) = \{f, g\}_M^A.$$

In this case, we will say that $\omega_{M^A}^A$ is the Poisson 2-form of the A -Poisson manifold M^A and we denote $(M^A, \omega_{M^A}^A)$ the A -Poisson manifold of Poisson 2-form $\omega_{M^A}^A$. \square

Proposition 13. *When (M, ω_M) is a Poisson manifold of Poisson 2-form ω_M , then for any $x, y \in \Omega_{\mathbb{R}}[C^\infty(M)]$ and for any $f, g \in C^\infty(M)$, we get*

$$(1) \quad [\widetilde{\text{ad}}^A(x^A)](f^A) = ([\widetilde{\text{ad}}(x)](f))^A.$$

$$(2) \quad [\widetilde{\text{ad}}^A(x^A)](y^A) = ([\widetilde{\text{ad}}(x)](y))^A.$$

$$(3) \quad \mathfrak{L}_{\widetilde{\text{ad}}^A[\delta_{M^A}^A(f^A)]}(g^A) = (\mathfrak{L}_{\widetilde{\text{ad}}[\delta_M(f)]}(g))^A.$$

Proof. 1. For any $x \in \Omega_{\mathbb{R}}[C^\infty(M)]$, $x^A = g^A \cdot \delta_{M^A}^A(h^A)$ with g and h in $C^\infty(M)$, and for any $f, g \in C^\infty(M)$, we have

$$\begin{aligned} [\widetilde{\text{ad}}^A(x^A)](f^A) &= [\widetilde{\text{ad}}^A(g^A \cdot \delta_{M^A}^A(h^A))](f^A) \\ &= [g^A \cdot \widetilde{\text{ad}}^A \circ \delta_{M^A}^A(h^A)](f^A) \\ &= [g^A \cdot \text{ad}^A(h^A)](f^A) \\ &= g^A \cdot [\text{ad}(h)]^A(f^A) \\ &= (g \cdot [\text{ad}(h)](f))^A \\ &= (g \cdot \widetilde{\text{ad}}[\delta_M(h)](f))^A \\ &= (\widetilde{\text{ad}}[g \cdot \delta_M(h)](f))^A \\ &= (\widetilde{\text{ad}}[g \cdot \delta_M(h)](f))^A \\ &= ([\widetilde{\text{ad}}(x)](f))^A. \end{aligned}$$

2. When $y \in \Omega_{\mathbb{R}}[C^\infty(M)]$, $y^A = g^A \cdot \delta_{M^A}^A(h^A)$ with g and h in $C^\infty(M)$

$$\begin{aligned} \widetilde{[\text{ad}^A(x^A)]}(y^A) &= \widetilde{[\text{ad}^A(x^A)]}(g^A \cdot \delta_{M^A}^A(h^A)) \\ &= g^A \cdot (\widetilde{[\text{ad}^A(x^A)]} \circ \delta_{M^A}^A)(h^A) \\ &= g^A \cdot \widetilde{[\text{ad}^A(x^A)]}(h^A) \\ &= g^A \cdot \widetilde{[\text{ad}^A(x^A)]}(h^A) \\ &= (g \cdot [\widetilde{\text{ad}}(x)](h))^A \\ &= (g \cdot [\widetilde{\text{ad}}(x)] \circ \delta_M)(h))^A \\ &= ([\widetilde{\text{ad}}(x)](g \cdot \delta_M(h)))^A \\ &= ([\widetilde{\text{ad}}(x)](y))^A. \end{aligned}$$

□

Proposition 14. *When (M, ω_M) is a Poisson manifold of Poisson 2-form ω_M , then for any $X, Y \in \Omega_A[C^\infty(M^A, A)]$ and for any $\varphi, \psi \in C^\infty(M^A, A)$, we get*

$$(1) \quad [\widetilde{\text{ad}}^A(X)](\varphi) = -\omega_{M^A}^A(X, \delta_{M^A}^A(\varphi));$$

$$(2) \quad [\widetilde{\text{ad}}^A(X)](Y) = -\omega_{M^A}^A(X, Y);$$

$$(3) \quad \mathfrak{L}_{\widetilde{\text{ad}}^A[\delta_{M^A}^A(\varphi)]} \delta_{M^A}^A(\psi) = \delta_{M^A}^A(\{\varphi, \psi\}_{M^A}).$$

Proof. When X and $Y \in \Omega_A[C^\infty(M^A, A)]$, $X = \sum_{i \in I: \text{fini}} \varphi_i \cdot \delta_{M^A}^A(\varphi'_i)$, $Y = \sum_{j \in J: \text{fini}} \psi_j \cdot \delta_{M^A}^A(\psi'_j)$ with $\varphi_i, \varphi'_i, \psi_j, \psi'_j \in C^\infty(M^A, A)$

$$\begin{aligned} 1. \quad [\widetilde{\text{ad}}^A(X)](\varphi) &= \left[\widetilde{\text{ad}}^A \left(\sum_{i \in I: \text{fini}} \varphi_i \cdot \delta_{M^A}^A(\varphi'_i) \right) \right] (\varphi) \\ &= \sum_{i \in I: \text{fini}} \varphi_i \cdot (\widetilde{\text{ad}}^A[\delta_{M^A}^A(\varphi'_i)])(\varphi) \\ &= \sum_{i \in I: \text{fini}} \varphi_i \cdot [\widetilde{\text{ad}}^A(\varphi'_i)](\varphi) \\ &= \sum_{i \in I: \text{fini}} \varphi_i \cdot \{\varphi'_i, \varphi\}_{M^A} \\ &= - \sum_{i \in I: \text{fini}} \varphi_i \cdot \omega_{M^A}^A(\delta_{M^A}^A(\varphi'_i), \delta_{M^A}^A(\varphi)) \\ &= -\omega_{M^A}^A \left(\sum_{i \in I: \text{fini}} \varphi_i \cdot \delta_{M^A}^A(\varphi'_i), \delta_{M^A}^A(\varphi) \right) \\ &= -\omega_{M^A}^A(X, \delta_{M^A}^A(\varphi)). \end{aligned}$$

$$\begin{aligned}
2. \quad & [\widetilde{\text{ad}}^A(X)](Y) = [\widetilde{\text{ad}}^A(X)]\left(\sum_{j \in J: \text{ fini}} \psi_j \cdot \delta_{M^A}^A(\psi'_j)\right) \\
&= \sum_{j \in J: \text{ fini}} \psi_j \cdot [\widetilde{\text{ad}}^A(X)](\delta_{M^A}^A(\psi'_j)) \\
&= \sum_{j \in J: \text{ fini}} \psi_j \cdot ([\widetilde{\text{ad}}^A(X)] \circ \delta_{M^A}^A)(\psi'_j) \\
&= \sum_{j \in J: \text{ fini}} \psi_j \cdot [\widetilde{\text{ad}}^A(X)](\psi'_j) \\
&= - \sum_{j \in J: \text{ fini}} \psi_j \cdot \omega_{M^A}^A(X, \delta_{M^A}^A(\psi'_j)) \\
&= -\omega_{M^A}^A\left(X, \sum_{j \in J: \text{ fini}} \psi_j \cdot \delta_{M^A}^A(\psi'_j)\right) \\
&= -\omega_{M^A}^A(X, Y);
\end{aligned}$$

3. $\mathfrak{L}_{\widetilde{\text{ad}}^A[\delta_{M^A}^A(\varphi)]}\delta_{M^A}^A(\psi) = \mathfrak{L}_{\text{ad}^A(\varphi)}\delta_{M^A}^A(\psi)$

$$\begin{aligned}
&= \delta_{M^A}^A[\text{ad}^A(\varphi)(\psi)] \\
&= \delta_{M^A}^A(\{\varphi, \psi\}_{M^A}).
\end{aligned}$$

□

REFERENCES

- [1] Bossoto, B.G.R., Okassa, E., *Champs de vecteurs et formes différentielles sur une variété des points proches*, Arch. Math. (Brno) **44** (2008), 159–171.
- [2] Bossoto, B.G.R., Okassa, E., *A-poisson structures on Weil bundles*, Int. J. Contemp. Math. Sci. **7** (16) (2012), 785–803.
- [3] Kolář, I., Michor, P.W., Slovák, J., *Natural Operations in Differential Geometry*, Springer, Berlin, 1993.
- [4] Laurent-Gengoux, C., Pichereau, A., Vanhaecke, P., *Poisson Structures*, Grundlehren Math. Wiss. **347** (2013), www.springer.com/series/138.
- [5] Lichnerowicz, A., *Les variétés de Poisson et leurs algèbres de Lie associées*, J. Differential Geom. **12** (1977), 253–300.
- [6] Morimoto, A., *Prolongation of connections to bundles of infinitely near points*, J. Differential Geom. **11** (1976), 479–498.
- [7] Moukala, N.M., Bossoto, B.G.R., *Hamiltonian vector fields on Weil bundles*, Journal of Mathematics Research **7** (3) (2015), 141–148.
- [8] Nkou, V.B., Bossoto, B.G.R., Okassa, E., *New characterization of vector field on Weil bundles*, Theoretical Mathematics & Applications **5** (2) (2015), 1–17, [arXiv:1504.04483 \[math.DG\]](https://arxiv.org/abs/1504.04483).
- [9] Okassa, E., *Prolongement des champs de vecteurs à des variétés des points proches*, Ann. Fac. Sci. Toulouse Math. (5) **8** (3) (1986–1987), 346–366.
- [10] Okassa, E., *Algèbres de Jacobi et algèbres de Lie-Rinehart-Jacobi*, J. Pure Appl. Algebra **208** (3) (2007), 1071–1089.

- [11] Okassa, E., *On Lie-Rinehart-Jacobi algebras*, J. Algebra Appl. **7** (2008), 749–772.
- [12] Okassa, E., *Symplectic Lie-Rinehart-Jacobi algebras and contact manifolds*, Canad. Math. Bull. **54** (4) (2011), 716–725.
- [13] Shurygin, V.V., *Some aspects of the theory of manifolds over algebras and of Weil bundles*, J. Math. Sci. (New York) **169** (3) (2010), 315–341.
- [14] Vaisman, I., *Lectures on the Geometry of Poisson Manifolds*, Progress in Math., vol. 118, Birkhäuser Verlag, Basel, 1994.
- [15] Weil, A., *Théorie des points proches sur les variétés différentiables*, Colloques Internationaux du Centre National de la Recherche Scientifique, Strasbourg (1953), 111–117.

MARIEN NGOUABI UNIVERSITY,

BP:69, BRAZZAVILLE, CONGO

AND

INSTITUT DE RECHERCHE EN SCIENCES EXACTES ET NATURELLES (IRSEN),

BRAZZAVILLE, CONGO

E-mail: bossotob@yahoo.fr nmahomouk@yahoo.fr