

Bilender P. Allahverdiev; Hüseyin Tuna

Spectral Theory of Singular Hahn Difference Equation of the Sturm-Liouville Type

Communications in Mathematics, Vol. 28 (2020), No. 1, 13–25

Persistent URL: <http://dml.cz/dmlcz/148258>

Terms of use:

© University of Ostrava, 2020

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Spectral Theory of Singular Hahn Difference Equation of the Sturm-Liouville Type

Bilender P. Allahverdiev, Hüseyin Tuna

Abstract. In this work, we consider the singular Hahn difference equation of the Sturm-Liouville type. We prove the existence of the spectral function for this equation. We establish Parseval equality and an expansion formula for this equation on a semi-unbounded interval.

1 Introduction

Spectral expansion theorems have attracted mathematicians for a long time. The first results of this type go back to Weyl [37]. Additional results were obtained by Stone [34], [35], Naimark [31], Berezanskii [13] and Titchmarsh [36]. Usually, if we want to solve a partial differential equation using the Fourier method (i.e., the separation of variables) then we consider the problem of expanding an arbitrary function as a series of eigenfunctions. Hence the eigenfunction expanding problem has been studied extensively in the literature (see [2], [3], [4], [5], [6], [7], [13], [15], [16], [21], [22], [29], [30], [31], [34], [35], [36], [38], [39]).

The study of the Hahn difference operator dates back to Hahn's works [17] and [18]. In 1949, Hahn introduced the quantum difference operator $D_{\omega,q}$ defined by

$$D_{\omega,q}f(x) = \frac{f(\omega + qx) - f(x)}{\omega + (q - 1)x},$$

where $q \in (0, 1)$ and $\omega > 0$ (see [17], [18]). The Hahn difference operator $D_{\omega,q}$ is a generalization of the two well-known difference operators; namely, the quantum

2020 MSC: 39A13, 39A12, 39A70, 34L10, 34B40

Key words: Hahn's Sturm-Liouville equation, spectral function, Parseval equality, spectral expansion.

Affiliation:

Bilender P. Allahverdiev – Department of Mathematics, Süleyman Demirel University, 32260 Isparta, Turkey

E-mail: bilenderpasaoglu@sdu.edu.tr

Hüseyin Tuna – Department of Mathematics, Mehmet Akif Ersoy University, 15030 Burdur, Turkey

E-mail: hustuna@gmail.com

q -difference operator (see [23]) and the forward difference operator (see [24], [25]). This operator has numerous applications in the construction of families of orthogonal polynomials and approximation problems (see [8], [12], [14], [27], [28], [32]). A proper inverse of $D_{\omega,q}$ and the associated integral calculus were studied in [1], [9]. Next, in [19], Hamza et al. established the theory of linear Hahn difference equations. They investigated also the existence and uniqueness of the solution of initial value problems for Hahn difference equations. Moreover, they obtained Gronwall's and Bernoulli's inequalities with respect to the Hahn difference operator, and investigated the mean value theorems for this calculus. Later, Hamza and Makharesh [20] studied Leibniz's rule and Fubini's theorem associated with the Hahn difference operator. Sitthiwirattam [33] investigated the nonlocal boundary value problem for nonlinear Hahn difference equation. Recently, in [10], Annaby et al. established a Sturm-Liouville theory associated with the Hahn difference operator in the regular setting. In [11], the authors introduce a couple of sampling theorems of Lagrange-type interpolation for ω, q -integral transforms whose kernels are either solutions or Green's function of the ω, q -Hahn-Sturm-Liouville problem.

In this paper, we study Hahn difference equations of the Sturm-Liouville type. Having the solutions of such equations we define the new Hilbert space and construct on it the Fourier transform and prove the Parseval equation. Therefore, we prove the existence of a spectral function for the Hahn difference equations of the Sturm-Liouville type in Lemma 3. In Theorem 5, a Parseval equality and an expansion formula in eigenfunctions are established in terms of this spectral function.

2 Notation and basic results

In this section, our aim is to present some basic concepts concerning the theory of Hahn calculus. For more details, the reader may refer to [9], [10], [17] and [18]. Throughout the paper, let $q \in (0, 1)$ and $\omega > 0$.

Let $\omega_0 := \omega/(1 - q)$ and let I be a real interval containing ω_0 .

Definition 1 ([17], [18]). Let $f: I \rightarrow \mathbb{R} := (-\infty, \infty)$ be a function. The Hahn difference operator is defined by

$$D_{\omega,q}f(x) = \begin{cases} \frac{f(\omega + qx) - f(x)}{\omega + (q-1)x}, & x \neq \omega_0, \\ f'(\omega_0), & x = \omega_0, \end{cases} \quad (1)$$

provided that f is differentiable at ω_0 . We call $D_{\omega,q}f$ a ω, q -derivative of f .

Remark 1. The Hahn difference operator unifies two well known operators. When $q \rightarrow 1$, we get the forward difference operator which is defined by

$$\Delta_{\omega}f(x) := \frac{f(\omega + x) - f(x)}{(\omega + x) - x}, \quad x \in \mathbb{R}.$$

When $\omega \rightarrow 0$, we get the Jackson q -difference operator which is defined by

$$D_qf(x) := \frac{f(qx) - f(x)}{(qx) - x}, \quad x \neq 0.$$

Furthermore, under appropriate conditions, we have

$$\lim_{\substack{q \rightarrow 1 \\ \omega \rightarrow 0}} D_{\omega, q} f(x) = f'(x).$$

In what follows, we present several important properties of the ω, q -derivative.

Theorem 1 ([9]). *Let $f, g: I \rightarrow \mathbb{R}$ be ω, q -differentiable at $x \in I$ and $h(x) := \omega + qx$. Then, for all $x \in I$ we have*

- i) $D_{\omega, q}(af + bg)(x) = aD_{\omega, q}f(x) + bD_{\omega, q}g(x), \quad a, b \in I,$
- ii) $D_{\omega, q}(fg)(x) = D_{\omega, q}(f(x))g(x) + f(\omega + qx)D_{\omega, q}g(x),$
- iii) $D_{\omega, q}\left(\frac{f}{g}\right)(x) = \frac{D_{\omega, q}(f(x))g(x) - f(x)D_{\omega, q}g(x)}{g(x)g(\omega + qx)},$
- iv) $D_{\omega, q}(h^{-1}(x)) = D_{-\omega q^{-1}, q^{-1}}f(x).$

The concept of the ω, q -integral of the function f can be defined as follows.

Definition 2 (Jackson-Nörlund Integral [9]). Let $f: I \rightarrow \mathbb{R}$ be a function and $a, b, \omega_0 \in I$. We define the ω, q -integral of the function f from a to b by

$$\int_a^b f(x) d_{\omega, q}(x) := \int_{\omega_0}^b f(x) d_{\omega, q}(x) - \int_{\omega_0}^a f(x) d_{\omega, q}(x),$$

where

$$\int_{\omega_0}^x f(t) d_{\omega, q}(t) := ((1-q)x - \omega) \sum_{n=0}^{\infty} q^n f\left(\omega \frac{1-q^n}{1-q} + xq^n\right), \quad x \in I$$

provided that the series converges at $x = a$ and $x = b$. In this case, f is called ω, q -integrable on $[a, b]$.

Similarly, one can define the ω, q -integral of a function f over (ω_0, ∞) by

$$\int_{\omega_0}^{\infty} f(x) d_{\omega, q}(x) := \lim_{b \rightarrow \infty} \int_{\omega_0}^b f(x) d_{\omega, q}(x).$$

The following properties of ω, q -integration can be found in [9].

Lemma 1 ([9]). *Let $f, g: I \rightarrow \mathbb{R}$ be ω, q -integrable on I and let $a, b, c \in I$ where $a < c < b$ and $\alpha, \beta \in \mathbb{R}$. Then the following formulae hold:*

- i) $\int_a^b (\alpha f(x) + \beta g(x)) d_{\omega, q}(x) = \alpha \int_a^b f(x) d_{\omega, q}(x) + \beta \int_a^b g(x) d_{\omega, q}(x),$
- ii) $\int_a^a f(x) d_{\omega, q}(x) = 0,$

$$\text{iii) } \int_a^b f(x) d_{\omega,q}(x) = \int_a^c f(x) d_{\omega,q}(x) + \int_c^b f(x) d_{\omega,q}(x),$$

$$\text{iv) } \int_a^b f(x) d_{\omega,q}(x) = - \int_b^a f(x) d_{\omega,q}(x).$$

Next, we present the ω, q -integration by parts.

Lemma 2 ([9]). *Let $f, g: I \rightarrow \mathbb{R}$ be ω, q -integrable on I and let $a, b \in I$ where $a < b$. Then the following formula holds:*

$$\int_a^b f(x) D_{\omega,q}g(x) d_{\omega,q}(x) + \int_a^b g(\omega + qx) D_{\omega,q}f(x) d_{\omega,q}(x) = f(b)g(b) - f(a)g(a).$$

The next result is the fundamental theorem of Hahn calculus.

Theorem 2 ([9]). *Let $f: I \rightarrow \mathbb{R}$ be continuous at ω_0 . Define*

$$F(x) := \int_{\omega_0}^x f(t) d_{\omega,q}(t), \quad x \in I.$$

Then F is continuous at ω_0 . Moreover, $D_{\omega,q}F(x)$ exists for every $x \in I$ and $D_{\omega,q}F(x) = f(x)$. Conversely,

$$\int_a^b D_{\omega,q}F(x) d_{\omega,q}(x) = F(b) - F(a).$$

Let $L_{\omega,q}^2(\omega_0, \infty)$ be the space of all complex-valued functions defined on $[\omega_0, \infty)$ such that

$$\|f\| := \left(\int_{\omega_0}^{\infty} |f(x)|^2 d_{\omega,q}x \right)^{1/2} < \infty.$$

The space $L_{\omega,q}^2(\omega_0, \infty)$ is a separable Hilbert space with the inner product

$$(f, g) := \int_{\omega_0}^{\infty} f(x) \overline{g(x)} d_{\omega,q}x, \quad f, g \in L_{\omega,q}^2(\omega_0, \infty)$$

(see [9]).

The ω, q -Wronskian of $y(\cdot), z(\cdot)$ is defined by

$$W_{\omega,q}(y, z)(x) := y(x)D_{\omega,q}z(x) - z(x)D_{\omega,q}y(x), \quad x \in [\omega_0, \infty). \quad (2)$$

Now we recall the following well-known theorems of Helly.

Theorem 3 ([26]). *Let $(w_n)_{n \in \mathbb{N}}$ ($\mathbb{N} := \{1, 2, \dots\}$) be a uniformly bounded sequence of real nondecreasing functions on a finite interval $a \leq \lambda \leq b$. Then there exists a subsequence $(w_{n_k})_{k \in \mathbb{N}}$ and a nondecreasing function w such that*

$$\lim_{k \rightarrow \infty} w_{n_k}(\lambda) = w(\lambda), \quad a \leq \lambda \leq b.$$

Theorem 4 ([26]). Assume $(w_n)_{n \in \mathbb{N}}$ is a real, uniformly bounded sequence of non-decreasing functions on a finite interval $a \leq \lambda \leq b$, and suppose

$$\lim_{n \rightarrow \infty} w_n(\lambda) = w(\lambda), \quad a \leq \lambda \leq b.$$

If f is any continuous function on $a \leq \lambda \leq b$, then

$$\lim_{n \rightarrow \infty} \int_a^b f(\lambda) dw_n(\lambda) = \int_a^b f(\lambda) dw(\lambda).$$

3 Main Results

Let us consider the Hahn difference equations of the Sturm-Liouville type given by

$$\Gamma(y) := -q^{-1}D_{-\omega q^{-1}, q^{-1}}D_{\omega, q}y(x) + v(x)y(x) = \lambda y(x), \quad x \in (\omega_0, \infty) \quad (3)$$

with the boundary condition

$$D_{-\omega q^{-1}, q^{-1}}y(\omega_0) \sin \beta + y(\omega_0) \cos \beta = 0, \quad \beta \in \mathbb{R}, \quad (4)$$

where λ is a complex eigenvalue parameter, v is a real-valued continuous function at ω_0 defined on $[\omega_0, \infty)$.

If we endow the problem (3)–(4) with the boundary condition

$$D_{-\omega q^{-1}, q^{-1}}y(q^{-n}) \sin \alpha + y(q^{-n}) \cos \alpha = 0, \quad \alpha \in \mathbb{R}, n \in \mathbb{N}, \quad (5)$$

then we deduce that the problem given by (3), (4) and (5) is a regular problem.

In [10], Annaby et al. showed that the boundary value problem (3) with the boundary conditions (4) and (5) has a compact resolvent, thus this problem has a purely discrete spectrum.

Let $\varphi(x)$ be a solution of the system (3) satisfying the initial conditions

$$\varphi(\omega_0) = \sin \beta, \quad D_{-\omega q^{-1}, q^{-1}}\varphi(\omega_0) = -\cos \beta. \quad (6)$$

Let ϱ be any nondecreasing function on \mathbb{R} . Denote by $L^2_{\varrho}(\mathbb{R})$ the Hilbert space of all functions $f: \mathbb{R} \rightarrow \mathbb{R}$ measurable with respect to the Lebesgue-Stieltjes measure defined by ϱ and such that

$$\int_{-\infty}^{\infty} f^2(\lambda) d\varrho(\lambda) < \infty,$$

with the inner product

$$(f, g)_{\varrho} := \int_{-\infty}^{\infty} f(\lambda)g(\lambda) d\varrho(\lambda).$$

The main result of this section is the following.

Theorem 5. *There exists a nondecreasing function $\varrho(\lambda)$ on $-\infty < \lambda < \infty$, a spectral function for the boundary value problem given by (3)–(4), with the following properties.*

i) If f is a real-valued function and $f \in L^2_{\omega, q}(\omega_0, q^{-n})$, then there exists a function $F \in L^2_{\varrho}(\mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} \left\{ F(\lambda) - \int_{\omega_0}^{q^{-n}} f(x) \varphi(x, \lambda) d_{\omega, q} x \right\} d\varrho(\lambda) = 0, \quad (7)$$

and the Parseval equality

$$\int_{\omega_0}^{\infty} f^2(x) d_{\omega, q} x = \int_{-\infty}^{\infty} F^2(\lambda) d\varrho(\lambda) \quad (8)$$

holds.

ii) The integral

$$\int_{-\infty}^{\infty} F(\lambda) \varphi(x, \lambda) d\varrho(\lambda)$$

converges to f in $L^2_{\omega, q}(\omega_0, \infty)$. That is,

$$\lim_{n \rightarrow \infty} \int_{\omega_0}^{q^{-n}} \left\{ f(x) - \int_{-\infty}^n F(\lambda) \varphi(x, \lambda) d\varrho(\lambda) \right\}^2 d_{\omega, q} x = 0. \quad (9)$$

Remark 2. The expression

$$f(x) = \int_{-\infty}^{\infty} F(\lambda) \varphi(x, \lambda) d\varrho(\lambda),$$

where the equality is in the sense of (9), is called the expansion theorem.

Let $\lambda_{m, q^{-n}}$ (where $m, n \in \mathbb{N}$) denote the eigenvalues of the regular problem given by (3)–(5), and $\varphi(x, \lambda)$ be the solution of the equation (3) satisfying the initial conditions in (6). The function $\varphi_{m, q^{-n}}(x) = \varphi(x, \lambda_{m, q^{-n}})$ will be an eigenfunction corresponding to the eigenvalue $\lambda_{m, q^{-n}}$.

Let $f(\cdot)$ be an arbitrary real-valued function on $L^2_{\omega, q}(\omega_0, q^{-n})$ and

$$\alpha_{m, q^{-n}}^2 = \int_{\omega_0}^{q^{-n}} \varphi_{m, q^{-n}}^2(x) d_{\omega, q} x$$

where $m \in \mathbb{N}$. Then we have

$$\int_{\omega_0}^{q^{-n}} f^2(x) d_{\omega, q} x = \sum_{m=1}^{\infty} \frac{1}{\alpha_{m, q^{-n}}^2} \left\{ \int_{\omega_0}^{q^{-n}} f(x) \varphi_{m, q^{-n}}(x) d_{\omega, q} x \right\}^2, \quad (10)$$

which is called the *Parseval equality* (see [10]).

Now we will introduce the nondecreasing step function $\varrho_{q^{-n}}$ on $[0, \infty)$ by

$$\varrho_{q^{-n}}(\lambda) = \begin{cases} - \sum_{\lambda < \lambda_{m, q^{-n}} < 0} \frac{1}{\alpha_{m, q^{-n}}^2}, & \text{for } \lambda \leq 0, \\ \sum_{0 \leq \lambda_{m, q^{-n}} < \lambda} \frac{1}{\alpha_{m, q^{-n}}^2}, & \text{for } \lambda \geq 0. \end{cases}$$

Then the equality (10) can be written as

$$\int_{\omega_0}^{q^{-n}} f^2(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F^2(\lambda) d_{\varrho_{q^{-n}}}(\lambda), \quad (11)$$

where

$$F(\lambda) = \int_{\omega_0}^{q^{-n}} f(x)\varphi(x, \lambda) d_{\omega, q}x.$$

We will show that the Parseval equality for the problem given by (3) and (4) can be obtained from (11) by letting $n \rightarrow \infty$. To this end, we shall prove the following lemma.

Lemma 3. *The total variation of the functions $\varrho_{q^{-n}}(\lambda)$ is uniformly bounded with respect to q^{-n} in each finite interval in the domain λ , i.e., for $s > 0$, there exists a constant $M = M(s) > 0$ not depending on q^{-n} such that*

$$\bigvee_{-s}^s \{\varrho_{q^{-n}}(\lambda)\} = \sum_{-s \leq \lambda_{m, q^{-n}} < s} \frac{1}{\alpha_{m, q^{-n}}^2} = \varrho_{q^{-n}}(s) - \varrho_{q^{-n}}(-s) < M. \quad (12)$$

Proof. Let $\sin \beta \neq 0$. Since $\varphi(x, \lambda)$ is continuous at ω_0 , we deduce from the condition $\varphi(\omega_0, \lambda) = \sin \beta$ that there exists a positive number k such that $k - \omega_0$ is so small and

$$\frac{1}{k} \left(\int_{\omega_0}^k \varphi(x, \lambda) d_{\omega, q}x \right)^2 > \frac{1}{2} \sin^2 \beta. \quad (13)$$

Let us define $f_k(x)$ by the formula

$$f_k(x) = \begin{cases} \frac{1}{k}, & \omega_0 \leq x < k \\ 0, & x > k. \end{cases}$$

By virtue of (11) and (13), we conclude that

$$\begin{aligned} \int_{\omega_0}^k f_k^2(x) d_{\omega, q}x &= \frac{1}{k} = \int_{-\infty}^{\infty} \left(\frac{1}{k} \int_{\omega_0}^k \varphi(x, \lambda) d_{\omega, q}x \right)^2 d_{\varrho_{q^{-n}}}(\lambda) \\ &\geq \int_{-s}^s \left(\frac{1}{k} \int_{\omega_0}^k \varphi(x, \lambda) d_{\omega, q}x \right)^2 d_{\varrho_{q^{-n}}}(\lambda) \\ &> \frac{1}{2} \sin^2 \beta \int_{-s}^s d_{\varrho_{q^{-n}}}(\lambda) \\ &= \frac{1}{2} \sin^2 \beta \{ \varrho_{q^{-n}}(s) - \varrho_{q^{-n}}(-s) \}, \end{aligned}$$

which is the desired result.

If $\sin \beta = 0$, then we can define the function $f_k(x)$ by the formula

$$f_k(x) = \begin{cases} \frac{1}{k^2}, & \omega_0 \leq x < k \\ 0, & x > k. \end{cases}$$

By virtue of (10), we get the inequality (12). □

Proof of Theorem 5. Suppose that the function $f_\xi(x)$ satisfies the following conditions:

- i) $f_\xi(x)$ vanishes outside the interval $[\omega_0, q^{-\xi}]$, $q^{-\xi} < q^{-n}$.
- ii) The functions $f_\xi(x)$ and $D_{\omega, q}f_\xi(x)$ are continuous on $[\omega_0, \infty)$.
- iii) $f_\xi(x)$ satisfies the boundary condition (4).

Applying (11) to $f_\xi(x)$, we deduce that

$$\int_{\omega_0}^{q^{-\xi}} f_\xi^2(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F^2(\lambda) d\rho(\lambda), \quad (14)$$

where

$$F(\lambda) = \int_{\omega_0}^{q^{-\xi}} f_\xi(x) \varphi(x, \lambda) d_{\omega, q}x. \quad (15)$$

Since $\varphi(x, \lambda)$ is a solution of the equation (3), we have

$$\varphi(x, \lambda) = \frac{1}{\lambda} \Gamma(\varphi(x, \lambda)).$$

By (15), we obtain

$$F_n(\lambda) = \frac{1}{\lambda} \int_{\omega_0}^{q^{-n}} f_\xi(x) \Gamma(\varphi(x, \lambda)) d_{\omega, q}x.$$

Since $f_\xi(x)$ vanishes in a neighbourhood of the point q^{-n} , and since $f_\xi(x)$ and $\varphi(x, \lambda)$ satisfy the boundary condition (6), by using Lemma 2 we obtain that

$$F_n(\lambda) = \frac{1}{\lambda} \int_{\omega_0}^{q^{-n}} \varphi(x, \lambda) \Gamma(f_\xi(x)) d_{\omega, q}x.$$

By virtue of (11), for any finite $s > 0$, we conclude that

$$\begin{aligned} \int_{|\lambda|>s} F_n^2(\lambda) d\rho_{q^{-n}}(\lambda) &\leq \frac{1}{s^2} \int_{|\lambda|>s} \left\{ \int_{\omega_0}^{q^{-n}} \varphi(x, \lambda) \Gamma(f_\xi(x)) d_{\omega, q}x \right\}^2 d\rho_{q^{-n}}(\lambda) \\ &\leq \frac{1}{s^2} \int_{-\infty}^{\infty} \left\{ \int_{\omega_0}^{q^{-n}} \varphi(x, \lambda) \Gamma(f_\xi(x)) d_{\omega, q}x \right\}^2 d\rho_{q^{-n}}(\lambda) \\ &= \frac{1}{s^2} \int_{\omega_0}^{q^{-\xi}} [\Gamma(f_\xi(x))]^2 d_{\omega, q}x. \end{aligned}$$

By the formula (14), we deduce that

$$\begin{aligned} &\left| \int_{\omega_0}^{q^{-\xi}} f_\xi^2(x) d_{\omega, q}x - \int_{-s}^s F_n^2(\lambda) d\rho_{q^{-n}}(\lambda) \right| \\ &< \frac{1}{s^2} \int_{\omega_0}^{q^{-\xi}} [-q^{-1} D_{-\omega q^{-1}, q^{-1}} D_{\omega, q} f_\xi(x) + v(x) f_\xi(x)]^2 d_{\omega, q}x. \quad (16) \end{aligned}$$

From Lemma 3, it follows that the set $\{\varrho_{q^{-n}}(\lambda)\}$ is bounded. Using Theorems 3 and 4, one can find a sequence $\{q^{-n_k}\}$ such that the sequence $\varrho_{q^{-n_k}}(\lambda)$ converges to a monotone function $\varrho(\lambda)$ as $k \rightarrow \infty$. Passing to the limit with respect to $\{q^{-n_k}\}$ in (16), we conclude that

$$\left| \int_{\omega_0}^{q^{-\varepsilon}} f_{\xi}^2(x) d_{\omega, q}x - \int_{-s}^s F_n^2(\lambda) d\varrho(\lambda) \right| < \frac{1}{s^2} \int_{\omega_0}^{q^{-\varepsilon}} [\Gamma(f_{\xi}(x))]^2 d_{\omega, q}x.$$

Letting $s \rightarrow \infty$, we see that

$$\int_{\omega_0}^{q^{-\varepsilon}} f_{\xi}^2(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F_n^2(\lambda) d\varrho(\lambda).$$

Now, let f be an arbitrary real-valued function on $L_{\omega, q}^2(\omega_0, \infty)$. It is known that there exists a sequence $\{f_{\xi}(x)\}$ satisfying the conditions i)–iii) and such that

$$\lim_{\xi \rightarrow \infty} \int_{\omega_0}^{\infty} (f(x) - f_{\xi}(x))^2 d_{\omega, q}x = 0.$$

Let

$$F_{\xi}(\lambda) = \int_{\omega_0}^{\infty} f_{\xi}(x) \varphi(x, \lambda) d_{\omega, q}x.$$

Then, we have

$$\int_{\omega_0}^{\infty} f_{\xi}^2(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F_{\xi}^2(\lambda) d\varrho(\lambda).$$

Since

$$\int_{\omega_0}^{\infty} (f_{\xi_1}(x) - f_{\xi_2}(x))^2 d_{\omega, q}x \rightarrow 0 \quad \text{as } \xi_1, \xi_2 \rightarrow \infty,$$

we have

$$\int_{-\infty}^{\infty} (F_{\xi_1}(\lambda) - F_{\xi_2}(\lambda))^2 d\varrho(\lambda) = \int_{\omega_0}^{\infty} (f_{\xi_1}(x) - f_{\xi_2}(x))^2 d_{\omega, q}x \rightarrow 0$$

as $\xi_1, \xi_2 \rightarrow \infty$. It follows from the completeness of the space $L_{\varrho}^2(\mathbb{R})$ that there exists a limit function F which satisfies

$$\int_{\omega_0}^{\infty} f^2(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F^2(\lambda) d\varrho(\lambda).$$

Our next aim is to prove that the function

$$K_{\xi}(\lambda) = \int_{\omega_0}^{q^{-\varepsilon}} f(x) \varphi(x, \lambda) d_{\omega, q}x$$

converges to F as $\xi \rightarrow \infty$, in the metric of the space $L_{\varrho}^2(\mathbb{R})$. Let g be another function in $L_{\omega, q}^2(\omega_0, \infty)$. Similarly, let $G(\lambda)$ be defined by g . It is evident that

$$\int_{\omega_0}^{\infty} (f(x) - g(x))^2 d_{\omega, q}x = \int_{-\infty}^{\infty} \{F(\lambda) - G(\lambda)\}^2 d\varrho(\lambda).$$

We define

$$g(x) = \begin{cases} f(x), & x \in [\omega_0, q^{-\xi}] \\ 0, & x \in (q^{-\xi}, \infty). \end{cases}$$

Then we obtain that

$$\int_{-\infty}^{\infty} \{F(\lambda) - K_{\xi}(\lambda)\}^2 d\rho(\lambda) = \int_{q^{-\xi}}^{\infty} f^2(x) d_{\omega, q}x \rightarrow 0 \quad (\xi \rightarrow \infty),$$

i.e., K_{ξ} converges to F in $L^2_{\rho}(\mathbb{R})$ as $\xi \rightarrow \infty$. This proves i).

Now, we will prove ii). Suppose that the real-valued functions f, g are in $L^2_{\omega, q}(\omega_0, q^{-n})$; and $F(\lambda)$ and $G(\lambda)$ are their Fourier transforms, respectively. Then $F \pm G$ are the transforms of $f \pm g$. Consequently, in view of (8), we have

$$\begin{aligned} \int_{\omega_0}^{\infty} [f(x) + g(x)]^2 d_{\omega, q}x &= \int_{-\infty}^{\infty} (F(\lambda) + G(\lambda))^2 d\rho(\lambda), \\ \int_{\omega_0}^{\infty} [f(x) - g(x)]^2 d_{\omega, q}x &= \int_{-\infty}^{\infty} (F(\lambda) - G(\lambda))^2 d\rho(\lambda). \end{aligned}$$

Subtracting the second relation from the first one, we deduce that

$$\int_{\omega_0}^{\infty} f(x)g(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F(\lambda)G(\lambda) d\rho(\lambda). \quad (17)$$

We set

$$f_{\tau}(x) = \int_{-\tau}^{\tau} F(\lambda)\varphi(x, \lambda) d\rho(\lambda),$$

where F is the function defined in (7). Let $g(\cdot)$ be a real-valued function which is equal to zero outside the finite interval $[\omega_0, q^{-\mu}]$. Thus we obtain

$$\begin{aligned} \int_{\omega_0}^{q^{-\mu}} f_{\tau}(x)g(x) d_{\omega, q}x &= \int_{\omega_0}^{q^{-\mu}} \left\{ \int_{-\tau}^{\tau} F(\lambda)\varphi(x, \lambda) d\rho(\lambda) \right\} g(x) d_{\omega, q}x \\ &= \int_{-\tau}^{\tau} F(\lambda) \left\{ \int_{\omega_0}^{q^{-\mu}} \varphi(x, \lambda)g(x) d_{\omega, q}x \right\} d\rho(\lambda) \\ &= \int_{-\tau}^{\tau} F(\lambda)G(\lambda) d\rho(\lambda). \end{aligned} \quad (18)$$

By (17), we have

$$\int_{\omega_0}^{\infty} f(x)g(x) d_{\omega, q}x = \int_{-\infty}^{\infty} F(\lambda)G(\lambda) d\rho(\lambda). \quad (19)$$

By (18) and (19), we obtain

$$\int_{\omega_0}^{\infty} (f(x) - f_{\tau}(x))g(x) d_{\omega, q}x = \int_{|\lambda| > \tau} F(\lambda)G(\lambda) d\rho(\lambda).$$

By using Cauchy-Schwarz inequality, we see that

$$\begin{aligned} \left| \int_{\omega_0}^{\infty} (f(x) - f_{\tau}(x))g(x) d_{\omega, q}x \right|^2 &\leq \int_{|\lambda| > \tau} F^2(\lambda) d\rho(\lambda) \int_{|\lambda| > \tau} G^2(\lambda) d\rho(\lambda) \\ &\leq \int_{|\lambda| > \tau} F^2(\lambda) d\rho(\lambda) \int_{-\infty}^{\infty} G^2(\lambda) d\rho(\lambda). \end{aligned}$$

Applying this inequality to the function

$$g(x) = \begin{cases} f_{\tau}(x) - f(x), & x \in [\omega_0, q^{-\mu}] \\ 0, & x \in (q^{-\mu}, \infty), \end{cases}$$

we deduce that

$$\int_{\omega_0}^{q^{-\mu}} (f(x) - f_{\tau}(x))^2 d_{\omega, q}x \leq \int_{|\lambda| > \tau} F^2(\lambda) d\rho(\lambda). \quad (20)$$

Let us mention that the right-hand side of the inequality (20) does not depend on μ . Hence passing to the limit as $\tau \rightarrow \infty$ gives the desired result. Thus Theorem 5 is proved. \square

References

- [1] Aldwoah K.A.: Generalized time scales and associated difference equations. Ph.D. Thesis (2009)
- [2] Allahverdiev B.P., Tuna H.: An expansion theorem for q -Sturm-Liouville operators on the whole line. *Turkish J. Math.* 42 (3) (2018) 1060–1071.
- [3] Allahverdiev B.P., Tuna H.: Spectral expansion for singular Dirac system with impulsive conditions. *Turkish J. Math.* 42 (5) (2018) 2527–2545.
- [4] Allahverdiev B.P., Tuna H.: Eigenfunction expansion in the singular case for Dirac systems on time scales. *Konuralp J. Math.* 7 (1) (2019) 128–135.
- [5] Allahverdiev B.P., Tuna H.: The spectral expansion for Hahn-Dirac system on the whole line. *Turkish J. Math.* 43 (2019) 1668–1687.
- [6] Allahverdiev B.P., Tuna H.: Eigenfunction expansion for singular Sturm-Liouville problems with transmission conditions. *Electron. J. Differ. Equat.* 2019 (3) (2019) 1–10.
- [7] Allahverdiev B.P., Tuna H.: The Parseval equality and expansion formula for singular Hahn-Dirac system. In: Alparslan-Gok Z.S.: *Emerging Applications of Differential Equations and Game Theory*. IGI Global (2020) 209–235.
- [8] Álvarez-Nodarse R.: On characterizations of classical polynomials. *J. Comput. Appl. Math.* 196 (1) (2006) 320–337.
- [9] Annaby M.H., Hamza A.E., Aldwoah K.A.: Hahn difference operator and associated Jackson-Nörlund integrals. *J. Optim. Theory Appl.* 154 (2012) 133–153.
- [10] Annaby M.H., Hamza A.E., Makharesh S.D.: A Sturm-Liouville theory for Hahn difference operator. In: Xin Li, Zuhair Nashed: *Frontiers of Orthogonal Polynomials and q -Series*. World Scientific, Singapore (2018) 35–84.
- [11] Annaby M.A., Hassan H.A.: Sampling theorems for Jackson-Nörlund transforms associated with Hahn-difference operators. *J. Math. Anal. Appl.* 464 (1) (2018) 493–506.

- [12] J. Arvesú: On some properties of q -Hahn multiple orthogonal polynomials. *J. Comput. Appl. Math.* 233 (6) (2010) 1462–1469. doi:10.1016/j.cam.2009.02.062
- [13] J.M. Berezanskii: *Expansions in Eigenfunctions of Selfadjoint Operators*. Amer. Math. Soc., Providence (1968).
- [14] A. Dobrogowska, A. Odziejewicz: Second order q -difference equations solvable by factorization method. *J. Comput. Appl. Math.* 193 (1) (2006) 319–346.
- [15] G.Sh. Guseinov: Eigenfunction expansions for a Sturm-Liouville problem on time scales. *Int. J. Difference Equat.* 2 (1) (2007) 93–104.
- [16] G.Sh. Guseinov: An expansion theorem for a Sturm-Liouville operator on semi-unbounded time scales. *Adv. Dyn. Syst. Appl.* 3 (1) (2008) 147–160.
- [17] W. Hahn: Über orthogonalpolynome, die q -Differenzgleichungen genügen. *Math. Nachr.* 2 (1949) 4–34.
- [18] W. Hahn: Ein Beitrag zur Theorie der Orthogonalpolynome. *Monatsh. Math.* 95 (1983) 19–24.
- [19] A.E. Hamza, S.A. Ahmed: Existence and uniqueness of solutions of Hahn difference equations. *Adv. Difference Equat.* 316 (2013) 1–15.
- [20] A.E. Hamza, S.D. Makhareh: Leibniz' rule and Fubini's theorem associated with Hahn difference operator. *J. Adv. Math.* 12 (6) (2016) 6335–6345.
- [21] A. Huseynov, E. Bairamov: On expansions in eigenfunctions for second order dynamic equations on time scales. *Nonlinear Dyn. Syst. Theory* 9 (1) (2009) 77–88.
- [22] A. Huseynov: Eigenfunction expansion associated with the one-dimensional Schrödinger equation on semi-infinite time scale intervals. *Rep. Math. Phys.* 66 (2) (2010) 207–235.
- [23] F.H. Jackson: q -Difference equations. *Amer. J. Math.* 32 (1910) 305–314.
- [24] D.L. Jagerman: *Difference Equations with Applications to Queues*. Dekker, New York (2000).
- [25] C. Jordan: *Calculus of Finite Differences, 3rd edn.* Chelsea, New York (1965).
- [26] A.N. Kolmogorov, S.V. Fomin: *Introductory Real Analysis. Translated by R.A. Silverman.* Dover Publications, New York (1970).
- [27] K.H. Kwon, D.W. Lee, S.B. Park, B.H. Yoo: Hahn class orthogonal polynomials. *Kyungpook Math. J.* 38 (1998) 259–281.
- [28] P.A. Lesky: *Eine Charakterisierung der klassischen kontinuierlichen, diskreten und q -Orthogonalpolynome.* Shaker, Aachen (2005).
- [29] N. Levinson: A simplified proof of the expansion theorem for singular second order linear differential equations. *Duke Math. J.* 18 (1951) 57–71.
- [30] B.M. Levitan, I.S. Sargsjan: *Sturm-Liouville and Dirac Operators.* Springer (1991).
- [31] M.A. Naimark: *Linear Differential Operators, 2nd edn., 1968.* Nauka, Moscow (1969). English translation of 1st edn.
- [32] J. Petronilho: Generic formulas for the values at the singular points of some special monic classical $H_{q,\omega}$ -orthogonal polynomials. *J. Comput. Appl. Math.* 205 (2007) 314–324.
- [33] T. Sitthiwiratham: On a nonlocal boundary value problem for nonlinear second-order Hahn difference equation with two different q, ω -derivatives. *Adv. Difference Equat.* 2016 (1) (2016). Article number 116

- [34] M.H. Stone: A comparison of the series of Fourier and Birkhoff. *Trans. Amer. Math. Soc.* 28 (1926) 695–761.
- [35] M.H. Stone: *Linear Transformations in Hilbert Space and Their Application to Analysis*. Amer. Math. Soc. (1932).
- [36] E.C. Titchmarsh: *Eigenfunction Expansions Associated with Second-Order Differential Equations. Part I. Second Edition*. Clarendon Press, Oxford (1962).
- [37] H. Weyl: Über gewöhnliche Differentialgleichungen mit Singuliritäten und die zugehörigen Entwicklungen willkürlicher Funktionen. *Math. Annal.* 68 (1910) 220–269.
- [38] K. Yosida: On Titchmarsh-Kodaira formula concerning Weyl-Stone eigenfunction expansion. *Nagoya Math. J.* 1 (1950) 49–58.
- [39] K. Yosida: *Lectures on Differential and Integral Equations*. Springer, New York (1960).

Received: 22 October 2018

Accepted for publication: 15 February 2019

Communicated by: Pasha Zusmanovich