

Alan S. Dow; Saharon Shelah

On the bounding, splitting, and distributivity numbers

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 64 (2023), No. 3, 331–351

Persistent URL: <http://dml.cz/dmlcz/152302>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2023

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

# On the bounding, splitting, and distributivity numbers

ALAN DOW, SAHARON SHELAH

*Abstract.* The cardinal invariants  $\mathfrak{h}, \mathfrak{b}, \mathfrak{s}$  of  $\mathcal{P}(\omega)$  are known to satisfy that  $\omega_1 \leq \mathfrak{h} \leq \min\{\mathfrak{b}, \mathfrak{s}\}$ . We prove that all inequalities can be strict. We also introduce a new upper bound for  $\mathfrak{h}$  and show that it can be less than  $\mathfrak{s}$ . The key method is to utilize finite support matrix iterations of ccc posets following paper Ultrafilters with small generating sets by A. Blass and S. Shelah (1989).

*Keywords:* cardinal invariants of the continuum; matrix forcing

*Classification:* 03E15

## 1. Introduction

Of course the cardinal invariants of the continuum discussed in this article are very well known, see [15, page 111], so we just give a brief reminder. They deal with the mod finite ordering of the infinite subsets of the integers. We follow convention and let  $[\omega]^\omega$  (or  $[\omega]^{\aleph_0}$ ) denote the family of infinite subsets of  $\omega$ . A set  $A$  is a pseudo-intersection of a family  $\mathcal{Y} \subset [\omega]^\omega$  if  $A$  is infinite and  $A \setminus Y$  is finite for all  $Y \in \mathcal{Y}$ . The family  $\mathcal{Y}$  has the strong finite intersection property (sfip) if every finite subset has infinite intersection and  $\mathfrak{p}$  is the minimum cardinal for which there is such a family with no pseudointersection. A family  $\mathcal{I} \subset \mathcal{P}(\omega)$  is an ideal if it is closed under finite unions and mod finite subsets. An ideal  $\mathcal{I} \subset \mathcal{P}(\omega)$  is dense if every  $Y \in [\omega]^\omega$  contains an infinite member of  $\mathcal{I}$ . A set  $S \subset \omega$  is *unsplit* by a family  $\mathcal{Y} \subset [\omega]^\omega$  if  $S$  is mod finite contained in one member of  $\{Y, \omega \setminus Y\}$  for each  $Y \in \mathcal{Y}$ . The splitting number  $\mathfrak{s}$  is the minimum cardinal of a family  $\mathcal{Y}$  for which there is no infinite set unsplit by  $\mathcal{Y}$  (i.e. every  $S \in [\omega]^\omega$  is *split* by some member of  $\mathcal{Y}$  and  $\mathcal{Y}$  is called a splitting family). The bounding number  $\mathfrak{b}$  can easily be defined in these same terms, but it is best defined by the mod finite ordering “ $<^*$ ” on the family of functions  $\omega^\omega$ . The cardinal  $\mathfrak{b}$  is the minimum cardinal for which there is a  $<^*$ -unbounded family  $B \subset \omega^\omega$  with  $|B| = \mathfrak{b}$ .

The finite support iteration of the standard Hechler poset was shown in [2] to produce models of  $\aleph_1 = \mathfrak{s} < \mathfrak{b}$ . The consistency of  $\aleph_1 = \mathfrak{b} < \mathfrak{s} = \aleph_2$  was established in [17] with a countable support iteration of a special poset we now call  $\mathcal{Q}_{\text{Bould}}$ . It is shown in [11] that one can use Cohen forcing to select countable chain condition (ccc) subposets of  $\mathcal{Q}_{\text{Bould}}$  and finite support iterations to obtain models of  $\aleph_1 < \mathfrak{b} < \mathfrak{s} = \mathfrak{b}^+$ . This result was improved in [5] to show that the gap between  $\mathfrak{b}$  and  $\mathfrak{s}$  can be made arbitrarily large. The papers [4], [5] and [6] are able to use ccc versions of the well-known Mathias forcing in their iterations in place of those discovered in [11]. The paper [5] also nicely expands on the method of matrix iterated forcing first introduced in [4], as do a number of more recent papers, see [9], [16] and [10] using template forcing. The distributivity number (degree)  $\mathfrak{h}$  was first studied in [1]. It equals the minimum number of dense ideals whose intersection is simply the Fréchet ideal  $[\omega]^{<\omega}$ . It was shown in [1], that  $\mathfrak{p} \leq \mathfrak{h} \leq \min\{\mathfrak{b}, \mathfrak{s}\}$ . Our goal is to separate all these cardinals. We succeed but confront a new problem since we use the result, also from [1], that  $\mathfrak{h} \leq \text{cof}(\mathfrak{c})$ .

## 2. A new bound on $\mathfrak{h}$

In [1], a family  $\mathfrak{A}$  of maximal almost disjoint families of infinite subsets of  $\omega$  is called a matrix. A matrix  $\mathfrak{A}$  is *shattering* if the entire collection  $\bigcup \mathfrak{A}$  is splitting. Evidently, if  $\{s_\alpha : \alpha < \kappa\}$  is a splitting family, then the family  $\mathfrak{A} = \{\{s_\alpha, \omega \setminus s_\alpha\} : \alpha < \kappa\}$  is a shattering matrix. A shattering matrix  $\mathfrak{A} = \{\mathcal{A}_\alpha : \alpha < \kappa\}$  is *refining*, if for all  $\alpha < \beta < \kappa$ ,  $\mathcal{A}_\beta$  refines  $\mathcal{A}_\alpha$  in the natural sense that each member of  $\mathcal{A}_\beta$  is mod finite contained in some member of  $\mathcal{A}_\alpha$ . Finally, a *base matrix* is a refining shattering matrix  $\mathfrak{A}$  satisfying that  $\bigcup \mathfrak{A}$  is dense in  $(\mathcal{P}(\omega)/\text{fin}, \subset^*)$  (i.e. a  $\pi$ -base for  $\omega^*$ ).

We add condition (6) to the following result from [1].

**Lemma 2.1.** *The value of  $\mathfrak{h}$  is the least cardinal  $\kappa$  such that any of the following holds:*

- (1) *the Boolean algebra  $\mathcal{P}(\omega)/\text{fin}$  is not  $\kappa$ -distributive;*
- (2) *there is a shattering matrix of cardinality  $\kappa$ ;*
- (3) *there is a shattering and refining matrix indexed by  $\kappa$ ;*
- (4) *there is a base matrix of cardinality  $\kappa$ ;*
- (5) *there is a family of  $\kappa$  many nowhere dense subsets of  $\omega^*$  whose union is dense;*
- (6) *there is a sequence  $\{\mathcal{S}_\alpha : \alpha < \kappa\}$  of splitting families satisfying that no 1-to-1 selection  $\langle s_\alpha : \alpha \in \kappa \rangle \in \Pi\{\mathcal{S}_\alpha : \alpha \in \kappa\}$  has a pseudo-intersection.*

PROOF: Since (1)–(5) are proven in [1], it is sufficient to prove that for a cardinal  $\kappa$  (3) and (6) are equivalent. First suppose that  $\mathfrak{A} = \{\mathcal{A}_\alpha : \alpha < \kappa\}$  is a refining and

shattering matrix. Since the matrix is refining, it follows easily that  $\{\mathcal{A}_\beta: \alpha \leq \beta < \kappa\}$  is a shattering matrix for each  $\alpha < \kappa$ . Therefore,  $\mathcal{S}_\alpha = \bigcup\{\mathcal{A}_\beta: \alpha \leq \beta\}$  is a splitting family for each  $\alpha < \kappa$ . Similarly, the refining property ensures that if  $\langle a_\alpha: \alpha \in \kappa \rangle \in \Pi\{\mathcal{S}_\alpha: \alpha \in \kappa\}$ , then  $\{a_\alpha: \alpha \in \kappa\}$  has no pseudo-intersection.

Now assume that  $\{\mathcal{S}_\alpha: \alpha < \kappa\}$  is a sequence of splitting families as in (6). By [1], it is sufficient to prove that  $\mathfrak{h} \leq \kappa$ , so let us assume that  $\kappa < \mathfrak{h}$ . We now make an observation about  $\kappa$ : for each infinite  $b \subset \omega$ ,  $\alpha < \kappa$  and family  $\mathcal{S}' \subset [\omega]^\omega$  of cardinality less than  $\kappa$ , there is an infinite  $a \subset b$  and an  $s \in \mathcal{S}_\alpha \setminus \mathcal{S}'$  such that  $a \subset s$  and  $s$  splits  $b$ . We prove this claim. We may ignore all members of  $\mathcal{S}'$  that are mod finite disjoint, or mod finite include,  $b$ . Since the family  $\{\{s' \cap b, b \setminus s'\}: s' \in \mathcal{S}'\}$  is not shattering (as a family of subsets of  $b$ ) there is an infinite  $b' \subset b$  that is not split by  $\mathcal{S}'$ . Choose any  $s \in \mathcal{S}_\alpha$  that splits  $b'$  and let  $a = s \cap b'$ . Evidently,  $s$  also splits  $b$ . Since the ideal generated by a splitting family is dense, we may choose a maximal almost disjoint family  $\mathcal{A}_0$  contained in the ideal generated by  $\mathcal{S}_0$ . Let  $s_0$  denote any mapping from  $\mathcal{A}_0$  into  $\mathcal{S}_0$  satisfying that  $a \subset s_0(a)$  for all  $a \in \mathcal{A}_0$ . Suppose that  $\alpha < \kappa$  and that we have chosen a refining sequence  $\{\mathcal{A}_\gamma: \gamma < \alpha\}$  of maximal almost disjoint families together with mappings  $\{s_\gamma: \gamma < \alpha\}$  so that for each  $a \in \mathcal{A}_\gamma$ ,  $a \subset s_\gamma(a) \in \mathcal{S}_\gamma$ . The extra induction assumption is that for all  $a \in \mathcal{A}_\gamma$ ,  $s_\gamma(a)$  is not an element of  $\{s_\beta(a'): \beta < \gamma \text{ and } a \subset^* a' \in \mathcal{A}_\beta\}$ . The existence of the family  $\mathcal{A}_\alpha$  and the mapping  $s_\alpha$  satisfying the induction conditions easily follows from the above observation. Now we verify that  $\mathfrak{A} = \{\mathcal{A}_\alpha: \alpha < \kappa\}$  satisfies that  $\bigcup \mathfrak{A}$  is splitting. Fix any infinite  $b \subset \omega$  and choose  $a_\alpha \in \mathcal{A}_\alpha$  for each  $\alpha \in \kappa$  so that  $b \cap a_\alpha$  is infinite. By construction,  $\{s_\alpha(a_\alpha): \alpha \in \kappa\}$  is a 1-to-1 selection from  $\Pi\{\mathcal{S}_\alpha: \alpha \in \kappa\}$ . Since  $b$  is therefore not a pseudo-intersection, there is an  $\alpha < \kappa$  such that  $b \setminus s_\alpha(a_\alpha) \subset b \setminus a_\alpha$  is infinite.  $\square$

The following is an immediate corollary to condition (6) in Lemma 2.1 and provide two approaches to bounding the value of  $\mathfrak{h}$ .

**Corollary 2.2** ([1], [3]). (1) If  $\mathfrak{c}$  is singular, then  $\mathfrak{h} \leq \text{cf}(\mathfrak{c})$ .

(2) A poset  $\mathbb{P}$  forces that  $\mathfrak{h} \leq \kappa$  if  $\mathbb{P}$  preserves  $\kappa$  and can be written as an increasing chain  $\{\mathbb{P}_\alpha: \alpha < \kappa\}$  of completely embedded posets satisfying that each  $\mathbb{P}_{\alpha+1}$  adds a real not added by  $\mathbb{P}_\alpha$ .

PROOF: For the statement in (1), let  $\{\kappa_\alpha: \alpha < \text{cf}(\mathfrak{c})\}$  be increasing and cofinal in  $\mathfrak{c}$ . Let  $\{x_\xi: \xi \in \mathfrak{c}\}$  be an enumeration of  $[\omega]^{\aleph_0}$ . To apply (6) from Lemma 2.1, let  $\mathcal{S}_\alpha = \{x_\xi: (\forall \eta < \kappa_\alpha) x_\eta \not\subset^* x_\xi\}$ . For the statement in (2), let  $G$  be a  $\mathbb{P}$ -generic filter and for each  $\alpha \in \kappa$ , let  $G_\alpha = G \cap \mathbb{P}_\alpha$ . To apply (6), let  $\mathcal{S}_\alpha$  be the set of  $x \in [\omega]^{\aleph_0}$  that contain no infinite  $y \in V[G_\alpha]$ . To see that  $\mathcal{S}_\alpha$  is splitting

in either case, given any infinite  $x \subset \omega$ , consider an enumeration  $\{x_t: t \in 2^{<\omega}\}$ . Then, for all  $\alpha \in \kappa$ , there is an  $f_\alpha \in 2^\omega$  so that  $\{x_{f_\alpha \upharpoonright n}: n \in \omega\} \in \mathcal{S}_\alpha$ .  $\square$

Our introduction of condition (6) in Lemma 2.1 is motivated by the fact that it provides us with a new approach to bounding  $\mathfrak{h}$ . We introduce the following variant of condition (6) in Lemma 2.1 and note that a shattering refining matrix will fail to satisfy the second condition.

**Definition 2.3.** Let  $\kappa < \lambda$  be cardinals and say that a family  $\{x_\alpha: \alpha < \lambda\}$  of infinite subsets of  $\omega$  is  $(\kappa, \lambda)$ -shattering if for all infinite  $b \subset \omega$

- (1) the set  $\{\alpha < \lambda: b \subset^* x_\alpha\}$  has cardinality less than  $\kappa$ ; and
- (2) the set  $\{\alpha < \lambda: b \cap x_\alpha =^* \emptyset\}$  has cardinality less than  $\lambda$ .

Say that  $\{x_\alpha: \alpha < \lambda\}$  is strongly  $(\kappa, \lambda)$ -shattering if it contains no splitting family of size less than  $\lambda$ .

Needless to say a  $(\kappa, \lambda)$ -shattering family is strongly  $(\kappa, \lambda)$ -shattering if  $\lambda = \mathfrak{s}$  and this is the kind of families we are interested in. However it seems likely that producing strongly  $(\kappa, \lambda)$ -shattering families would be interesting (and as difficult) even without requiring that  $\lambda = \mathfrak{s}$ . Nevertheless  $\mathfrak{s}$  is necessarily less than  $\lambda$  as we show next.

**Proposition 2.4.** *If there is a  $(\kappa, \lambda)$ -shattering family, then  $\mathfrak{h} \leq \kappa$  and  $\mathfrak{s} \leq \lambda$ .*

PROOF: Let  $\mathcal{S} = \{x_\alpha: \alpha < \lambda\}$  be a  $(\kappa, \lambda)$ -shattering family. Given any infinite  $b \subset \omega$ , there is a  $\beta < \lambda$  such that each of  $b \subset^* x_\beta$  and  $b \cap x_\beta =^* \emptyset$  fail. This means that  $\mathcal{S}$  is splitting. By condition (1) in Definition 2.3 and applying condition (6) of Lemma 2.1 with  $\mathcal{S}_\alpha = \mathcal{S}$  for all  $\alpha < \kappa$ , it follows that  $\mathfrak{h} \leq \kappa$ .  $\square$

For any index set  $I$  the standard poset for adding Cohen reals,  $\mathcal{C}_I$ , is the set of all finite functions into 2 with domain a subset of  $I$  where  $p < q$  providing  $p \supset q$ . If  $I = \lambda$  is an ordinal, then we may use  $\dot{x}_\alpha$  to be the canonical  $\mathcal{C}_\lambda$ -name  $\{(\check{n}, \{\langle \alpha + n, 1 \rangle\}: n \in \omega)\}$  (i.e., for  $s \in \mathcal{C}_\lambda$ ,  $s \Vdash n \in \dot{x}_\alpha$  providing  $s(\alpha + n) = 1$ ).

It is routine to verify that, for any regular cardinal  $\lambda > \aleph_1$ , forcing with  $\mathcal{C}_\lambda$  will naturally add an  $(\aleph_1, \lambda)$ -shattering family but it is clear that this family would not be strongly  $(\aleph_1, \lambda)$ -shattering. Nevertheless, it may be possible with further forcing, to have it become strongly  $(\kappa, \lambda)$ -shattering for some  $\aleph_1 \leq \kappa < \mathfrak{s}$ .

In Theorem 5.9 we will prove that it is consistent with  $\aleph_2 < \kappa^+ < \mathfrak{c}$  that there is a strongly  $(\kappa, \kappa^+)$ -shattering family.

**Question 2.1.** Assume that  $\kappa < \lambda$  are regular cardinals and that there is a strongly  $(\kappa, \lambda)$ -shattering family. We pose the following questions.

- (1) Is it consistent that  $\kappa^+ < \lambda$ ?

- (2) Is it consistent that  $\lambda < \mathfrak{b}$ ?
- (3) Is it consistent that  $\kappa < \mathfrak{b} < \lambda$ ?

### 3. Matrix forcing and distinguishing $\mathfrak{h}, \mathfrak{s}, \mathfrak{b}$

In this section we recall the forcing methods for distinguishing  $\mathfrak{b}$  and  $\mathfrak{s}$  and apply them to prove the main results. We denote by  $\mathbb{D}$  the standard (Hechler) poset for adding a dominating real. The poset  $\mathbb{D}$  is an ordering on  $\omega^{<\omega} \times \omega^\omega$  where  $(s, f) < (t, g)$  providing  $g \leq f$  and  $s$  extends  $t$  by values that are coordinatewise above  $g$ . Given a sfip family  $\mathcal{F}$  of subsets of  $\omega$ , there are two main posets for adding a pseudo-intersection. The Mathias–Prikry style poset is  $\mathbb{M}(\mathcal{F})$  and consists of pairs  $(a, A)$  where  $A$  is in the filter base generated by  $\mathcal{F}$ ,  $a \subset \min(A)$ , and  $\mathbb{M}(\mathcal{F})$  is ordered by  $(a_1, A_1) < (a_2, A_2)$  providing  $a_2 \subset a_1 \subset a_2 \cup A_2$  and  $A_1 \subset A_2$ . When the context is clear, we will let  $\dot{x}_{\mathcal{F}}$  denote the canonical name,  $\{(\check{n}, (a, \omega \setminus n + 1)) : n \in a \subset n + 1\}$ , which is forced to be the desired pseudo-intersection. When  $\mathcal{U}$  is a free ultrafilter on  $\omega$ ,  $\mathbb{M}(\mathcal{U})$  was the poset used in [4] and [5] and, in this case,  $\dot{x}_{\mathcal{U}}$  is unsplit by the set of ground model subsets of  $\omega$ . When mixed with matrix iteration methods, the ultrafilter  $\mathcal{U}$  can be constructed so as to not add a dominating real.

The Laver style poset,  $\mathbb{L}(\mathcal{F})$ , is also very useful in matrix iterations and is defined as follows. The members of  $\mathbb{L}(\mathcal{F})$  are subtrees  $T$  of  $\omega^{<\omega}$  with a root or stem,  $\text{root}(T)$ , and for all  $\text{root}(T) \subseteq t \in T$ , the set  $\text{Br}(T, t) = \{j \in \omega : t \frown j \in T\}$  is an element of the filter generated by  $\mathcal{F}$ . This poset is ordered by “ $\subset$ ”. For each  $T \in \mathbb{L}(\mathcal{F})$  and  $t \in T$ , the subtree  $T_t = \{t' \in T : t \cup t' \in \omega^{<\omega}\}$  is also a condition. The generic function,  $\dot{f}_{\mathbb{L}(\mathcal{F})}$ , added by  $\mathbb{L}(\mathcal{F})$  can be described by the name of the union of the branch of  $\omega^{<\omega}$  named by  $\{(\check{t}, (\omega^{<\omega})_t) : t \in \omega^{<\omega}\}$ . This poset forces that  $\dot{f}_{\mathbb{L}(\mathcal{F})}$  dominates the ground model reals and the range of  $\dot{f}_{\mathbb{L}(\mathcal{F})}$  is a pseudo-intersection of  $\mathcal{F}$ . Again, if  $\mathcal{F}$  is an ultrafilter, this pseudo-intersection is not split by any ground model set.

For each sfip family  $\mathcal{U}$  on  $\omega$ , each of the posets  $\mathbb{D}$ ,  $\mathbb{M}(\mathcal{U})$ , and  $\mathbb{L}(\mathcal{U})$  is  $\sigma$ -centered. We just need this for the fact that this ensures that they are upwards ccc.

For a poset  $P$  and a set  $X$ , a canonical  $P$ -name for a subset of  $X$  will be a name of the form  $\bigcup \{\check{x} \times A_x : x \in X\}$  where for each  $x \in X$ ,  $A_x$  is an antichain of  $P$ . Of course if  $\dot{Y}$  is any  $P$ -name of a subset of  $X$ , there is a canonical name that is forced to equal it. When we say that a poset  $P$  forces a statement, we intend the meaning that every element (i.e.  $1_P$ ) of  $P$  forces that statement. We write  $P \triangleleft Q$  to mean that  $P$  is a complete suborder of  $Q$ .

The terminology “matrix iterations” is used in [5], see also forthcoming preprint (F1222) from the second author.

**Definition 3.1.** For an infinite cardinal  $\kappa$  with uncountable cofinality, and an ordinal  $\zeta$ , a  $\kappa \times \zeta$ -matrix iteration is a family

$$\langle \langle \mathbb{P}_{\alpha, \xi} : \alpha \leq \kappa, \xi \leq \zeta \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa, \xi < \zeta \rangle \rangle$$

where for each  $\alpha < \beta \leq \kappa$  and  $\xi < \eta \leq \zeta$ :

- (1)  $\mathbb{P}_{\beta, \xi}$  is a ccc poset;
- (2)  $\mathbb{P}_{\alpha, \xi} < \cdot \mathbb{P}_{\beta, \xi} < \cdot \mathbb{P}_{\beta, \eta}$ ;
- (3)  $\mathbb{P}_{\kappa, \xi}$  is the union of the chain  $\{\mathbb{P}_{\gamma, \xi} : \gamma < \kappa\}$ ;
- (4)  $\dot{\mathbb{Q}}_{\alpha, \xi}$  is a  $\mathbb{P}_{\alpha, \xi}$ -name of a ccc poset and  $\mathbb{P}_{\alpha, \xi+1} = \mathbb{P}_{\alpha, \xi} * \dot{\mathbb{Q}}_{\alpha, \xi}$ ;
- (5) if  $\eta$  is a limit, then  $\mathbb{P}_{\beta, \eta} = \bigcup \{\mathbb{P}_{\beta, \gamma} : \gamma < \eta\}$ .

One constructs  $\kappa \times \zeta$ -matrices by recursion on  $\zeta$  and, for successor steps, by careful choice of the component sequence  $\{\dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa\}$ . An important observation is that all the work is in the successor steps. The following is from [5, Lemma 3.10]

**Lemma 3.2.** *If  $\zeta$  is a limit then a family*

$$\langle \langle \mathbb{P}_{\alpha, \xi} : \alpha \leq \kappa, \xi \leq \zeta \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa, \xi < \zeta \rangle \rangle$$

*is a  $\kappa \times \zeta$ -matrix iteration provided that for all  $\eta < \zeta$  and  $\beta \leq \kappa$ :*

- (1)  $\langle \langle \mathbb{P}_{\alpha, \xi} : \alpha \leq \kappa, \xi \leq \eta \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa, \xi < \eta \rangle \rangle$  is a  $\kappa \times \eta$ -matrix iteration;  
and
- (2)  $\mathbb{P}_{\beta, \zeta} = \bigcup \{\mathbb{P}_{\beta, \xi} : \xi < \zeta\}$ .

The following is well-known, see for example [16, Section 5] and [13].

**Proposition 3.3.** *For any  $\zeta$  and  $\kappa \times \zeta$ -matrix iteration*

$$\langle \langle \mathbb{P}_{\alpha, \xi} : \alpha \leq \kappa, \xi \leq \zeta \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa, \xi < \zeta \rangle \rangle$$

*the extension*

$$\langle \langle \mathbb{P}_{\alpha, \xi} : \alpha \leq \kappa, \xi \leq \zeta + 1 \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa, \xi < \zeta + 1 \rangle \rangle$$

*is a  $\kappa \times (\zeta + 1)$ -matrix iteration if either the following holds:*

- (1)<sub>Q</sub> for all  $\alpha \leq \kappa$ ,  $\dot{\mathbb{Q}}_{\alpha, \zeta}$  is the  $\mathbb{P}_{\alpha, \zeta}$ -name for  $\mathbb{D}$ ;
- (2)<sub>Q</sub> there is an  $\alpha < \kappa$  such that  $\dot{\mathbb{Q}}_{\beta, \zeta}$  is the trivial poset for  $\beta < \alpha$ ,  $\dot{\mathbb{Q}}_{\alpha, \zeta}$  is a  $\mathbb{P}_{\alpha, \zeta}$ -name of a  $\sigma$ -centered poset, and  $\dot{\mathbb{Q}}_{\beta, \zeta} = \dot{\mathbb{Q}}_{\alpha, \zeta}$  for all  $\alpha \leq \beta \leq \kappa$ .

Notice that if we define the extension as in (1)<sub>Q</sub> then we will be adding a dominating real, but even if  $\dot{\mathbb{Q}}_{\alpha, \zeta}$  is forced to equal  $\mathbb{D}$  in (2)<sub>Q</sub>, the real added will only dominate the reals added by  $\mathbb{P}_{\alpha, \zeta}$ .

**Proposition 3.4** ([4]). *Let  $M$  be a model of (a sufficient amount of) set-theory and  $P \in M$  be a poset that is also contained in  $M$ . Then for any  $f \in \omega^\omega$  that is*

not dominated by any  $g \in M \cap \omega^\omega$ ,  $P$  forces that  $f \not\leq \dot{g}$  for all  $P$ -names  $\dot{g} \in M$  of elements of  $\omega^\omega$ .

PROOF: Let  $p \in P$  and  $n \in \omega$ . It suffices to prove that there is a  $q < p$  in  $P$  and a  $k > n$  and  $m < f(k)$  such that  $q \Vdash \dot{g}(k) = m$ . Since  $p \in M$ , we can work in  $M$  and define a function  $h \in \omega^\omega$  by the rule that, for all  $k \in \omega$ , there is a  $q_k < p$  such that  $q_k \Vdash \dot{g}(k) = h(k)$ . Choose any  $k > n$  so that  $h(k) < f(k)$ . Then  $q_k \Vdash \dot{g}(k) < f(k)$  and proves that  $p \nVdash f \leq \dot{g}$ .  $\square$

An analogous result, with the same proof, holds for splitting.

**Proposition 3.5.** *Let  $M$  be a model of (a sufficient amount of) set-theory and  $P \in M$  be a poset that is also contained in  $M$ . If  $x \in [\omega]^\omega$  satisfies that  $y \not\subset x$  for all  $y \in M \cap [\omega]^\omega$ , then  $P$  forces that  $\dot{y} \not\subset x$  for all  $P$ -names  $\dot{y} \in M$  for elements of  $[\omega]^\omega$ .*

We also use the main construction from [4].

**Proposition 3.6.** *Suppose that*

$$\langle \langle \mathbb{P}_{\alpha,\xi}: \alpha \leq \kappa, \xi \leq \zeta \rangle, \langle \dot{\mathbb{Q}}_{\alpha,\xi}: \alpha \leq \kappa, \xi < \zeta \rangle \rangle$$

is a  $\kappa \times \zeta$ -matrix iteration and that  $\{\dot{f}_\alpha: \alpha < \kappa\}$  is a sequence satisfying that for all  $\alpha < \kappa$ :

- (1)  $\dot{f}_\alpha$  is a  $\mathbb{P}_{\alpha,\zeta}$ -name that is forced to be in  $\omega^\omega$ ;
- (2) for all  $\beta < \alpha$  and  $\mathbb{P}_{\beta,\zeta}$ -name  $\dot{g}$  of a member of  $\omega^\omega$ ,  $\mathbb{P}_{\alpha,\zeta}$  forces that  $\dot{f}_\alpha \not\leq \dot{g}$ .

Then there is a sequence  $\{\dot{\mathcal{U}}_{\alpha,\zeta}: \alpha \leq \kappa\}$  such that for all  $\alpha < \kappa$ :

- (3)  $\dot{\mathcal{U}}_{\alpha,\zeta}$  is a  $\mathbb{P}_{\alpha,\zeta}$ -name of an ultrafilter on  $\omega$ ;
- (4) for  $\beta < \alpha$ ,  $\dot{\mathcal{U}}_{\beta,\zeta}$  is a subset of  $\dot{\mathcal{U}}_{\alpha,\zeta}$ ;
- (5) for each  $\beta < \alpha$  and each  $\mathbb{P}_{\beta,\zeta} * \mathbb{M}(\dot{\mathcal{U}}_{\beta,\zeta})$ -name  $\dot{g}$  of an element of  $\omega^\omega$ ,  $\mathbb{P}_{\alpha,\zeta} * \mathbb{M}(\dot{\mathcal{U}}_{\alpha,\zeta})$  forces that  $\dot{f}_\alpha \not\leq \dot{g}$ ; and
- (6)  $\langle \langle \mathbb{P}_{\alpha,\xi}: \alpha \leq \kappa, \xi \leq \zeta + 1 \rangle, \langle \dot{\mathbb{Q}}_{\alpha,\xi}: \alpha \leq \kappa, \xi < \zeta + 1 \rangle \rangle$  is a  $\kappa \times (\zeta + 1)$ -matrix iteration, where for each  $\alpha \leq \kappa$ ,  $\mathbb{P}_{\alpha,\zeta+1} = \mathbb{P}_{\alpha,\zeta} * \dot{\mathbb{Q}}_{\alpha,\zeta}$  and  $\dot{\mathbb{Q}}_{\alpha,\zeta}$  is the  $\mathbb{P}_{\alpha,\zeta}$ -name for  $\mathbb{M}(\dot{\mathcal{U}}_{\alpha,\zeta})$ .

We record two more well-known preparatory preservation results.

**Proposition 3.7** ([2]). *Suppose that  $M \subset N$  are models of (a sufficient amount of) set-theory and that  $G$  is  $\mathbb{D}$ -generic over  $N$ . If  $x \in N \cap [\omega]^\omega$  does not include any  $y \in M \cap [\omega]^\omega$ , it will not include any  $y \in M[G] \cap [\omega]^\omega$ .*

**Proposition 3.8.** *Assume that  $\{P_\alpha: \alpha \leq \delta\}$  is a  $<\cdot$ -increasing chain of ccc posets with  $P_\delta = \bigcup \{P_\alpha: \alpha < \delta\}$ . Let  $G_\delta$  be  $P_\delta$ -generic. Let  $x \in [\omega]^\omega$  and  $f \in \omega^\omega$ . Then each of the following holds:*

- (1) If  $f \not\leq g$  for each  $g \in V[G_\alpha]$  and for all  $\alpha < \delta$ , then  $f \not\leq g$  for each  $g \in V[G_\delta]$ .
- (2) If  $x$  does not contain any  $y \in [\omega]^\omega \cap V[G_\alpha]$  for all  $\alpha < \kappa$ , then  $x$  does not contain any  $y \in [\omega]^\omega \cap V[G_\delta]$ .

PROOF: We prove only (1) since the proof of (2) is similar. If  $\delta$  has uncountable cofinality, then there is nothing to prove since  $V[G_\delta] \cap \omega^\omega$  would then equal  $\bigcup \{V[G_\alpha] \cap \omega^\omega : \alpha < \delta\}$ . Otherwise, consider any  $P_\delta$ -name  $\dot{g}$  and condition  $p \in P_\delta$  forcing that  $\dot{g} \in \omega^\omega$ . We prove that  $p$  does not force that  $\dot{g}(n) > f(n)$  for all  $k < n$ . We may assume that  $\dot{g}$  is a canonical name, so let  $\dot{g} = \bigcup \{(\overline{n}, \overline{m}) \times A_{n,m} : n, m \in \omega \times \omega\}$ . Choose any  $\alpha < \delta$  so that  $p \in P_\alpha$  and work in  $V[G_\alpha]$ . We define a function  $h \in \omega^\omega \cap V[G_\alpha]$ . For each  $n \in \omega$ , we set  $h(n)$  to be the minimum  $m$  such that there is  $q_{n,m} \in A_{n,m}$  having a  $P_\alpha$ -reduct  $p_{n,m} \in G_\alpha$ . Since  $A_n = \bigcup \{A_{n,m} : m \in \omega\}$  is predense in  $P_\kappa$ , the set of  $P_\alpha$ -reducts of members of  $A_n$  is predense in  $P_\alpha$ . By hypothesis, there is a  $k < n$  such that  $h(n) < f(n)$ . Since  $q_{n,h(n)}$  is compatible with  $p$ , this prove that  $p \nVdash \dot{g}(n) > f(n)$ .  $\square$

#### 4. Building the models to distinguish $\mathfrak{h}, \mathfrak{b}, \mathfrak{s}$

For simplicity we assume GCH. Let  $\aleph_1 \leq \mu < \kappa < \lambda$  be regular cardinals and assume that  $\theta > \lambda$  is a cardinal with cofinality  $\mu$ . We will need to enumerate names in order to force that  $\mathfrak{p} \geq \mu$ . For each ccc poset  $\tilde{P} \in H(\theta^+)$  let  $\{\dot{Y}(\tilde{P}, \xi) : \xi < \theta\}$  be an enumeration of the set of all canonical  $\tilde{P}$ -names of subsets of  $\omega$ . Also let  $\{S_\xi : \xi < \theta\}$  be an enumeration of all subsets of  $\theta$  that have cardinality less than  $\mu$ . For each  $\eta < \lambda$ , let  $\zeta_\eta$  denote the ordinal product  $\theta \cdot \eta$ .

**Theorem 4.1.** *There is a ccc poset that forces  $\mathfrak{p} = \mathfrak{h} = \mu$ ,  $\mathfrak{b} = \kappa$ ,  $\mathfrak{s} = \lambda$  and  $\mathfrak{c} = \theta$ .*

PROOF: The poset will be obtained by constructing a  $\kappa \times \zeta$ -matrix iteration where  $\zeta$  is the ordinal product  $\theta \cdot \lambda$  (the lexicographic ordering on  $\lambda \times \theta$ ). We begin with the  $\kappa \times \kappa$ -matrix iteration

$$\langle \langle \mathbb{P}_{\alpha, \xi} : \alpha \leq \kappa, \xi \leq \kappa \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi} : \alpha \leq \kappa, \xi < \kappa \rangle \rangle$$

where, for each  $\alpha < \kappa$ ,  $\mathbb{P}_{\alpha, \alpha}$  forces that  $\dot{\mathbb{Q}}_{\alpha, \alpha}$  is  $\mathbb{D}$ , for  $\beta < \alpha$ ,  $\dot{\mathbb{Q}}_{\beta, \alpha}$  is the trivial poset, and for  $\alpha \leq \beta \leq \kappa$ ,  $\dot{\mathbb{Q}}_{\beta, \alpha}$  equals  $\dot{\mathbb{Q}}_{\alpha, \alpha}$ . By Proposition 3.3, there is such a matrix. For each  $\alpha < \kappa$ , let  $\dot{f}_\alpha$  be the canonical name for the dominating real added by  $\mathbb{P}_{\alpha, \alpha+1}$ . By Propositions 3.4 and 3.8, it follows that for all  $\beta < \alpha < \kappa$ ,  $\mathbb{P}_{\alpha, \kappa}$  forces that  $\dot{f}_\alpha \not\leq \dot{g}$  for all  $\mathbb{P}_{\beta, \kappa}$ -names  $\dot{g}$  of elements of  $\omega^\omega$ .

We omit the routine enumeration details involved in the recursive construction and state the properties we require of our  $\kappa \times \zeta$ -matrix iteration. Each step of the

construction uses either (2) of Proposition 3.3 or Proposition 3.6 to choose the next sequence  $\{\dot{Q}_{\alpha,\xi}: \alpha \leq \kappa\}$ . In the case of Proposition 3.3 (2), the preservation of inductive condition (1) follows from Proposition 3.4. The preservation through limit steps follows from Proposition 3.8.

There is a matrix-iteration sequence

$$\langle \langle \mathbb{P}_{\alpha,\xi}: \alpha \leq \kappa, \xi \leq \zeta \rangle, \langle \dot{Q}_{\alpha,\xi}: \alpha \leq \kappa, \xi < \zeta \rangle \rangle$$

satisfying each of the following for each  $\xi < \zeta$ :

- (1) for each  $\beta < \alpha < \kappa$  and each  $\mathbb{P}_{\beta,\xi}$ -name  $\dot{g}$  for an element of  $\omega^\omega$ ,  $\mathbb{P}_{\alpha,\xi}$  forces that  $\dot{f}_\alpha \not\leq \dot{g}$ ;
- (2) for each  $\beta < \lambda$  with  $\zeta_{\beta+1} \leq \xi$  and each  $\eta < \theta$ , if  $\mathbb{P}_{\kappa,\zeta_\beta}$  forces that the family  $\mathcal{F}_{\beta,\eta} = \{\dot{Y}(\mathbb{P}_{\kappa,\zeta_\beta}, \gamma): \gamma \in S_\eta\}$  has the sfip, then there is a  $\bar{\eta} < \zeta_{\beta+1}$  and an  $\alpha < \kappa$  such that  $\dot{Q}_{\beta,\bar{\eta}}$  equals the  $\mathbb{P}_{\alpha,\bar{\eta}}$ -name for  $\mathbb{M}(\mathcal{F}_{\beta,\eta})$  for all  $\alpha \leq \beta \leq \kappa$ ;
- (3) for each  $\beta < \lambda$  such that  $\zeta_\beta < \xi$ ,  $\mathbb{P}_{\kappa,\zeta_{\beta+1}}$  equals  $\mathbb{P}_{\kappa,\zeta_\beta} * \mathbb{M}(\dot{\mathcal{U}}_{\kappa,\zeta_\beta})$  and  $\dot{\mathcal{U}}_{\kappa,\zeta_\beta}$  is a  $\mathbb{P}_{\kappa,\zeta_\beta}$ -name of an ultrafilter on  $\omega$ ;
- (4) for each  $\eta < \lambda$  and each  $\alpha < \kappa$  such that  $\zeta_\eta < \xi$ , then  $\dot{Q}_{\alpha,\zeta_\eta+\alpha}$  is the  $\mathbb{P}_{\alpha,\zeta_\eta+\alpha}$ -name for  $\mathbb{D}$ , and  $\dot{Q}_{\beta,\zeta_\eta+\alpha} = \dot{Q}_{\alpha,\zeta_\eta+\alpha}$  for all  $\alpha \leq \beta \leq \kappa$ .

Now we verify that  $P = \mathbb{P}_{\kappa,\zeta}$  has the desired properties. Since  $P$  is ccc, it preserves cardinals and clearly forces that  $\mathfrak{c} = \theta$ . It thus follows from Corollary 2.2 that  $\mathfrak{p} \leq \mathfrak{h} \leq \mu = \text{cf}(\mathfrak{c})$ . If  $\mathcal{Y}$  is a family of fewer than  $\mu$  many canonical  $P$ -names of subsets of  $\omega$ , then there is an  $\alpha < \kappa$  and  $\eta < \lambda$  such that  $\mathcal{Y}$  is a family of  $\mathbb{P}_{\alpha,\zeta_\eta}$ -names. It follows that there is a  $\beta < \theta$  such that  $\mathcal{Y}$  is equal to the set  $\{\dot{Y}(\mathbb{P}_{\kappa,\zeta_\beta}, \gamma): \gamma \in S_\eta\}$ . If  $\mathbb{P}_{\kappa,\zeta_\beta}$  forces that  $\mathcal{Y}$  has the sfip, then inductive condition 2 ensures that there is a  $P$ -name for a pseudo-intersection for  $\mathcal{Y}$ . This shows that  $P$  forces that  $\mathfrak{p} \geq \mu$ . It is clear that inductive condition 1 ensures that  $\mathfrak{b} \leq \kappa$ . We check that condition 4 ensure that  $\mathfrak{b} \geq \kappa$ . Suppose that  $\mathcal{G}$  is a family of fewer than  $\kappa$  many canonical  $P$ -names of members of  $\omega^\omega$ . We again find  $\eta < \lambda$  and  $\alpha < \kappa$  such that  $\mathcal{G}$  is a family of  $\mathbb{P}_{\alpha,\zeta_\eta}$ -names. Condition 4 forces there is a function that dominates  $\mathcal{G}$ . Finally we verify that condition 3 ensures that  $P$  forces that  $\mathfrak{s} = \lambda$ . If  $\mathcal{S}$  is any family of fewer than  $\lambda$ -many canonical  $P$ -names of subsets of  $\omega$ , then there is an  $\eta < \lambda$  such that  $\mathcal{S}$  is a family of  $\mathbb{P}_{\kappa,\zeta_\eta}$ -names. Evidently,  $\mathbb{P}_{\kappa,\zeta_{\eta+1}}$  adds a subset of  $\omega$  that is not split by  $\mathcal{S}$ . There are a number of ways to observe that for each  $\eta < \lambda$ ,  $\mathbb{P}_{\kappa,\zeta_{\eta+1}}$  adds a real that is Cohen over the extension by  $\mathbb{P}_{\kappa,\zeta_\eta}$ . This ensures that  $P$  forces that  $\mathfrak{s} \leq \lambda$ .  $\square$

In the next result we proceed similarly except that we first add  $\kappa$  many Cohen reals and preserve that they are splitting. We then cofinally add dominating reals with Hechler's  $\mathbb{D}$  and again use small posets to ensure  $\mathfrak{p} \geq \mu$ .

**Theorem 4.2.** *There is a ccc poset that forces  $\mathfrak{p} = \mathfrak{h} = \mu$ ,  $\mathfrak{s} = \kappa$ ,  $\mathfrak{b} = \lambda$  and  $\mathfrak{c} = \theta$ .*

PROOF: We begin with the  $\kappa \times \kappa$ -matrix iteration

$$\langle \langle \mathbb{P}_{\alpha, \xi}: \alpha \leq \kappa, \xi \leq \kappa \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi}: \alpha \leq \kappa, \xi < \kappa \rangle \rangle$$

where  $\mathbb{P}_{\alpha, \alpha}$  forces that  $\dot{\mathbb{Q}}_{\alpha, \alpha}$  is  $\mathcal{C}_\omega$ , for  $\beta < \alpha$ ,  $\dot{\mathbb{Q}}_{\beta, \alpha}$  is the trivial poset, and for  $\alpha \leq \beta \leq \kappa$ ,  $\dot{\mathbb{Q}}_{\beta, \alpha}$  equals  $\dot{\mathbb{Q}}_{\alpha, \alpha}$ . We let  $\dot{x}_\alpha$  denote the canonical Cohen real added by  $\mathbb{P}_{\alpha, \alpha+1}$ . Of course  $\mathbb{P}_{\alpha, \alpha+1}$  forces that neither  $\dot{x}_\alpha$  nor its complement include any infinite subsets of  $\omega$  that have, for any  $\beta < \alpha$ , a  $\mathbb{P}_{\beta, \alpha+1}$ -name. By Proposition 3.8, the inductive condition 4 below holds for  $\xi = \kappa$ .

Then, proceeding as in the proof of Theorem 4.1, we just assert the existence of a  $\kappa \times \zeta$ -matrix iteration

$$\langle \langle \mathbb{P}_{\alpha, \xi}: \alpha \leq \kappa, \xi \leq \zeta \rangle, \langle \dot{\mathbb{Q}}_{\alpha, \xi}: \alpha \leq \kappa, \xi < \zeta \rangle \rangle$$

satisfying each of the following for each  $\kappa \leq \xi < \zeta$ :

- (1) for each  $\beta < \alpha < \kappa$ ,  $\mathbb{P}_{\alpha, \xi}$  forces that neither  $\dot{x}_\alpha$  nor  $\omega \setminus \dot{x}_\alpha$  include any infinite subset of  $\omega$  that has a  $\mathbb{P}_{\beta, \xi}$ -name;
- (2) for each  $\eta < \lambda$  with  $\zeta_{\eta+1} \leq \xi$  and each  $\delta < \theta$ , if  $\mathbb{P}_{\kappa, \zeta_\eta}$  forces that the family  $\mathcal{F}_{\eta, \delta} = \{\dot{Y}(\mathbb{P}_{\kappa, \zeta_\eta}, \gamma): \gamma \in S_\delta\}$  has the sfip, then there is a  $\bar{\delta} < \zeta_{\eta+1}$  and an  $\alpha < \kappa$  such that  $\dot{\mathbb{Q}}_{\beta, \bar{\delta}}$  equals the  $\mathbb{P}_{\alpha, \bar{\delta}}$ -name for  $\mathbb{M}(\mathcal{F}_{\eta, \delta})$  for all  $\alpha \leq \beta \leq \kappa$ ;
- (3) for each  $\eta < \lambda$  and each  $\alpha < \kappa$  such that  $\zeta_\eta < \xi$ , then  $\dot{\mathbb{Q}}_{\alpha, \zeta_\eta + \alpha}$  is the  $\mathbb{P}_{\alpha, \zeta_\eta + \alpha}$ -name for  $\mathbb{M}(\dot{\mathcal{U}}_{\alpha, \zeta_\beta})$  where  $\dot{\mathcal{U}}_{\alpha, \zeta_\beta}$  is a  $\mathbb{P}_{\alpha, \zeta_\beta}$ -name of an ultrafilter on  $\omega$ , and  $\dot{\mathbb{Q}}_{\beta, \zeta_\eta + \alpha} = \dot{\mathbb{Q}}_{\alpha, \zeta_\eta + \alpha}$  for all  $\alpha \leq \beta \leq \kappa$ ;
- (4) for each  $\eta < \lambda$  such that  $\zeta_\eta < \xi$ ,  $\mathbb{P}_{\kappa, \zeta_\eta + 1}$  equals  $\mathbb{P}_{\kappa, \zeta_\eta} * \mathbb{D}$ .

Evidently conditions (2) and (3) are similar and can be achieved while preserving condition (1) by Proposition 3.3 (2). The fact that  $\mathbb{P}_{\kappa, \zeta_\eta} * \mathbb{D}$  preserves condition (1) follows from Proposition 3.7. Condition (1) ensures that  $\mathfrak{s} \leq \kappa$ , and by arguments similar to those in Theorem 4.1, condition (3) ensures that  $\mathfrak{s} \geq \kappa$ . The fact that  $\mathfrak{b} = \lambda$  (in fact  $\mathfrak{d} = \lambda$ ) follows easily from condition (4). The facts that  $\mathfrak{c} = \theta$ ,  $\mathfrak{p} \geq \mu$  and  $\mathfrak{h} = \mu$  are proven exactly as in Theorem 4.1.  $\square$

## 5. On $(\kappa, \lambda)$ -shattering

In this section we prove, see Theorem 5.9, that it is consistent that strongly  $(\kappa, \kappa^+)$ -shattering families exist. The method used in this section is the following generalization of matrix iterations used in [8]. A chain  $\{P_\alpha: \alpha < \delta\}$  is continuous if for every limit  $\alpha < \delta$ ,  $P_\alpha = \bigcup \{P_\beta: \beta < \alpha\}$ .

**Definition 5.1.** Let  $\kappa > \omega_1$  be a regular cardinal. For an ordinal  $\zeta$ , a  $\kappa \times \zeta$ -matrix of posets is a family  $\{P_{\alpha,\xi} : \alpha \leq \kappa, \xi < \zeta\}$  of ccc posets satisfying for each  $\alpha < \kappa$ , and  $\xi < \eta < \zeta$ :

- (1)  $P_{\alpha,\xi} < \cdot P_{\beta,\xi}$  for all  $\alpha < \beta \leq \kappa$ ;
- (2)  $P_{\beta,\xi} = \bigcup \{P_{\eta,\xi} : \eta < \beta\}$  for  $\beta \leq \kappa$  with  $\text{cf}(\beta) > \omega$ ; and
- (3) for some  $\gamma < \kappa$ ,  $P_{\beta,\xi} < \cdot P_{\beta,\eta}$  for all  $\gamma \leq \beta \leq \kappa$ ;
- (4) if  $\eta$  is a limit ordinal, there is a cub  $C \subset \eta$  and a  $\gamma < \kappa$  such that, for all  $\gamma \leq \beta < \kappa$ ,  $\{P_{\beta,\delta} : \delta \in C \cup \{\eta\}\}$  is a continuous  $< \cdot$ -increasing chain.

One must be careful with a  $\kappa \times \zeta$ -matrix since there is no natural extension or definition of  $P_{\alpha,\zeta}$  for  $\alpha < \kappa$ . However, when  $\text{cf}(\zeta) > \omega_1$  the matrix can be viewed as a matrix type construction of a ccc poset  $P_{\kappa,\zeta}$ .

**Lemma 5.2.** If  $\{P_{\alpha,\xi} : \alpha \leq \kappa, \xi < \zeta\}$  is a  $\kappa \times \zeta$ -matrix of posets with  $\kappa > \omega_1$  regular and  $\text{cf}(\zeta) > \omega_1$ , then the poset  $P_{\kappa,\zeta} = \bigcup \{P_{\kappa,\xi} : \xi < \zeta\}$  is ccc and satisfies that  $P_{\alpha,\xi} < \cdot P_{\kappa,\zeta}$  for all  $\alpha \leq \kappa$  and  $\xi < \zeta$ .

PROOF: Let  $\alpha < \kappa$  and  $\xi < \zeta$ . It follows from property (1) in Definition 5.1 that  $P_{\alpha,\xi} < \cdot P_{\kappa,\xi}$ . By (3) of Definition 5.1, we have that  $\{P_{\kappa,\eta} : \xi \leq \eta < \zeta\}$  is a  $< \cdot$ -chain. This implies that  $P_{\kappa,\xi} < \cdot P_{\kappa,\zeta}$ . Now we check that  $P_{\kappa,\zeta}$  is ccc. Assume that  $A \subset P_{\kappa,\zeta}$  has cardinality  $\aleph_1$ . Choose any  $\gamma_0 < \kappa$  so that  $A \subset \bigcup \{P_{\beta,\xi} : \beta < \gamma_0, \xi < \zeta\}$ . Similarly choose  $\eta < \zeta$  minimal so that  $A \subset \bigcup \{P_{\beta,\xi} : \beta < \gamma_0, \xi < \eta\}$ . By property (2) of Definition 5.1, there is a  $\gamma_0 \leq \gamma_1 < \kappa$  such that  $A \subset \bigcup \{P_{\gamma_1,\xi} : \xi < \eta\}$ . Now choose a cub  $C \subset \eta$  as in condition (4) of Definition 5.1, and, using conditions (2) and (3) of Definition 5.1, we can choose  $\zeta_1 \leq \zeta_2 < \kappa$  so that  $A \subset \bigcup \{P_{\zeta_2,\delta} : \delta \in C\} \subset P_{\zeta_2,\eta}$ . Since  $P_{\zeta_2,\eta}$  is ccc, it follows that  $A$  is not an antichain.  $\square$

We will use the method of matrix of posets from Definition 5.1 in which our main component posets to raise the value of  $\mathfrak{s}$  will be the Laver style posets. Before proceeding it may be helpful to summarize the rough idea of how we generalize the fundamental preservation technique of a matrix iteration. In a  $\kappa \times \kappa^+$ -matrix iteration, one may introduce a sequence  $\{\dot{a}_\alpha : \alpha < \kappa\}$  of  $P_{\kappa,1}$ -names that have no infinite pseudointersection. With this fixed enumeration, one then ensures that no  $P_{\alpha,\gamma}$ -name will be forced to be a subset of  $\dot{a}_\beta$  for any  $\alpha \leq \beta < \kappa$ . In the construction introduced in [8], we instead continually add to the list a  $P_{0,\gamma+1}$ -name  $\dot{a}_\gamma$  and at stage  $\mu < \kappa^+$ , we adopt a new enumeration of  $\{\dot{a}_\alpha : \alpha < \mu\}$  in order-type  $\kappa$  (coherent with previous listings) and again ensure that no  $P_{\alpha,\mu+1}$ -name is a subset of any  $\dot{a}_\beta$  for  $\beta$  not listed before  $\alpha$  in this new  $\mu$ th listing. We utilize a  $\square$ -principle to make these enumerations sufficiently coherent and to use as the required cub's in condition (4) of Definition 5.1. The greater flexibility in the definition of  $\kappa \times \kappa^+$ -matrix of posets makes this possible.

We recall some notions and results about these studied in [7], [8].

**Proposition 5.3.** *If  $P < P'$  are ccc posets, and  $\dot{D} \subset \dot{E}$  are, respectively, a  $P$ -name and a  $P'$ -name of ultrafilters on  $\omega$ , then  $P * \mathbb{L}(\dot{D}) < P' * \mathbb{L}(\dot{E})$ .*

**Definition 5.4.** A family  $\mathcal{A} \subset [\omega]^\omega$  is thin over a model  $M$  if for every  $I$  in the ideal generated by  $\mathcal{A}$  and every infinite family  $\mathcal{F} \in M$  consisting of pairwise disjoint finite sets of bounded size,  $I$  is disjoint from some member of  $\mathcal{F}$ .

It is routine to prove that for each limit ordinal  $\delta$ ,  $\mathcal{C}_\delta$  forces that the family  $\{\dot{x}_\alpha : \alpha \in \delta\}$ , as defined above, is thin over the ground model. In fact if  $\mathcal{A}$  is thin over some model  $M$ , then  $\mathcal{C}_\delta$  forces that  $\mathcal{A} \cup \{\dot{x}_\alpha : \alpha \in \delta\}$  is also thin over  $M$ . This is the notion we use to control that property (1) of the definition of a  $(\kappa, \kappa^+)$ -shattering sequence will be preserved while at the same time raising the value of  $\mathfrak{s}$ .

We first note that Proposition 3.5 extends to include this concept.

**Proposition 5.5.** *Suppose that  $M$  is a model of a sufficient amount of set-theory and that  $\mathcal{A} \subset [\omega]^\omega$  is thin over  $M$ . Then for any poset  $P$  such that  $P \in M$  and  $P \subset M$ ,  $\mathcal{A}$  is thin over the forcing extension by  $P$ .*

PROOF: Let  $\{\dot{F}_l : l \in \omega\}$  be  $P$ -names and suppose that  $p \in P$  forces that  $\{\dot{F}_l : l \in \omega\}$  are pairwise disjoint subsets of  $[\omega]^k$ ,  $k \in \omega$ . Also let  $I$  be any member of the ideal generated by  $\mathcal{A}$ . Working in  $M$ , recursively choose  $q_j < p$ ,  $j \in \omega$ , and  $H_j, l_j$  so that  $q_j \Vdash \dot{F}_{l_j} = \check{H}_j$  and  $H_j \cap \bigcup \{H_i : i < j\} = \emptyset$ . The sequence  $\{H_j : j \in \omega\}$  is a family in  $M$  of pairwise disjoint sets of cardinality  $k$ . Therefore there is a  $j$  with  $H_j \cap I = \emptyset$ . This proves that  $p$  does not force that  $I$  meets every member of  $\{\dot{F}_l : l \in \omega\}$ .  $\square$

**Lemma 5.6** ([8, 3.8]). *Let  $\kappa$  be a regular uncountable cardinal and let  $\{P_\beta : \beta \leq \kappa\}$  be a  $<\cdot$ -increasing chain of ccc posets with  $P_\kappa = \bigcup \{P_\alpha : \alpha < \kappa\}$ . Assume that, for each  $\beta < \kappa$ ,  $\dot{A}_\beta$  is a  $P_{\beta+1}$ -name of a subset of  $[\omega]^\omega$  that is forced to be thin over the forcing extension by  $P_\beta$ . Also let  $\dot{D}_0$  be a  $P_0 * \mathcal{C}_{\{0\} \times \mathfrak{c}}$ -name that is forced to be a Ramsey ultrafilter on  $\omega$ . Then there is a sequence  $\langle \dot{D}_\beta : 0 < \beta < \kappa \rangle$  such that for all  $\alpha < \beta < \kappa$ :*

- (1)  $\dot{D}_\beta$  is a  $P_\beta * \mathcal{C}_{(\beta+1) \times \mathfrak{c}}$ -name;
- (2)  $\dot{D}_\alpha$  is a subset of  $\dot{D}_\beta$ ;
- (3)  $P_\beta * \mathcal{C}_{(\beta+1) \times \mathfrak{c}}$  forces that  $\dot{D}_\beta$  is a Ramsey ultrafilter;
- (4)  $P_\alpha * \mathcal{C}_{(\alpha+1) \times \mathfrak{c}} * \mathbb{L}(\dot{D}_\alpha) < P_\beta * \mathcal{C}_{(\beta+1) \times \mathfrak{c}} * \mathbb{L}(\dot{D}_\beta)$ ; and
- (5)  $P_\beta * \mathcal{C}_{(\beta+1) \times \mathfrak{c}} * \mathbb{L}(\dot{D}_\beta)$  forces that  $\dot{A}_\beta$  is thin over the forcing extension by  $P_\alpha * \mathcal{C}_{(\alpha+1) \times \mathfrak{c}} * \mathbb{L}(\dot{D}_\alpha)$ .

**Lemma 5.7** ([8, 2.7]). *Assume that  $P_{0,0} < P_{1,0}$  and that  $\dot{A}$  is a  $P_{1,0}$ -name of a subset of  $[\omega]^\omega$ . Assume that  $\langle P_{0,\xi} : \xi < \delta \rangle$  and  $\langle P_{1,\xi} : \xi < \delta \rangle$  are  $<\cdot$ -chains such*

that  $P_{0,\xi} \dot{<} P_{1,\xi}$  for all  $\xi < \delta$ , and that  $P_{1,\xi}$  forces that  $\dot{\mathcal{A}}$  is thin over the forcing extension by  $P_{0,\xi}$  for all  $\xi < \delta$ . Then  $P_{1,\delta} = \bigcup \{P_{1,\xi} : \xi < \delta\}$  forces that  $\mathcal{A}$  is thin over the forcing extension by  $P_{0,\delta} = \bigcup \{P_{0,\xi} : \xi < \delta\}$ .

Before proving the next result we recall the notion of a  $\square_\kappa$ -sequence. For a set  $C$  of ordinals, let  $\sup(C)$  be the supremum,  $\bigcup C$ , of  $C$  and let  $C'$  denote the set of limit ordinals  $\alpha < \sup(C)$  such that  $C \cap \alpha$  is cofinal in  $\alpha$ . For a limit ordinal  $\alpha$ , a set  $C$  is a cub in  $\alpha$  if  $C \subset \alpha = \sup(C)$  and  $C' \subset C$ .

**Definition 5.8** ([14]). For a cardinal  $\kappa$ , the family  $\{C_\alpha : \alpha \in (\kappa^+)'\}$  is a  $\square_\kappa$ -sequence if for each  $\alpha \in (\kappa^+)'$ :

- (1)  $C_\alpha$  is a cub in  $\alpha$ ;
- (2) if  $\text{cf}(\alpha) < \kappa$ , then  $|C_\alpha| < \kappa$ ;
- (3) if  $\beta \in C'_\alpha$ , then  $C_\beta = C_\alpha \cap \beta$ .

If there is a  $\square_\kappa$ -sequence, then  $\square_\kappa$  is said to hold.

**Theorem 5.9.** *It is consistent with  $\aleph_1 < \mathfrak{h} < \mathfrak{s} < \text{cf}(\mathfrak{c}) = \mathfrak{c}$  that there is an  $(\mathfrak{h}, \mathfrak{s})$ -shattering family.*

PROOF: We start in a model of GCH satisfying  $\square_\kappa$  for some regular cardinal  $\kappa > \aleph_1$ . Choose any regular  $\lambda > \kappa^+$ . Fix a  $\square_\kappa$ -sequence  $\{C_\alpha : \alpha \in (\kappa^+)'\}$ . We may assume that  $C_\alpha = \alpha$  for all  $\alpha \in \kappa'$ . For each  $\alpha \in (\kappa^+)'$ , let  $o(C_\alpha)$  denote the order-type of  $C_\alpha$ . When  $C'_\alpha$  is bounded in  $\alpha$  with  $\eta = \max(C'_\alpha)$ , then let  $\{\varphi_l^\alpha : l \in \omega\}$  enumerate  $C_\alpha \setminus \eta$  in increasing order.

We will construct a  $\kappa \times \kappa^+$ -matrix of posets,  $\langle P_{\alpha,\xi} : \alpha \leq \kappa, \xi < \kappa^+ \rangle \in H(\lambda^+)$  and prove that the poset  $P_{\kappa,\kappa^+}$  as in Lemma 5.2 has the desired properties. For each  $\xi < \eta i < \kappa^+$ , we will also choose an  $\iota(\xi, \eta) < \kappa$  satisfying, as in (3) of the definition of  $\kappa \times (\xi + 1)$ -matrix that  $P_{\alpha,\xi} \dot{<} P_{\alpha,\eta}$  for all  $\iota(\xi, \eta) \leq \alpha < \kappa$ . We construct this family by constructing  $\langle P_{\alpha,\xi} : \alpha \leq \kappa, \xi < \zeta \rangle$  by recursion on limit  $\zeta < \kappa^+$ .

We will recursively define two other families. For each  $\alpha < \kappa$  and  $\xi < \kappa^+$ , we will define a set  $\text{supp}(P_{\alpha,\xi}) \subset \xi$  that can be viewed as the union of the supports of the elements of  $P_{\alpha,\xi}$  and will satisfy that  $\{\text{supp}(P_{\alpha,\xi}) : \alpha < \kappa\}$  is increasing and covers  $\xi$ . For each limit  $\eta < \kappa^+$  of cofinality less than  $\kappa$  and each  $n \in \omega$ , we will select a canonical  $P_{\kappa,\eta+n+1}$ -name,  $\dot{a}_{\eta+n}$  of a subset  $\omega$  that is forced to be Cohen over the forcing extension by  $P_{\kappa,\eta}$ . While this condition looks awkward, we simply want to avoid this task at limits of cofinality  $\kappa$ . Needing notation for this, let  $E = \kappa^+ \setminus \bigcup \{[\eta, \eta + \omega) : \text{cf}(\eta) = \kappa\}$ .

For each  $\alpha < \kappa$  and  $\xi < \eta < \kappa^+$ , we define  $\mathcal{A}_{\alpha,\xi,\eta}$  to be the family  $\{\dot{a}_\gamma : \gamma \in E \cap \eta \setminus \text{supp}(P_{\alpha,\xi})\}$ . The intention is that for all  $\alpha < \xi \leq \eta$ ,  $\mathcal{A}_{\alpha,\xi,\eta}$  is a family of  $P_{\kappa,\eta}$ -names which is forced by the poset  $P_{\kappa,\eta}$  to be thin over the forcing extension by  $P_{\alpha,\xi}$ . Let us note that if  $\alpha < \beta$  and  $\xi \leq \eta$ , then  $\mathcal{A}_{\alpha,\xi,\eta} \setminus \mathcal{A}_{\beta,\eta,\eta}$

should then be a set of  $P_{\beta,\eta}$ -names. By ensuring that  $\text{supp}(P_{\alpha,\xi})$  has cardinality less than  $\kappa$  for all  $\alpha < \kappa$  and  $\xi < \kappa^+$ , this will ensure that the family  $\{\dot{a}_\eta : \eta \in E\}$  is  $(\kappa, \kappa^+)$ -shattering. For each  $\eta < \kappa^+$  with cofinality  $\kappa$  we will ensure that  $P_{\kappa,\eta+1}$  has the form  $P_{\kappa,\eta} * \mathcal{C}_{\kappa \times \lambda}$  and that  $P_{\kappa,\eta+2} = P_{\kappa,\eta+1} * \mathbb{L}(\dot{D}_{\kappa,\eta})$  for a  $P_{\kappa,\eta+1}$ -name  $\dot{D}_{\kappa,\eta}$  of an ultrafilter on  $\omega$ . This will ensure that  $\mathfrak{c} \geq \lambda$  and  $\mathfrak{s} = \kappa^+$ . The sequence defining  $P_{\kappa,\eta+3}$  will be devoted to ensuring that  $\mathfrak{p} \geq \kappa$ .

We start the recursion in a rather trivial fashion. For each  $\alpha < \kappa$ ,  $P_{\alpha,0} = \mathcal{C}_\omega$  and for each  $n \in \omega$ ,  $P_{\alpha,n+1} = P_{\alpha,n} * \mathcal{C}_\omega$ . We may also let  $\iota(n, m) = 0$  for all  $n < m < \omega$ . For each  $n \in \omega$ , let  $\dot{a}_n$  be the canonical name of the Cohen real added by the second coordinate of  $P_{\kappa,n+1} = P_{\kappa,n} * \mathcal{C}_\omega$ . For each  $\alpha < \kappa$  and  $n \in \omega$ , define  $\text{supp}(P_{\alpha,n})$  to be  $n$ .

It should be clear that  $P_{\kappa,\omega}$  forces that for each  $\alpha < \kappa$  and  $n \in \omega$ , the family  $\{\dot{a}_m : n \leq m \in \omega\}$  is thin over the forcing extension by  $P_{\alpha,n}$ . Assume that  $P$  is a poset whose elements are functions with domain a subset of an ordinal  $\xi$ . We adopt the notational convention that for a  $P$ -name  $\dot{Q}$  for a poset,  $P * \xi \dot{Q}$  will denote the representation of  $P * \dot{Q}$  whose elements have the form  $p \cup \{(\xi, \dot{q})\}$  for  $(p, \dot{q}) \in P * \dot{Q}$ .

We will prove, by induction on limit  $\zeta < \kappa^+$ , there is a  $\kappa \times (\zeta + 1)$ -matrix  $\{P_{\alpha,\xi} : \alpha \leq \kappa, \xi \leq \zeta\}$  and families  $\{\mathcal{A}_{\alpha,\xi,\eta} : \alpha < \kappa, \xi \leq \eta \leq \zeta\}$  satisfying conditions (1)–(10).

- (1) For all  $\alpha < \beta < \kappa$  and  $\xi < \eta < \zeta$ , if  $P_{\alpha,\xi} < P_{\beta,\eta}$ , then the poset  $P_{\beta,\eta}$  forces that the family  $\mathcal{A}_{\alpha,\xi,\eta} \setminus \mathcal{A}_{\beta,\eta,\eta}$  is thin over the forcing extension by  $P_{\alpha,\xi}$ ;
- (2) for all  $\alpha < \kappa$  and  $\xi < \zeta$ , the elements  $p$  of the poset  $P_{\alpha,\xi}$  are functions that have a finite domain,  $\text{dom}(p)$ , contained in  $\xi$ ;
- (3) if  $C'_\zeta$  is cub in  $\zeta$  and  $\eta \in C'_\zeta$ , then
  - (a)  $P_{n,\zeta}$  is the trivial poset and  $\text{supp}(P_{n,\zeta}) = \emptyset$  for  $n \in \omega$ ;
  - (b)  $P_{\alpha,\zeta} = P_{\alpha,\eta}$  and  $\text{supp}(P_{\alpha,\zeta}) = \text{supp}(P_{\alpha,\eta})$  for all  $o(C_\eta) \leq \alpha < o(C_\eta) + \omega$ ; and
  - (c)  $P_{\alpha,\zeta} = \bigcup \{P_{\alpha,\eta} : \eta \in C'_\zeta\}$  and  $\text{supp}(P_{\alpha,\zeta}) = \bigcup \{\text{supp}(P_{\alpha,\eta}) : \eta \in C'_\zeta\}$  for all  $o(C_\zeta) \leq \alpha < \kappa$ ;

also, let  $\iota(\eta, \zeta) = o(C_\eta)$  for all  $\eta \in C'_\zeta$  and for all  $\gamma < \zeta \setminus C'_\zeta$  let  $\iota(\gamma, \zeta) = \iota(\gamma, \eta)$  where  $\eta = \min(C'_\zeta \setminus \gamma)$ ;

- (4) if  $\max(C'_\zeta) < \zeta$  then let

$$\iota_\zeta = \max(o(C_\zeta), \sup\{\iota(\varphi_l^\zeta, \varphi_{l'}^\zeta + n) : l \leq l' < n < \omega\})$$

and

- (a) set  $P_{\alpha,\zeta} = P_{\alpha,\varphi_0^\zeta}$  and  $\text{supp}(P_{\alpha,\zeta}) = \text{supp}(P_{\alpha,\varphi_0^\zeta})$  for all  $\alpha < \iota_\zeta$ ;
- (b) set, for  $\iota_\zeta \leq \alpha < \kappa$ ,  $P_{\alpha,\zeta} = \bigcup \{P_{\alpha,\varphi_l^\zeta + n} : l, n \in \omega\}$  and  $\text{supp}(P_{\alpha,\zeta}) = \bigcup \{\text{supp}(P_{\alpha,\varphi_l^\zeta + n}) : l, n \in \omega\}$ ;

- (c) for each  $\gamma \in \varphi_0^\zeta$  let  $\iota(\gamma, \zeta) = \iota(\gamma, \varphi_0^\zeta)$ , let  $\iota(\varphi_0^\zeta, \zeta) = o(C_\gamma)$ , and for each  $\varphi_0^\zeta < \gamma < \zeta$ ,  $\iota(\gamma, \zeta)$  is the maximum of  $\iota_\zeta$  and  $\min\{\iota(\gamma, \varphi_l^\zeta + n) : l, n \in \omega \text{ and } \gamma < \varphi_l^\zeta + n\}$ ;
- (5) if  $o(C_\zeta) < \kappa$ , then for all  $\alpha < \kappa$  and  $n \in \omega$ :
- (a)  $P_{\alpha, \zeta+n+1} = P_{\alpha, \zeta+n} *_{\zeta+n} \mathcal{C}_\omega$ ;
  - (b)  $\dot{a}_{\zeta+n}$  in the canonical  $P_{0, \zeta+n} *_{\zeta+n} \mathcal{C}_\omega$ -name for the Cohen real added by the second coordinate copy of  $\mathcal{C}_\omega$ ;
  - (c)  $\text{supp}(P_{\alpha, \zeta+n+1}) = \text{supp}(P_{\alpha, \zeta}) \cup [\zeta, \zeta + n]$ ; and
  - (d)  $\iota(\zeta + k, \zeta + n + 1) = 0$  for all  $k \leq n$ , and for all  $\gamma < \zeta$ ,  $\iota(\gamma, \zeta + n + 1) = \iota(\gamma, \zeta)$ ;
- (6) if  $o(C_\zeta) = \kappa$ , then for all  $\alpha < \kappa$ ,  $P_{\alpha, \zeta+1} = P_{\alpha, \zeta} *_{\zeta} \mathcal{C}_{\alpha+1 \times \lambda}$ ;
- (7) if  $o(C_\zeta) = \kappa$ , then for all  $n \in \omega$  and all  $\alpha < \kappa$ ,  $P_{\alpha, \zeta+3+n} = P_{\alpha, \zeta+3}$ ;
- (8) if  $o(C_\zeta) = \kappa$ , then there is an  $\iota_\zeta < \kappa$  such that  $P_{\beta, \zeta+2} = P_{\beta, \zeta+1}$  for all  $\beta < \iota_\zeta$ , and there is a sequence  $\langle \dot{D}_{\alpha, \zeta} : \iota_\zeta \leq \alpha < \kappa \rangle$  such that for each  $\iota_\zeta \leq \alpha < \kappa$ :
- (a)  $\dot{D}_{\alpha, \zeta}$  is a  $P_{\alpha, \kappa+1}$ -name of a Ramsey ultrafilter on  $\omega$ ;
  - (b) for each  $\iota_\zeta \leq \beta < \alpha$ ,  $\dot{D}_{\beta, \zeta} \subset \dot{D}_{\alpha, \zeta}$ ;
  - (c)  $P_{\alpha, \zeta+2} = P_{\alpha, \zeta+1} *_{\zeta+1} \mathbb{L}(\dot{D}_{\alpha, \kappa})$ ;
- (9) if  $o(C_\zeta) = \kappa$ , then for  $\iota_\zeta$  chosen as in (8)
- (a) for each  $\alpha < \iota_\zeta$ ,  $P_{\alpha, \kappa+3} = P_{\alpha, \kappa+2}$ ;
  - (b)  $P_{\iota_\zeta, \zeta+3} = P_{\iota_\zeta, \zeta+2} *_{\zeta+2} \dot{Q}_{\iota_\zeta, \zeta+2}$  for some  $P_{\iota_\zeta, \zeta}$ -name,  $\dot{Q}_{\iota_\zeta, \zeta+2}$  in  $H(\lambda^+)$  of a finite support product of  $\sigma$ -centered posets;
  - (c) for each  $\iota_\zeta < \alpha < \kappa$ ,  $P_{\alpha, \zeta+3} = P_{\alpha, \zeta+2} *_{\zeta+2} \dot{Q}_{\iota_\zeta, \zeta+2}$ ;
- (10) if  $o(C_\zeta) = \kappa$ , then for all  $\alpha < \kappa$ ,  $n \in \omega$ , and  $\gamma < \zeta$ ,  $\text{supp}(P_{\alpha, \zeta+n+1}) = \text{supp}(P_{\alpha, \zeta}) \cup [\zeta, \zeta+n]$ ,  $\iota(\gamma, \zeta+n) = \iota(\gamma, \zeta)$ , and  $\iota(\zeta+k, \zeta+n) = \iota_\zeta$  for all  $k < n \in \omega$ .

It should be clear from the properties, and by induction on  $\zeta$ , that for all  $\alpha < \kappa$  and  $\xi < \zeta$ , each  $p \in P_{\alpha, \xi}$  is a function with finite domain contained in  $\text{supp}(P_{\alpha, \xi})$ . Similarly, it is immediate from the hypotheses that  $\text{supp}(P_{\alpha, \xi})$  has cardinality less than  $\kappa$  for all  $(\alpha, \xi) \in \kappa \times \kappa^+$ .

Before verifying the construction, we first prove, by induction on  $\zeta$ , that the conditions (2)–(10) ensure that for all  $\xi \leq \zeta$  and  $\eta \in C'_\xi$ :

Claim (a):  $P_{\alpha, \eta} < P_{\alpha, \xi}$  for all  $o(C_\eta) + \omega \leq \alpha < \kappa$ .

Claim (b):  $P_{\alpha, \eta} = P_{\alpha, \xi}$  for all  $\alpha < o(C_\eta) + \omega$ .

If  $o(C_\xi) \leq \alpha$ , then  $P_{\alpha, \eta} < P_{\alpha, \xi}$  follows immediately from clause 2 (c) and, by induction, clauses 3 (a). Now assume  $\alpha < o(C_\xi) + \omega$ . If  $C'_\xi$  is not cofinal in  $\xi$ , then, by induction,  $P_{\alpha, \eta} = P_{\alpha, \varphi_0^\xi}$  and, by clause 3 (a),  $P_{\alpha, \varphi_0^\xi} = P_{\alpha, \xi}$ . If  $C'_\xi$  is cofinal in  $\xi$ , then choose  $\bar{\eta} \in C'_\xi$  so that  $o(C_{\bar{\eta}}) \leq \alpha < o(C_{\bar{\eta}}) + \omega$ . By clause 2 (b),  $P_{\alpha, \xi} = P_{\alpha, \bar{\eta}}$ . By the inductive assumption,  $P_{\alpha, \eta} = P_{\alpha, \bar{\eta}}$  since one of  $\eta = \bar{\eta}$ ,  $\eta \in C'_\eta$  or  $\bar{\eta} \in C'_\eta$  must hold.

The second thing we check is that the conditions (2)–(10) also ensure that for each  $\zeta < \kappa^+$ ,  $\langle P_{\alpha,\eta} : \alpha \leq \kappa, \eta \leq \zeta \rangle$  is a  $\kappa \times \zeta$ -matrix. We assume, by induction on limit  $\zeta$ , that for  $\gamma < \eta < \zeta$ ,  $\{P_{\alpha,\gamma} : \alpha \leq \kappa\}$  is a  $<\cdot$ -chain and that  $P_{\alpha,\gamma} < P_{\alpha,\eta}$  for all  $\eta$  with  $\iota(\gamma, \eta) \leq \alpha \leq \kappa$ . Note that clauses 3 (c) and 4 (b) of the construction ensure that condition (4) of Definition 5.1 holds. We check the details for  $\zeta + 1$  and skip the easy subsequent verification for  $\zeta + n$ ,  $n \in \omega$ . Suppose first that  $C'_\zeta$  is cofinal in  $\zeta$  and let  $\iota(\gamma, \zeta) \leq \alpha < \kappa$  for some  $\gamma < \zeta$ . Of course we may assume that  $\gamma \notin C'_\zeta$ . Since  $C'_\zeta$  is cofinal in  $\zeta$ , let  $\eta = \min(C'_\zeta \setminus \gamma)$ . By induction,  $P_{\alpha,\gamma} < P_{\alpha,\eta} < P_{\alpha,\zeta}$ . Now assume that  $C'_\zeta$  is not cofinal in  $\zeta$ . If  $\gamma \leq \varphi_0^\zeta$ , then  $\iota(\gamma, \zeta) = \iota(\gamma, \varphi_0^\zeta)$ , and so we have that  $P_{\alpha,\gamma} < P_{\alpha,\varphi_0^\zeta} < P_{\alpha,\zeta}$ . If  $\varphi_0^\zeta < \gamma$ , then choose any  $l \in \omega$  so that  $\gamma < \varphi_l^\zeta$ . By construction,  $\iota(\gamma, \zeta) \geq \iota(\gamma, \varphi_l^\zeta)$  and so for  $\iota(\gamma, \zeta) \leq \alpha < \kappa$ ,  $P_{\alpha,\gamma} < P_{\alpha,\varphi_l^\zeta} < P_{\alpha,\zeta}$ .

Now we consider the values of  $\mathcal{A}_{\alpha,\xi,\eta}$  for  $\alpha < \kappa$  and  $\omega \leq \xi \leq \eta$  by examining the names  $\dot{a}_\gamma$  for  $\gamma \in E$ .

By clause (5),  $\dot{a}_\gamma$  is a  $P_{0,\gamma+1}$ -name and  $\gamma$  is in the domain of each  $p \in P_{0,\gamma+1}$  appearing in the name. One direction of this next claim is then obvious given that the domain of every element of  $P_{\alpha,\xi}$  is a subset of  $\text{supp}(P_{\alpha,\xi})$ .

Claim (c):  $\dot{a}_\gamma$  is a  $P_{\alpha,\xi}$ -name if and only if  $\gamma \in \text{supp}(P_{\alpha,\xi})$ .

Assume that  $\gamma \in \text{supp}(P_{\alpha,\xi})$ . We prove this by induction on  $\xi$ . If  $\xi$  is a limit, then  $\text{supp}(P_{\alpha,\xi})$  is defined as a union, hence there is an  $\eta < \xi$  such that  $\gamma \in \text{supp}(P_{\alpha,\eta})$  and  $P_{\alpha,\eta} < P_{\alpha,\xi}$ . If  $\xi = \eta + n$  for some limit  $\eta$  and  $n \in \omega$ , then  $P_{\alpha,\eta} < P_{\alpha,\xi}$  and so we may assume that  $\eta \leq \gamma = \eta + k < \eta + n$  and that  $o(C_\eta) < \kappa$ . Since  $P_{0,\eta+k} < P_{\alpha,\eta+k} < P_{\alpha,\eta+n} = P_{\alpha,\xi}$ , it follows that  $\dot{a}_\gamma$  is a  $P_{\alpha,\xi}$ -name.

We prove by induction on  $\xi$ ,  $\xi$  a limit, that for all  $\gamma < \xi$ :

Claim (d): for all  $\alpha < \iota(\gamma + 1, \xi)$ ,  $\gamma$  is not in  $\text{supp}(P_{\alpha,\xi})$ .

First consider the case that  $C'_\xi$  is cofinal in  $\xi$  and let  $\eta$  be the minimum element of  $C'_\xi \setminus (\gamma + 1)$ . By definition  $\iota(\gamma + 1, \xi)$  is equal to  $\iota(\gamma + 1, \eta)$  and the claim follows since we have that  $\text{supp}(P_{\iota(\gamma+1,\xi),\zeta}) = \text{supp}(P_{\iota(\gamma+1,\xi),\eta})$ . Now assume that  $C'_\xi$  is not cofinal in  $\xi$  and assume that  $\alpha < \iota(\gamma + 1, \xi)$ . We break into cases:  $\gamma < \varphi_0^\xi$  and  $\varphi_0^\xi \leq \gamma < \xi$ . In the first case  $\iota(\gamma, \xi) = \iota(\gamma, \varphi_0^\xi)$  and the claim follows by induction and the fact that  $\text{supp}(P_{\alpha,\varphi_0^\xi}) = \text{supp}(P_{\alpha,\xi})$  for all  $\alpha < \iota(\gamma, \xi)$ . Now consider  $\varphi_0^\xi \leq \gamma < \xi$ . If  $\alpha < \iota_\xi$ , then  $P_{\alpha,\xi} = P_{\alpha,\varphi_0^\xi}$  and, since  $\iota_\xi \leq \iota(\gamma + 1, \xi)$ ,  $\gamma$  is not in  $\text{supp}(P_{\alpha,\varphi_0^\xi})$ . Otherwise, choose  $l, n \in \omega$  so that  $\iota_\xi \leq \alpha < \iota(\gamma + 1, \xi) = \iota(\gamma + 1, \varphi_l^\xi + n)$  as in the definition of  $\iota(\gamma, \xi)$ . By the minimality in the choice of  $\varphi_l^\xi + n$ , it follows that  $\gamma$  is not in  $\text{supp}(P_{\alpha,\varphi_l^\xi+n})$  for

all  $l', n \in \omega$ . Since  $\text{supp}(P_{\alpha,\xi})$  is the union of all such sets, it follows that  $\gamma$  is not in  $\text{supp}(P_{\alpha,\xi})$ .

Next we prove, by induction on  $\zeta$ , that the matrix so chosen will additionally satisfy condition (1). We first find a reformulation of condition (1). Note that by Claim (c),  $\mathcal{A}_{\alpha,\xi,\eta} = \{\dot{a}_\gamma : \gamma \in E \cap \eta \setminus \text{supp}(P_{\alpha,\xi})\}$ .

Claim (e): For each  $\alpha < \kappa$  and  $\xi < \eta < \zeta$  and finite subset  $\{\gamma_i : i < m\}$  of  $E \cap \eta \setminus \text{supp}(P_{\alpha,\xi})$  there is a  $\beta < \kappa$  such that  $\iota(\xi, \eta) \leq \beta$ ,  $\{\gamma_i : i < m\} \subset \text{supp}(P_{\beta,\eta})$  and  $P_{\beta,\eta}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the forcing extension by  $P_{\alpha,\xi}$ .

Let us verify that Claim (e) follows from condition (1). Let  $\alpha, \xi, \eta$  and  $\{\gamma_i : i < m\}$  be as in the statement of Claim (e). Choose  $\beta < \kappa$  so that  $\iota(\xi, \eta)$  and each  $\iota(\gamma_i + 1, \eta)$  is less than  $\beta$ . Then  $P_{\alpha,\xi} \dot{<} P_{\beta,\eta}$  and  $\{\dot{a}_{\gamma_i} : i < m\} \subset \mathcal{A}_{\alpha,\xi,\eta} \setminus \mathcal{A}_{\beta,\eta,\eta}$ . This value of  $\beta$  satisfies the conclusion of Claim (e).

Now assume that Claim (e) holds and we prove that condition (1) holds. Assume that  $P_{\alpha,\xi} \dot{<} P_{\delta,\eta}$ . To prove that  $\mathcal{A}_{\alpha,\xi,\eta} \setminus \mathcal{A}_{\delta,\eta,\eta}$  is forced by  $P_{\delta,\eta}$  to be thin over the forcing extension by  $P_{\alpha,\xi}$ , it suffices to prove this for any finite subset of  $\mathcal{A}_{\alpha,\xi,\eta} \setminus \mathcal{A}_{\delta,\eta,\eta}$ . Thus, let  $\{\gamma_i : i < m\}$  be any finite subset of  $\text{supp}(P_{\delta,\eta}) \cap E \cap \eta \setminus \text{supp}(P_{\alpha,\xi})$ . Choose  $\beta$  as in the conclusion of the claim. If  $\beta \leq \delta$ , then  $P_{\delta,\eta}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the forcing extension because  $P_{\beta,\eta} \dot{<} P_{\delta,\eta}$  does. Similarly, if  $\delta < \beta$ , then  $P_{\delta,\eta}$  being completely embedded in  $P_{\beta,\eta}$  cannot force that  $\{\dot{a}_{\gamma_i} : i < m\}$  is not thin over the forcing extension by  $P_{\alpha,\xi}$ .

We assume that  $\omega \leq \zeta < \kappa^+$  is a limit and that  $\langle P_{\alpha,\xi} : \alpha \leq \kappa, \xi < \zeta \rangle$  have been chosen so that conditions (1)–(10) are satisfied. We prove, by induction on  $n \in \omega$ , that there is an extension  $\langle P_{\alpha,\xi} : \alpha \leq \kappa, \xi < \zeta + n \rangle$  that also satisfies conditions (1)–(10).

For  $n = 1$ , we define the sequence  $\langle P_{\alpha,\zeta} : \alpha < \kappa \rangle$  according to the requirement of (3) or (4) as appropriate. It follows from Lemma 5.7 that (2) will hold for the extension  $\langle P_{\alpha,\xi} : \alpha < \kappa, \xi < \zeta + 1 \rangle$ . Conditions (3)–(10) hold since there are no new requirements. We must verify that the condition in Claim (e) holds for  $\eta = \zeta$ . Let  $\alpha, \xi$  and  $\{\gamma_i : i < m\}$  be as in the statement of Claim (e) with  $\eta = \zeta$ . Let  $C_\zeta = \{\eta_\beta : \beta < o(C_\zeta)\}$  be an order-preserving enumeration. We first deal with case that  $C'_\zeta$  is cofinal in  $\zeta$ . Choose any  $\beta_0 < \kappa$  large enough so that  $\gamma_i \in \text{supp}(P_{\beta_0,\zeta})$  for all  $i < m$ . Choose  $\beta_0 < \beta$  so that  $\iota(\xi, \eta_{\beta_0}) \leq \beta$ . Now we have that  $P_{\alpha,\xi} \dot{<} P_{\beta,\eta_{\beta_0}}$  and  $P_{\beta,\eta_{\beta_0}} \dot{<} P_{\beta,\zeta}$ . Applying Claim (e) to  $\eta_{\beta_0}$ , we have that  $P_{\beta,\eta_{\beta_0}}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the forcing extension by  $P_{\alpha,\xi}$ . As in the proof of Claim (e), this implies that  $P_{\beta,\zeta}$  forces the same thing.

Now the case that  $C'_\zeta$  is not cofinal in  $\zeta$ . If  $\alpha < \iota_\zeta$ , then apply Claim (e) to choose  $\beta$  so that  $P_{\beta, \iota_\zeta}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is not thin over the extension by  $P_{\alpha, \xi}$ . Since  $P_{\beta, \iota_\zeta} < P_{\beta, \zeta}$  holds for all  $\beta$ ,  $P_{\beta, \zeta}$  also forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is not thin over the extension by  $P_{\alpha, \xi}$ . If  $\iota_\zeta \leq \alpha$ , first choose  $\delta < \kappa$  large enough so that  $\iota(\xi, \zeta)$  and each  $\iota(\gamma_i + 1, \zeta)$  is less than  $\delta$ . Since  $\{\gamma_i : i < m\}$  is a subset of  $\text{supp}(P_{\delta, \zeta})$ , we can choose  $l < \omega$  large enough so that  $\{\gamma_i : i < \omega\} \subset \text{supp}(P_{\delta, \varphi_l^\zeta})$ . Applying Claim (e) to  $\eta = \varphi_l^\zeta$ , we choose  $\beta$  as in the claim. As we have seen, there is no loss to assuming that  $\delta \leq \beta$  and, since  $P_{\beta, \varphi_l^\zeta} < P_{\beta, \zeta}$ , this completes the proof.

If  $o(C'_\zeta) < \kappa$ , then the construction of  $\langle P_{\alpha, \zeta+n} : n \in \omega, \alpha < \kappa \rangle$  is canonical so that conditions (2)–(10) hold. We again verify that Claim (e) holds for all values of  $\eta$  with  $\zeta < \eta < \zeta + \omega$ . Let  $\alpha, \xi$  and  $\{\gamma_i : i < m\}$  be as in Claim (e) for  $\eta = \zeta + n$ . We may assume that  $\{\gamma_i : i < m\} \cap \zeta = \{\gamma_i : i < \overline{m}\}$  for some  $\overline{m} \leq m$ . If  $\xi < \zeta$ , let  $\bar{\xi} = \xi$ , otherwise, choose any  $\bar{\xi} < \zeta$  so that  $P_{\alpha, \zeta} = P_{\alpha, \bar{\xi}}$ . Note that  $\{\gamma_i : \overline{m} \leq i < m\}$  is disjoint from the interval  $[\zeta, \xi)$ . Choose  $\beta < \kappa$  to be greater than  $\iota(\bar{\xi}, \zeta)$  and each  $\iota(\gamma_i + 1, \zeta)$ ,  $i < \overline{m}$ , and so that  $P_{\beta, \zeta}$  forces that  $\{\dot{a}_{\gamma_i} : i < \overline{m}\}$  is thin over the extension by  $P_{\alpha, \bar{\xi}}$ . If  $\overline{m} = m$  we are done by the fact that  $P_{\alpha, \xi}$  is isomorphic to  $P_{\alpha, \bar{\xi}} * \mathcal{C}_\omega$ . In fact, we similarly have that  $P_{\beta, \xi}$  forces that  $\{\dot{a}_{\gamma_i} : i < \overline{m}\}$  is thin over the forcing extension by  $P_{\alpha, \xi}$ . Since  $P_{\beta, \zeta+n}$  forces that  $\bigcup \{\dot{a}_{\gamma_i} : \overline{m} \leq i < m\}$  is a Cohen real over the forcing extension by  $P_{\beta, \xi}$  it also follows that  $P_{\beta, \zeta+n}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the extension by  $P_{\alpha, \xi}$ .

Now we come to the final case where  $o(C'_\zeta) = \kappa$  and the main step to the proof. The fact that Claim (e) will hold for  $\eta = \zeta + 1$  is proven as above for the case when  $o(C'_\zeta) < \kappa$  and  $C'_\zeta$  is cofinal in  $\zeta$ . For values of  $n > 3$ , there is nothing to prove since  $P_{\alpha, \zeta+3+k} = P_{\alpha, \zeta+3}$  for all  $k \in \omega$ . We also note that  $\zeta + n \notin E$  for all  $n \in \omega$ .

At step  $\eta = \zeta + 2$  we must take great care to preserve Claim (e) and at step  $\zeta + 3$  we make a strategic choice towards ensuring that  $\mathfrak{p}$  will equal  $\kappa$ . Indeed, we begin by choosing the lexicographic minimal pair,  $(\xi_\zeta, \alpha_\zeta)$ , in  $\zeta \times \kappa$  with the property that there is a family of fewer than  $\kappa$  many canonical  $P_{\alpha_\zeta, \xi_\zeta}$ -names of subsets of  $\omega$  and a  $p \in P_{\alpha_\zeta, \xi_\zeta}$  that forces over  $P_{\kappa, \zeta}$  that there is no pseudo-intersection. If there is no such pair, then let  $(\alpha_\zeta, \xi_\zeta) = (\omega, \zeta + 1)$ . Choose  $\iota_\zeta$  so that  $P_{\alpha_\zeta, \xi_\zeta} < P_{\iota_\zeta, \zeta+1}$ .

Assume that  $\alpha, \xi, \{\gamma_i : i < m\}$  are as in Claim (e). We first check that if  $\xi < \zeta + 2$ , then there is nothing new to prove. Indeed, simply choose  $\beta < \kappa$  large enough so that  $P_{\beta, \zeta+1}$  has the properties required in Claim (e) for  $P_{\alpha, \xi}$ . Of course it follows that  $P_{\beta, \zeta+2}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the extension by  $P_{\alpha, \xi}$  since  $P_{\beta, \zeta+1}$  already forces this.

This means that we need only consider instances of Claim (e) in which  $\xi = \zeta + 2$ . The analogous statement also holds when we move to  $\zeta + 3$ . For each  $\beta < \kappa$ , let

$$T_\beta = E \cap \text{supp}(P_{\beta+1, \zeta}) \setminus \text{supp}(P_{\beta, \zeta})$$

and note that  $P_{\beta+1, \zeta+1}$  forces that  $\{\dot{a}_\gamma : \gamma \in T_\beta\}$  is thin over the extension by  $P_{\beta, \zeta+1}$ . Most of the work has been done for us in Lemma 5.6. Except for some minor re-indexing, we can assume that the sequence  $\{P_\beta : \beta < \kappa\}$  in the statement of Lemma 5.6 is the sequence  $\{P_{\beta, \zeta} : \beta < \kappa\}$ . We also have that  $P_{\beta, \zeta} * \mathcal{C}_{(\beta+1) \times \kappa}$  is isomorphic to  $P_{\beta, \zeta+1}$ . We can choose any  $P_{0, \zeta+1}$ -name  $\dot{D}_{0, \zeta}$ -name of a Ramsey ultrafilter on  $\omega$ . The family  $\{\dot{a}_\gamma : \gamma \in T_\beta\}$  will play the role of  $\dot{A}_\beta$  in the statement of Lemma 5.6, and we let  $\{\dot{D}_{\beta, \zeta} : 0 < \beta < \kappa\}$  be the sequence as supplied in Lemma 5.6.

Now assume that  $\alpha < \kappa$  and that  $\{\gamma_i : i < m\} \subset E \cap \zeta \setminus \text{supp}(P_{\alpha, \zeta+1})$ . Let  $\{\dot{F}_l : l \in \omega\}$  be any sequence of  $P_{\alpha, \zeta+2}$ -names of pairwise disjoint elements of  $[\omega]^k$  for some  $k \in \omega$ . We must find a sufficiently large  $\beta < \kappa$  so that  $P_{\beta, \zeta+2}$  forces that  $\dot{a}_{\gamma_0} \cup \dots \cup \dot{a}_{\gamma_{m-1}}$  is disjoint from  $\dot{F}_l$  for some  $l \in \omega$ . Let  $\{\beta_j : j < \overline{m}\}$  be the set (listed in increasing order) of  $\beta < \kappa$  such that  $T_\beta \cap \{\gamma_i : i < m\}$  is not empty and let  $\beta_m = \beta_{m-1} + 1$ . By re-indexing we can assume there is a sequence  $\{m_j : j \leq \overline{m}\} \subset m+1$  so that  $\gamma_i \in T_{\beta_j}$  for  $m_j \leq i < m_{j+1}$ . Although  $P_{\beta, \zeta+2} = P_{\beta, \zeta+1}$  for values of  $\beta < \iota_\zeta$ , we will let  $\overline{P}_{\beta, \zeta+2} = P_{\beta, \zeta+1} *_{\zeta+1} \mathbb{L}(\dot{D}_{\beta, \zeta})$  for  $\beta < \iota_\zeta$ , and for consistent notation, let  $\overline{P}_{\beta, \zeta+2} = P_{\beta, \zeta+2}$  for  $\iota_\zeta \leq \beta < \kappa$ . We note that  $\{\dot{F}_l : l \in \omega\}$  is also sequence of  $\overline{P}_{\alpha, \zeta+2}$ -names of pairwise disjoint elements of  $[\omega]^k$ .

For each  $j < \overline{m}$ , let  $\dot{L}_{j+1}$  be the  $\overline{P}_{\beta_j+1, \zeta+2}$ -name of those  $l$  such that  $\dot{F}_l$  is disjoint from  $\bigcup\{\dot{a}_{\gamma_i} : i < m_{j+1}\}$ . It follows, by induction on  $j < \overline{m}$ , that  $\overline{P}_{\beta_j+1, \zeta+2}$  forces that  $\dot{L}_{j+1}$  is infinite since  $\overline{P}_{\beta_j+1, \zeta+2}$  forces that  $\{\dot{a}_{\gamma_i} : m_j \leq i < m_{j+1}\}$  is thin over the forcing extension by  $\overline{P}_{\beta_j, \zeta+2}$ . It now follows  $\overline{P}_{\beta_m, \zeta+2}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the forcing extension by  $\overline{P}_{\alpha, \zeta+2}$ . If  $\beta_m < \iota_\zeta$ , let  $\beta = \iota_\zeta$ , otherwise, let  $\beta = \beta_m$ . It follows that  $P_{\beta, \zeta+2}$  forces that  $\{\dot{a}_{\gamma_i} : i < m\}$  is thin over the forcing extension by  $P_{\alpha, \zeta+2} < \overline{P}_{\alpha, \zeta+2}$ . This completes the verification of Claim (e) for the case  $\eta = \zeta + 2$  and we now turn to the final case of  $\eta = \zeta + 3$ .

We have chosen the pair  $(\alpha_\zeta, \xi_\zeta)$  when choosing  $\iota_\zeta$ . Let  $\dot{Q}_{\iota_\zeta, \zeta+2}$  be the  $P_{\iota_\zeta, \zeta+2}$ -name of the finite support product of all posets of the form  $\mathbb{M}(\mathcal{F})$  where  $\mathcal{F}$  is a family of fewer than  $\kappa$  canonical  $P_{\alpha_\zeta, \xi_\zeta}$ -names of subsets of  $\omega$  that is forced to have the sfip. Since  $P_{\alpha_\zeta, \xi_\zeta} \in H(\lambda^+)$  the set of all such families  $\mathcal{F}$  is an element of  $H(\lambda^+)$ . This is our value of  $\dot{Q}_{\iota_\zeta, \zeta+2}$  as in condition (9) for the definition of  $P_{\beta, \zeta+3}$  for all  $\beta < \kappa$ . The fact that Claim (e) holds in this case follows immediately from the induction hypothesis and Proposition 5.5. We also note that  $P_{\iota_\zeta, \zeta+3}$

forces that every family of fewer than  $\kappa$  many canonical  $P_{\alpha_\zeta, \xi_\zeta}$ -names that is forced to have the sfip is also forced by  $P_{\kappa, \zeta+3}$  to have a pseudo-intersection. This means that for values of  $\zeta' > \zeta$  with  $o(C'_{\zeta'}) = \kappa$ , the pair  $(\alpha_{\zeta'}, \xi_{\zeta'})$  will be lexicographically strictly smaller than the choice for  $\zeta'$ . In other words, the family  $\{(\xi_\zeta, \alpha_\zeta) : \zeta < \kappa^+, \text{cf}(\zeta) = \kappa\}$  is strictly increasing in the lexicographic ordering.

Now we can verify that  $P_{\kappa, \kappa^+}$  forces that  $\mathfrak{p} \geq \kappa$ . If it does not, then there is a  $\delta < \kappa$  and a family,  $\{\dot{y}_\gamma : \gamma < \delta\}$  of canonical  $P_{\kappa, \kappa^+}$ -names of subsets of  $\omega$  with some  $p \in P_{\kappa, \kappa^+}$  forcing that the family has sfip but has no pseudo-intersection. By an easy modification of the names, we can assume that every condition in  $P_{\kappa, \kappa^+}$  forces that the family  $\{\dot{y}_\gamma : \gamma < \delta\}$  is forced to have sfip. Choose any  $\xi < \kappa^+$  so that  $p \in P_{\kappa, \xi}$  and every  $\dot{y}_\gamma$  is a  $P_{\kappa, \xi}$ -name. Choose  $\alpha < \kappa$  large enough so that  $p \in P_{\alpha, \xi}$ ,  $\iota(\zeta, \xi)$ , and each  $\alpha_\gamma$ ,  $\gamma < \delta$ , is less than  $\alpha$ . It follows that  $\dot{y}_\gamma$  is a  $P_{\alpha, \xi}$ -name for all  $\gamma < \delta$ . Since the family  $\{(\xi_\zeta, \alpha_\zeta) : \zeta < \kappa^+, \text{cf}(\zeta) = \kappa\}$  is strictly increasing in the lexicographic ordering, and this ordering on  $\kappa^+ \times \kappa$  has order type  $\kappa^+$ , there is a minimal  $\zeta < \kappa^+$  (with  $\text{cf}(\zeta) = \kappa$ ) such that  $(\xi, \alpha) \leq (\xi_\zeta, \alpha_\zeta)$ . By the assumption on  $(\alpha, \xi)$ ,  $(\xi_\zeta, \alpha_\zeta)$  will be chosen to equal  $(\xi, \alpha)$ . One of the factors of the poset  $\dot{Q}_{\iota_\zeta, \zeta+2}$  will be chosen to be  $\mathbb{M}(\{\dot{y}_\gamma : \gamma < \delta\})$ . This proves that  $P_{\kappa, \zeta+3}$  forces  $\{\dot{y}_\gamma : \gamma < \delta\}$  does have a pseudo-intersection.

It should be clear from condition (8) in the construction that  $P_{\kappa, \kappa^+}$  forces that  $\mathfrak{s} \geq \kappa^+$ . To finish the proof we must show that  $P_{\kappa, \kappa^+}$  forces that  $\{\dot{a}_\gamma : \gamma \in E\}$  is  $(\kappa, \kappa^+)$ -shattering. Since  $\dot{a}_\gamma$  is forced to be a Cohen real over the extension by  $P_{\kappa, \gamma}$ , condition (2) in Definition 2.3 of  $(\kappa, \kappa^+)$ -shattering holds. Finally, we verify condition (1) of Definition 2.3. Choose any  $P_{\kappa, \kappa^+}$ -name  $\dot{b}$  of an infinite subset of  $\omega$ . Choose any  $(\alpha, \xi) \in \kappa \times \kappa^+$  so that  $\dot{b}$  is a  $P_{\alpha, \xi}$ -name. The set  $E \cap \text{supp}(P_{\alpha, \xi})$  has cardinality less than  $\kappa$ . For any  $\gamma \in E \setminus \text{supp}(P_{\alpha, \xi})$ , there is a  $(\beta, \zeta) \in \kappa \times \kappa^+$  such that  $\{\dot{a}_\gamma\}$  is thin over the forcing extension by  $P_{\alpha, \xi}$ . It follows trivially that  $P_{\beta, \zeta}$  forces that  $\dot{b}$  is not a (mod finite) subset of  $\dot{a}_\gamma$ .  $\square$

## 6. Questions

- (1) Is it consistent to have  $\omega_1 < \mathfrak{h} < \mathfrak{b} < \mathfrak{s}$  and  $\mathfrak{c}$  regular?
- (2) Is it consistent to have  $\omega_1 < \mathfrak{h} < \mathfrak{s} < \mathfrak{b}$  and  $\mathfrak{c}$  regular?

Question (2) has been answered in [12].

## REFERENCES

- [1] Balcar B., Pelant J., Simon P., *The space of ultrafilters on  $\mathbb{N}$  covered by nowhere dense sets*, Fund. Math. **110** (1980), no. 1, pages 11–24.
- [2] Baumgartner J.E., Dordal P., *Adjoining dominating functions*, J. Symbolic Logic **50** (1985), no. 1, 94–101.

- [3] Blass A., *Applications of superperfect forcing and its relatives*, Conf. Set Theory and Its Applications, Toronto, 1987, Lecture Notes in Math., 1401, Springer, Berlin, 1989, pages 18–40.
- [4] Blass A., Shelah S., *Ultrafilters with small generating sets*, Israel J. Math. **65** (1989), no. 3, 259–271.
- [5] Brendle J., Fischer V., *Mad families, splitting families and large continuum*, J. Symbolic Logic **76** (2011), no. 1, 198–208.
- [6] Brendle J., Raghavan D., *Bounding, splitting, and almost disjointness*, Ann. Pure Appl. Logic **165** (2014), no. 2, 631–651.
- [7] Dow A., Shelah S., *On the cofinality of the splitting number*, Indag. Math. (N.S.) **29** (2018), no. 1, 382–395.
- [8] Dow A., Shelah S., *Pseudo P-points and splitting number*, Arch. Math. Logic **58** (2019), no. 7–8, 1005–1027.
- [9] Fischer V., Friedman S. D., Mejía D. A., Montoya D. C., *Coherent systems of finite support iterations*, J. Symb. Log. **83** (2018), no. 1, 208–236.
- [10] Fischer V., Mejía D. A., *Splitting, bounding, and almost disjointness can be quite different*, Canad. J. Math. **69** (2017), no. 3, 502–531.
- [11] Fischer V., Steprāns J., *The consistency of  $\mathfrak{b} = \kappa$  and  $\mathfrak{s} = \kappa^+$* , Fund. Math. **201** (2008), no. 3, 283–293.
- [12] Goldstern M., Kellner J., Mejía D. A., Shelah S., *Preservation of splitting families and cardinal characteristics of the continuum*, Israel J. Math. **246** (2021), no. 1, 73–129.
- [13] Ihoda J. I., Shelah S., *Souslin forcing*, J. Symbolic Logic **53** (1988), no. 4, 1188–1207.
- [14] Jech T., *Set Theory*, Springer Monographs in Mathematics, Springer, Berlin, 2003.
- [15] Kunen K., Vaughan J. E., eds., *Handbook of Set-theoretic Topology*, North-Holland Publishing Co., Amsterdam, 1984.
- [16] Mejía D. A., *Matrix iterations and Cichon’s diagram*, Arch. Math. Logic **52** (2013), no. 3–4, 261–278.
- [17] Shelah S., *On cardinal invariants of the continuum*, Conf. Axiomatic Set Theory, Boulder, 1983, Contemp. Math., 31, Amer. Math. Soc., Providence, 1984, pages 183–207.

A. Dow:

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE,  
9201 UNIVERSITY CITY BLVD, CHARLOTTE, NC 28223-0001, NORTH CAROLINA,  
U.S.A.

*E-mail:* adow@charlotte.edu

S. Shelah:

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY,  
HILL CENTER FOR THE MATHEMATICAL SCIENCES, 110 FRELINGHUYSEN RD.,  
PISCATAWAY, NJ 08854-8019, NEW JERSEY, U.S.A.

current address:

EINSTEIN INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, GIVAT RAM,  
JERUSALEM 9190401, ISRAEL

*E-mail:* shelah@math.rutgers.edu

(Received June 6, 2022, revised October 1, 2023)