

Zdeněk Dušek

## Explicit geodesic graphs on some H-type groups

In: Jan Slovák and Martin Čadek (eds.): Proceedings of the 21st Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 2002. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 69. pp. [77]--88.

Persistent URL: <http://dml.cz/dmlcz/701689>

### Terms of use:

© Circolo Matematico di Palermo, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## EXPLICIT GEODESIC GRAPHS ON SOME H-TYPE GROUPS

ZDENĚK DUŠEK

**ABSTRACT.** A *g.o. space* is a homogeneous Riemannian manifold  $(M = G/H, g)$  on which every geodesic is an orbit of a one-parameter subgroup of the group  $G$ . ( $G$  acts transitively on  $M$  as a group of isometries.) Each *g.o. space* gives rise to certain rational maps called "geodesic graphs". We are particularly interested in the case when the geodesic graphs are of non-linear character.

H-type groups provide the examples of these spaces. In this article we study H-type groups with 2-dimensional and 3-dimensional center and we present geodesic graphs with respect to various groups of isometries.

### 1. INTRODUCTION

Let  $(M, g)$  be a connected Riemannian manifold,  $p \in M$  a fixed point and let  $G$  be a connected group of isometries which acts transitively on  $M$ . Then  $M$  can be viewed as a homogeneous space  $(G/H, g)$ , where  $H$  is the isotropy subgroup at  $p$ . The Lie algebra of  $G$ , or  $H$ , respectively, will be denoted by  $\mathfrak{g}$ , or  $\mathfrak{h}$ , respectively.

**Definition 1.** A homogeneous space  $(G/H, g)$  is called a (*Riemannian*) *g.o. space*, if each geodesic of  $(G/H, g)$  (with respect to the Riemannian connection) is an orbit of a one-parameter subgroup  $\{\exp(tZ)\}$ ,  $Z \in \mathfrak{g}$ , of the group of isometries  $G$ .

**Definition 2.** Let  $(G/H, g)$  be a Riemannian *g.o. space*. A vector  $X \in \mathfrak{g} \setminus \{0\}$  is called a *geodesic vector* if the curve  $\exp(tX)(p)$  is a geodesic.

In a *g.o. space* we investigate those sets of geodesic vectors which generate all geodesics through a fixed point. These sets are called "geodesic graphs". Let us recall basic facts about geodesic graphs. (A comprehensive information can be found in [1].)

On the Lie algebra  $\mathfrak{g}$  of the group  $G$  there exists an  $\text{Ad}(H)$ -invariant decomposition (reductive decomposition)  $\mathfrak{g} = \mathfrak{m} + \mathfrak{h}$ , where  $\mathfrak{h}$  is the Lie algebra of the group  $H$  and  $\mathfrak{m}$  is a vector space  $\mathfrak{m} \subset \mathfrak{g}$ . (Such a decomposition is not unique.) On the vector space  $\mathfrak{m}$  there is a natural  $\text{Ad}(H)$ -invariant scalar product. It comes from the identification of  $\mathfrak{m} \subset T_e G$  with the tangent space  $T_p M$  via the projection  $\pi : G \mapsto M$ .

We define equivariant subalgebras  $\mathfrak{q}_X \subset \mathfrak{h}$  for  $X \in \mathfrak{m}$  in the following way

$$\mathfrak{q}_X = \{A \in \mathfrak{h} \mid [A, X] = 0\}$$

and we choose an invariant scalar product on  $\mathfrak{h}$ .

**Definition 3.** Let  $(G/H, g)$  be a g.o. space and  $\mathfrak{g} = \mathfrak{m} + \mathfrak{h}$  an  $\text{Ad}(H)$ -invariant decomposition of the Lie algebra  $\mathfrak{g}$ . The *canonical geodesic graph* is an  $\text{Ad}(H)$ -equivariant map  $\xi : \mathfrak{m} \rightarrow \mathfrak{h}$  (defined on an open dense subset of  $\mathfrak{m}$ ) such that  $X + \xi(X)$  is a geodesic vector and  $\xi(X) \perp \mathfrak{q}_X$  for each  $X \in \mathfrak{m} \setminus \{0\}$ .

For the existence of the canonical geodesic graph see [4], [2]. It is analytic on an open dense subset of  $\mathfrak{m}$ .

**Definition 4.** Let  $(G/H, g)$  be a g.o. space and  $\mathfrak{g} = \mathfrak{m} + \mathfrak{h}$  an  $\text{Ad}(H)$ -invariant decomposition of the Lie algebra  $\mathfrak{g}$ . A *general geodesic graph* is an  $\text{Ad}(H)$ -equivariant map  $\eta : \mathfrak{m} \rightarrow \mathfrak{h}$  which is analytic on an open dense subset of  $\mathfrak{m}$  and such that  $X + \eta(X)$  is a geodesic vector for each  $X \in \mathfrak{m} \setminus \{0\}$ .

**Remark.** The subalgebras  $\mathfrak{q}_X$  have the following property: If  $X \in \mathfrak{m}$ ,  $A \in \mathfrak{h}$  are the vectors such that  $X + A$  is a geodesic vector then all geodesic vectors "based on  $X$ " are of the form  $X + A + Q$ , where  $Q \in \mathfrak{q}_X$ . If the algebra  $\mathfrak{q}_X$  is nontrivial, this gives us the possibility to find more geodesic graphs than the canonical one. If the algebras  $\mathfrak{q}_X$  are trivial, then only canonical geodesic graph exists.

An essential tool for constructing geodesic graphs is the following

**Proposition 1** (cf. [2], Corollary 2.2). A vector  $Z \in \mathfrak{g} \setminus \{0\}$  is geodesic if and only if

$$(1) \quad \langle [Z, Y]_{\mathfrak{m}}, Z_{\mathfrak{m}} \rangle = 0 \quad \text{for all } Y \in \mathfrak{m}.$$

Here the subscript  $\mathfrak{m}$  indicates the projection into  $\mathfrak{m}$ .

We replace the vector  $Z$  by a vector  $X + \xi(X)$  expressed with respect to the bases  $\{X_i\}$  of  $\mathfrak{m}$  and  $\{D_j\}$  of  $\mathfrak{h}$  as

$$X = \sum_{i=1}^{\dim \mathfrak{m}} x_i X_i, \quad \xi(X) = \sum_{j=1}^{\dim \mathfrak{h}} \xi_j D_j$$

and for  $Y$  we substitute step by step all the elements  $X_i$ .

We obtain a system of linear equations for  $\xi_j$  with coefficients and right-hand sides depending on  $x_i$ . If this system doesn't have the unique solution ( $\dim \mathfrak{q}_X = q > 0$  for generic  $X \in \mathfrak{m}$ ) then we add  $q$  additional linear equations, which characterize the orthogonality  $\xi(X) \perp \mathfrak{q}_X$  (see [1] for detailed construction).

This extended system has the unique solution and by using the Cramer's rule we obtain a vector  $\xi(X)$ , whose components with respect to the basis of  $\mathfrak{h}$  are of the form  $\xi_j = P_j/P$ , where  $P_j$  and  $P$  are homogeneous polynomials in variables  $x_i$  and  $\deg(P_j) = \deg(P) + 1$ .

In the examples already known these polynomials have the common factor and the degree of the polynomials can be decreased. We define the *degree of a geodesic graph* as the degree of the denominator after cancelling the common factor out.

## 2. H-TYPE GROUPS

**Definition 5.** Let  $\mathfrak{n}$  be a 2-step nilpotent Lie algebra with an inner product  $\langle \cdot, \cdot \rangle$ . Let  $\mathfrak{z}$  be the center of  $\mathfrak{n}$  and let  $\mathfrak{v}$  be it's orthogonal complement. For each vector  $Z \in \mathfrak{z}$

define the operator  $J_Z : \mathfrak{v} \mapsto \mathfrak{v}$  by the relation

$$(2) \quad \langle J_Z X, Y \rangle = \langle Z, [X, Y] \rangle \quad \text{for all } X, Y \in \mathfrak{v}.$$

The algebra  $\mathfrak{n}$  is called a *generalized Heisenberg algebra (H-type algebra)* if, for each  $Z \in \mathfrak{z}$ , the operator  $J_Z$  satisfies the identity

$$(3) \quad J_Z^2 = -\langle Z, Z \rangle \text{id}_{\mathfrak{v}}.$$

A connected, simply connected Lie group whose Lie algebra is an H-type algebra is diffeomorphic to  $\mathbb{R}^n$  and it is called an *H-type group*. It is endowed with a left-invariant metric.

H-type algebras are completely classified (see [3]). For each dimension of the center  $\mathfrak{z}$  there is a series of H-type algebras. Each algebra of the series contains the center  $\mathfrak{z}$  and the complement  $\mathfrak{v}$  which decomposes into irreducible  $\mathfrak{z}$ -modules (the operators  $J_Z$  make  $\mathfrak{v}$  a  $\mathfrak{z}$ -module). Irreducible  $\mathfrak{z}$ -modules are all equivalent if  $\dim \mathfrak{z} \not\equiv 3 \pmod{4}$ , otherwise there exist two nonequivalent irreducible modules of the same dimension (called non-isotypic modules).

The H-type group is a g.o. space if and only if (see [7] or [3])

- $\dim \mathfrak{z} \in \{1, 2, 3\}$  or
- $\dim \mathfrak{z} \in \{5, 6, 7\}$  and  $\dim \mathfrak{v} = 8$  or
- $\dim \mathfrak{z} = 7$  and  $\dim \mathfrak{v} \in \{16, 24\}$  and  $\mathfrak{v}$  is decomposed into 8-dimensional modules of the same type.

Each H-type group with  $\dim \mathfrak{z} = 1$  is a naturally reductive space. The geodesic graph for naturally reductive spaces is linear - of degree 0. H-type groups with  $\dim \mathfrak{z} = 3$  are naturally reductive if and only if the complement  $\mathfrak{v}$  is decomposed into equivalent modules. Other H-type groups which are g.o. spaces are not naturally reductive. In the following sections we will concentrate on H-type groups with  $\dim \mathfrak{z} = 2$  or 3. The case  $\dim \mathfrak{z} = 5$  is investigated in [5].

### 2.1 $\dim \mathfrak{z} = 2$

Let  $\mathfrak{n}$  be a vector space of dimension  $4n + 2$  equipped with a scalar product and let  $\{E_1, \dots, E_{4n}, Z_1, Z_2\}$  form an orthonormal basis. We define the structure of a Lie algebra on  $\mathfrak{n}$  by the following relations. For  $p = 0, \dots, n - 1$

$$\begin{aligned} [E_{4p+1}, E_{4p+2}] &= 0, \\ [E_{4p+1}, E_{4p+3}] &= Z_1, \quad [E_{4p+2}, E_{4p+3}] = Z_2, \\ [E_{4p+1}, E_{4p+4}] &= Z_2, \quad [E_{4p+2}, E_{4p+4}] = -Z_1, \quad [E_{4p+3}, E_{4p+4}] = 0, \end{aligned}$$

for other  $k, l = 1, \dots, 4n$  we put  $[E_k, E_l] = 0$ , further  $[Z_1, Z_2] = 0$ , and for  $k = 1, \dots, 4n$  and  $l = 1, 2$  we put  $[E_k, Z_l] = 0$ .

The elements  $Z_1$  and  $Z_2$  span the center  $\mathfrak{z}$  of the Lie algebra  $\mathfrak{n}$  and one easily verifies the condition (3) for the operators  $J_Z$ , so this relations define an H-type algebra. Each quadruplet  $\mathfrak{v}_p = \text{span}(E_{4p+1}, \dots, E_{4p+4})$  for  $0 \leq p \leq n - 1$  is an irreducible  $\mathfrak{z}$ -module

and these modules are equivalent to each other. Summarizing, we have

$$\mathfrak{n} = \mathfrak{z} + \mathfrak{v} = \mathfrak{z} + \sum_{p=0}^{n-1} \mathfrak{v}_p.$$

If  $n = 1$  then we have the Lie algebra of the simplest (6-dimensional) H-type group with 2-dimensional center. It was the first example (by A. Kaplan) of a g.o. space which is not naturally reductive. Its geodesic graph was described in [2] and this section is a generalization to other H-type groups with  $\dim \mathfrak{z} = 2$ .

Let us express the H-type group  $N$  corresponding to  $\mathfrak{n}$  as a homogeneous space  $G/H$ . For  $p = 0, \dots, n-1$ , the following operators acting on  $\mathfrak{v}$  are skew-symmetric derivations of the Lie algebra  $\mathfrak{n}$ :

$$D_{3p+1} = -A_{(4p+1, 4p+2)} + A_{(4p+3, 4p+4)},$$

$$D_{3p+2} = +A_{(4p+1, 4p+3)} + A_{(4p+2, 4p+4)},$$

$$D_{3p+3} = +A_{(4p+1, 4p+4)} - A_{(4p+2, 4p+3)}.$$

Here  $A_{(k,l)}$  are the elements of  $\mathfrak{so}(\mathfrak{v})$  acting on  $\mathfrak{v}$  by  $A_{(k,l)}(E_i) = \delta_{ki}E_l - \delta_{li}E_k$ . So each subalgebra  $\mathfrak{h}_p = \text{span}(D_{3p+1}, \dots, D_{3p+3})$  acts effectively only on  $\mathfrak{v}_p$ .

We put

$$\mathfrak{h} = \text{span}(D_1, \dots, D_{3n}) = \bigoplus_{p=0}^{n-1} \mathfrak{h}_p \cong \bigoplus_{p=0}^{n-1} \mathfrak{su}(2).$$

and consider the decomposition  $\mathfrak{g} = \mathfrak{n} + \mathfrak{h}$ . Obviously,  $\mathfrak{g}$  is a well-defined Lie algebra. If we express the H-type group  $N$  corresponding to  $\mathfrak{n}$  as a homogeneous space  $G/H$  then  $G$  can be considered as a transitive group of *isometries* of  $N$ .

Hence we have  $N = G/H$ , where  $H \cong [\text{SU}(2)]^n$  and  $G = N \rtimes H$ . Here the group  $G$  is not the full isometry group of  $N$ . But the group  $N$  is a g.o. space with respect to this group.

Now, we shall construct the canonical geodesic graph  $\xi : \mathfrak{n} \mapsto \mathfrak{h}$ . We put

$$X = \sum_{k=1}^{4n} x_k E_k + \sum_{l=1}^2 z_l Z_l, \quad \xi(X) = \sum_{i=1}^{3n} \xi_i D_i.$$

We check easily that the subalgebras  $\mathfrak{q}_X$  from the Introduction are trivial. From the equation (1) we obtain  $4n + 2$  linear equations for the components  $\xi_i$  ( $i = 1, \dots, 3n$ ) of the vector  $\xi(X)$  depending on the variables  $x_k$  and  $z_l$  ( $k = 1, \dots, 4n$  and  $l = 1, 2$ ).

For each quadruplet of these equations corresponding to  $Y = E_{4p+1}, \dots, E_{4p+4}$  only three of them are linearly independent. Hence we omit the fourth equation from each quadruplet (corresponding to  $Y = E_{4p+4}$  for  $p = 0, \dots, n-1$ ). The last two equations (corresponding to  $Y = Z_1$  and  $Y = Z_2$ ) are trivial.

The matrix of this system of equations is equivalent to the block square matrix  $A$  (of rank  $3n$ ) with nonzero  $3 \times 3$  blocks just along the diagonal. These blocks are

$$A_p = \begin{pmatrix} -x_{4p+2} & x_{4p+3} & x_{4p+4} \\ x_{4p+1} & x_{4p+4} & -x_{4p+3} \\ x_{4p+4} & -x_{4p+1} & x_{4p+2} \end{pmatrix} \quad \text{for } p = 0, \dots, n-1.$$

The right-hand side vector  $\mathbf{b}$  (of  $3n$  entries) can be written in block form as  $\mathbf{b} = (\mathbf{b}_0, \dots, \mathbf{b}_{n-1})^t$ , where

$$\mathbf{b}_p = \begin{pmatrix} x_{4p+3}z_1 + x_{4p+4}z_2 \\ -x_{4p+4}z_1 + x_{4p+3}z_2 \\ -x_{4p+1}z_1 - x_{4p+2}z_2 \end{pmatrix} \quad \text{for } p = 0, \dots, n-1.$$

Hence we solve the matrix equation  $A\xi = \mathbf{b}$ , where  $\xi = (\xi_1, \dots, \xi_{3n})^t$ . Using the Cramer's rule we get explicitly

$$\begin{aligned} \xi_{3p+1} &= \frac{-2(x_{4p+2}x_{4p+3} + x_{4p+4}x_{4p+1})z_1 - 2(x_{4p+2}x_{4p+4} - x_{4p+1}x_{4p+3})z_2}{x_{4p+1}^2 + x_{4p+2}^2 + x_{4p+3}^2 + x_{4p+4}^2}, \\ \xi_{3p+2} &= \frac{(x_{4p+1}^2 - x_{4p+2}^2 + x_{4p+3}^2 - x_{4p+4}^2)z_1 + 2(x_{4p+3}x_{4p+4} + x_{4p+1}x_{4p+2})z_2}{x_{4p+1}^2 + x_{4p+2}^2 + x_{4p+3}^2 + x_{4p+4}^2}, \\ \xi_{3p+3} &= \frac{2(x_{4p+3}x_{4p+4} - x_{4p+1}x_{4p+2})z_1 + (x_{4p+1}^2 - x_{4p+2}^2 - x_{4p+3}^2 + x_{4p+4}^2)z_2}{x_{4p+1}^2 + x_{4p+2}^2 + x_{4p+3}^2 + x_{4p+4}^2}, \\ &\quad \text{for } 0 \leq p \leq n-1. \end{aligned}$$

Thus, there is a canonical geodesic graph of degree 2 for every  $H$ -type group with  $\dim \mathfrak{g} = 2$ . Our choice of the group  $G$  doesn't involve other geodesic graphs, because we have  $\dim \mathfrak{q}_X = 0$  for generic  $X$ .

Now, let us express the group  $N$  in the new form  $G'/H'$ , where  $G'$  is the full isometry group and look for geodesic graphs with respect to bigger groups of isometries. The 6-dimensional  $H$ -type group was treated in [1]. Here the full isometry group  $G'$  is one dimension bigger than  $G$ . In the decomposition  $\mathfrak{g}' = \mathfrak{n} + \mathfrak{h}'$  we have  $\mathfrak{h}' = \text{span}(\mathfrak{h}, R)$ .  $R$  is the operator

$$R = 2B_{(1,2)} + A_{(1,2)} + A_{(3,4)}.$$

(Here  $B_{(1,2)}$  is the operator on  $\mathfrak{g}$  acting by  $B_{(1,2)}(Z_i) = \delta_{1i}Z_2 - \delta_{2i}Z_1$ .) But the equation (1) implies that the component of the operator  $R$  in any geodesic graph is zero. In this case only canonical geodesic graph exists.

In the 10-dimensional case ( $\mathfrak{n} = \mathfrak{g} + \sum_{p=0}^1 \mathfrak{v}_p$ ) the algebra  $\mathfrak{h}'$  in the decomposition  $\mathfrak{g}' = \mathfrak{n} + \mathfrak{h}'$  is spanned by 11 skew-symmetric derivations on  $\mathfrak{n}$ . We denote them  $D_1, \dots, D_6, P_1, \dots, P_5$ . The new elements act on  $\mathfrak{n}$  by

$$\begin{aligned} P_1 &= +A_{(1,5)} + A_{(2,6)} + A_{(3,7)} + A_{(4,8)}, \\ P_2 &= +A_{(1,6)} - A_{(2,5)} - A_{(3,8)} + A_{(4,7)}, \\ P_3 &= +A_{(1,7)} + A_{(2,8)} - A_{(3,5)} - A_{(4,6)}, \\ P_4 &= +A_{(1,8)} - A_{(2,7)} + A_{(3,6)} - A_{(4,5)}, \end{aligned}$$

$$P_5 = 2B_{(1,2)} + A_{(1,2)} + A_{(3,4)} + A_{(5,6)} + A_{(7,8)}.$$

Again, the equation (1) implies that the component of the operator  $P_5$  in any geodesic graph is zero. We denote

$$\mathfrak{h}'' = \text{span}(D_1, \dots, D_6, P_1, \dots, P_4) \cong \mathfrak{so}(5).$$

We are given the new expression for the group  $N$  as  $G''/H''$ , where  $H'' = \text{Spin}(5)$  and  $G'' = N \rtimes H''$ . In this case we have  $\dim_{\mathbb{R}} \mathfrak{n} = 3$  for generic  $X \in \mathfrak{n}$ . Hence general geodesic graphs do exist. Conjecture: there is no geodesic graph of degree 1.

## 2.2. $\dim \mathfrak{z} = 3$ .

In this case we have a vector space  $\mathfrak{n}$  of dimension  $4(n+m)+3$  equipped with a scalar product and the elements  $\{E_1, \dots, E_{4n}, F_1, \dots, F_{4m}, Z_1, \dots, Z_3\}$  form an orthonormal basis. The structure of a Lie algebra on  $\mathfrak{n}$  is defined by the following relations. For  $p = 0, \dots, n-1$

$$\begin{aligned} [E_{4p+1}, E_{4p+2}] &= Z_1, \\ [E_{4p+1}, E_{4p+3}] &= Z_2, \quad [E_{4p+2}, E_{4p+3}] = Z_3, \\ [E_{4p+1}, E_{4p+4}] &= Z_3, \quad [E_{4p+2}, E_{4p+4}] = -Z_2, \quad [E_{4p+3}, E_{4p+4}] = Z_1, \end{aligned}$$

for  $q = 0, \dots, m-1$

$$\begin{aligned} [F_{4q+1}, F_{4q+2}] &= Z_1, \\ [F_{4q+1}, F_{4q+3}] &= Z_2, \quad [F_{4q+2}, F_{4q+3}] = -Z_3, \\ [F_{4q+1}, F_{4q+4}] &= Z_3, \quad [F_{4q+2}, F_{4q+4}] = Z_2, \quad [F_{4q+3}, F_{4q+4}] = -Z_1. \end{aligned}$$

For other  $i, j = 1, \dots, 4n$  and  $k, l = 1, \dots, 4m$  we put  $[E_i, E_j] = 0$ ,  $[F_k, F_l] = 0$ , and for  $i = 1, \dots, 4n$ ,  $j = 1, \dots, 4m$  and  $k, l = 1, \dots, 3$  we put

$$[E_i, Z_k] = 0, \quad [F_j, Z_k] = 0, \quad [E_i, F_j] = 0, \quad [Z_k, Z_l] = 0.$$

We have  $\mathfrak{z} = \text{span}(Z_1, \dots, Z_3)$ ,  $\mathfrak{v}_p = \text{span}(E_{p+1}, \dots, E_{p+4})$  for  $0 \leq p \leq n-1$  and  $\bar{\mathfrak{v}}_q = \text{span}(F_{q+1}, \dots, F_{q+4})$  for  $0 \leq q \leq m-1$ . The action of  $\mathfrak{z}$  on  $\mathfrak{v}_p$  (via the operators  $J_Z$ ) can be viewed as multiplication of quaternions by imaginary quaternions on the left and the action on  $\bar{\mathfrak{v}}_q$  as multiplication on the right. The modules  $\mathfrak{v}_p$  and  $\bar{\mathfrak{v}}_q$  are not equivalent.

We start with the simplest case  $n = 1, m = 0$ . It is the seven-dimensional algebra  $\mathfrak{n}_{(1,0)} = \mathfrak{z} + \mathfrak{v}$  with  $\mathfrak{v} = \mathfrak{v}_0$ . (The double index at  $\mathfrak{n}$  shows the number of modules of each type in the complement of  $\mathfrak{z}$ .) We will show geodesic graphs with respect to various groups of isometries and apply the results to the general case.

To express  $N_{(1,0)} = G/H$  we put  $\mathfrak{h} = \text{Der}(\mathfrak{n}) \cap \mathfrak{so}(\mathfrak{n})$  in the decomposition  $\mathfrak{g} = \mathfrak{n} + \mathfrak{h}$ . We get the following operators on  $\mathfrak{n}$

$$\begin{aligned} D_1 &= -A_{(1,2)} + A_{(3,4)}, \quad D_4 = 2B_{(2,3)} + A_{(1,2)} + A_{(3,4)}, \\ D_2 &= +A_{(1,3)} + A_{(2,4)}, \quad D_5 = 2B_{(1,3)} - A_{(1,3)} + A_{(2,4)}, \\ D_3 &= +A_{(1,4)} - A_{(2,3)}, \quad D_6 = 2B_{(1,2)} + A_{(1,4)} + A_{(2,3)}. \end{aligned}$$

Again,  $A_{(k,l)}$  are the elements of  $\mathfrak{so}(\mathfrak{v})$  acting on  $\mathfrak{v}$  by  $A_{(k,l)}(E_i) = \delta_{ki}E_l - \delta_{li}E_k$  and  $B_{(k,l)}$  are the elements of  $\mathfrak{so}(\mathfrak{z})$  acting on  $\mathfrak{z}$  by  $B_{(k,l)}(Z_i) = \delta_{ki}Z_l - \delta_{li}Z_k$ .

We have

$$\mathfrak{h} = \text{span}(D_1, \dots, D_6) \cong \mathfrak{su}(2) \oplus \tilde{\mathfrak{su}}(2),$$

where  $\tilde{\mathfrak{su}}(2)$  means another representation of  $\mathfrak{su}(2)$  on  $\mathfrak{n}$ . The group  $G$  corresponding to the algebra  $\mathfrak{g}$  is the maximal connected isometry group of  $N$ .

The system of equations obtained from the equation (1) in the same way as in 2.1. is equivalent to the matrix equation  $\mathbf{A}\xi = \mathbf{b}$  with

$$\mathbf{A} = \begin{pmatrix} -x_2 & x_3 & x_4 & x_2 & -x_3 & x_4 \\ x_1 & x_4 & -x_3 & -x_1 & x_4 & x_3 \\ x_4 & -x_1 & x_2 & x_4 & x_1 & -x_2 \\ 0 & 0 & 0 & 0 & 2z_3 & 2z_2 \\ 0 & 0 & 0 & 2z_3 & 0 & -2z_1 \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} x_2z_1 + x_3z_2 + x_4z_3 \\ -x_1z_1 - x_4z_2 + x_3z_3 \\ x_4z_1 - x_1z_2 - x_2z_3 \\ 0 \\ 0 \end{pmatrix}.$$

The solution of this system is not unique ( $\dim \mathfrak{q}_X = 1$ ). We find the generator of the algebra  $\mathfrak{q}_X$  as the solution of the homogeneous system  $\mathbf{A} \cdot \mathbf{Q}(X) = 0$  (see [1]). The components  $(Q_i)_{i=1}^6$  of the vector  $\mathbf{Q}(X)$  may be chosen as corresponding maximal subdeterminants with the corresponding signs of the matrix  $\mathbf{A}$ . But all these determinants have the common factor  $4x_4z_3$  and therefore we can cancel out by this common factor and get the simpler components

$$\begin{aligned} Q_1 &= (-x_1^2 - x_2^2 + x_3^2 + x_4^2)z_1 - 2(x_3x_2 + x_1x_4)z_2 + 2(x_1x_3 - x_4x_2)z_3, \\ Q_2 &= 2(x_3x_2 - x_1x_4)z_1 + (x_1^2 - x_2^2 + x_3^2 - x_4^2)z_2 + 2(x_1x_2 + x_4x_3)z_3, \\ Q_3 &= 2(x_2x_4 + x_1x_3)z_1 + 2(x_4x_3 - x_1x_2)z_2 + (x_1^2 - x_2^2 - x_3^2 + x_4^2)z_3, \\ Q_4 &= -(x_2^2 + x_1^2 + x_3^2 + x_4^2)z_1, \\ Q_5 &= (x_2^2 + x_1^2 + x_3^2 + x_4^2)z_2, \\ Q_6 &= -(x_2^2 + x_1^2 + x_3^2 + x_4^2)z_3. \end{aligned}$$

We extend the matrix  $\mathbf{A}$  by the row vector  $\mathbf{Q}(X)^t$  and the vector  $\mathbf{b}$  by the sixth component equal to 0. So we have added the condition  $\mathbf{Q}(X) \perp \xi(X)$  (the invariant scalar product on  $\mathfrak{h}$  is chosen so that  $\{D_i\}_{i=1}^6$  form an orthonormal basis). The solution of the extended matrix equation (obtained by the Cramer's rule) is

$$\begin{aligned} \xi_1 &= \frac{(-x_1^2 - x_2^2 + x_3^2 + x_4^2)z_1 - 2(x_1x_4 + x_3x_2)z_2 + 2(x_1x_3 - x_2x_4)z_3}{2(x_1^2 + x_2^2 + x_3^2 + x_4^2)}, \\ \xi_2 &= \frac{2(x_2x_3 - x_1x_4)z_1 + (x_1^2 - x_2^2 + x_3^2 - x_4^2)z_2 + 2(x_1x_2 + x_4x_3)z_3}{2(x_1^2 + x_2^2 + x_3^2 + x_4^2)}, \\ \xi_3 &= \frac{2(x_2x_4 + x_1x_3)z_1 + 2(x_3x_4 - x_1x_2)z_2 + (x_1^2 - x_2^2 - x_3^2 + x_4^2)z_3}{2(x_1^2 + x_2^2 + x_3^2 + x_4^2)}, \\ \xi_4 &= 1/2 z_1, \\ \xi_5 &= -1/2 z_2, \\ \xi_6 &= 1/2 z_3. \end{aligned}$$



Hence the degree of the canonical geodesic graph in the full isometry group is equal to 2. But we may consider the map

$$\eta(X) = \xi(X) - \frac{1}{2(x_1^2 + x_2^2 + x_3^2 + x_4^2)} Q(X).$$

We have  $\text{Ad}(h)\mathfrak{q}_X = \mathfrak{q}_{\text{Ad}(h)X}$  and for the scalar product on  $\mathfrak{h}$  such that  $\{D_i\}_{i=1}^6$  form an orthonormal basis we have

$$\|Q(X), Q(X)\|^2 = 2(z_1^2 + z_2^2 + z_3^2)(x_2^2 + x_1^2 + x_3^2 + x_4^2)^2,$$

which is an invariant function with respect to the representation  $\text{Ad}(H)|_{\mathfrak{n}}$ . The function  $\frac{1}{2(x_1^2 + x_2^2 + x_3^2 + x_4^2)}$  is invariant as well, so the map  $\eta$  is  $\text{Ad}(H)$ -equivariant. It is a general geodesic graph and it is obvious, that this geodesic graph is linear. Indeed

$$\eta_1 = \eta_2 = \eta_3 = 0, \quad \eta_4 = z_1, \quad \eta_5 = -z_2, \quad \eta_6 = z_3.$$

The linear geodesic graph shows, that the space  $N_{(1,0)}$  is naturally reductive. An interesting observation shows, that this linear geodesic graph can be obtained as the canonical geodesic graph in the smaller group of isometries. If we put  $N_{(1,0)} = G'/H'$ , where

$$\mathfrak{h}' = \text{span}(D_4, \dots, D_6) = \tilde{\mathfrak{su}}(2),$$

we get the same map. We will use this idea in constructing linear geodesic graphs in a more general case  $N_{(n,0)}$  or  $N_{(0,m)}$ .

Let  $\mathfrak{n}_{(n,0)} = \mathfrak{z} + \mathfrak{v}$  where  $\mathfrak{v} = \sum_{p=0}^{n-1} \mathfrak{v}_p$  be an  $H$ -type algebra with  $\dim \mathfrak{z} = 3$  and the complement  $\mathfrak{v}$  decomposed into  $n$  irreducible modules of the first type. We put  $N_{(n,0)} = G/H$  where  $\mathfrak{h} = \text{span}(\tilde{D}_1, \dots, \tilde{D}_3)$ , acting by

$$\tilde{D}_1 = 2B_{(2,3)} + \sum_{p=0}^{n-1} (A_{(4p+1,4p+2)} + A_{(4p+3,4p+4)}),$$

$$\tilde{D}_2 = 2B_{(1,3)} + \sum_{p=0}^{n-1} (-A_{(4p+1,4p+3)} + A_{(4p+2,4p+4)}),$$

$$\tilde{D}_3 = 2B_{(1,2)} + \sum_{p=0}^{n-1} (A_{(4p+1,4p+4)} + A_{(4p+2,4p+3)}).$$

We have  $H \cong \text{SU}(2)$ ,  $G = N \rtimes H$ . From (1) we get  $4n + 3$  linear equations with right-hand sides for 3 components  $(\xi_i)_{i=1}^3$ . This system splits into  $n$  quadruplets of equations corresponding to different values of  $p$  and one triplet of equations. In each quadruplet (for  $0 \leq p \leq n-1$ ) only three equations are linearly independent and they can be expressed as a matrix equation  $\tilde{\mathbf{A}}_p \xi = \tilde{\mathbf{b}}_p$  where

$$\tilde{\mathbf{A}}_p = \begin{pmatrix} x_{4p+2} & -x_{4p+3} & x_{4p+4} \\ -x_{4p+1} & x_{4p+4} & x_{4p+3} \\ x_{4p+4} & x_{4p+1} & -x_{4p+2} \end{pmatrix}, \quad \tilde{\mathbf{b}}_p = \begin{pmatrix} x_{4p+2}z_1 + x_{4p+3}z_2 + x_{4p+4}z_3 \\ -x_{4p+1}z_1 - x_{4p+4}z_2 + x_{4p+3}z_3 \\ x_{4p+4}z_1 - x_{4p+1}z_2 - x_{4p+2}z_3 \end{pmatrix}.$$

The components of the solution of each of these subsystems are

$$\xi_1 = z_1, \quad \xi_2 = -z_2, \quad \xi_3 = z_3$$

and one easily verifies, that it is the solution of the whole system.

Similarly, in the case of an H-type algebra with  $\dim \mathfrak{g} = 3$  and the complement  $\bar{\mathfrak{v}}$  decomposed into  $m$  irreducible modules of the second type ( $\mathfrak{n}_{(0,m)} = \mathfrak{g} + \bar{\mathfrak{v}}$  where  $\bar{\mathfrak{v}} = \sum_{q=0}^{m-1} \bar{\mathfrak{v}}_q$ ) we take  $\mathfrak{h} = \text{span}(\widehat{D}_1, \dots, \widehat{D}_3)$ , acting by

$$\begin{aligned} \widehat{D}_1 &= 2B_{(2,3)} + \sum_{q=0}^{m-1} (-\bar{A}_{(4q+1,4q+2)} + \bar{A}_{(4q+3,4q+4)}), \\ \widehat{D}_2 &= 2B_{(1,3)} + \sum_{q=0}^{m-1} (\bar{A}_{(4q+1,4q+3)} + \bar{A}_{(4q+2,4q+4)}), \\ \widehat{D}_3 &= 2B_{(1,2)} + \sum_{q=0}^{m-1} (-\bar{A}_{(4q+1,4q+4)} + \bar{A}_{(4q+2,4q+3)}). \end{aligned}$$

(Here  $\bar{A}_{(k,l)}$  acts on  $\bar{\mathfrak{v}}$  in the same way as  $A_{(k,l)}$  acts on  $\mathfrak{v}$ , namely  $\bar{A}_{(k,l)}(F_i) = \delta_{ki}F_l - \delta_{li}F_k$ .) We have  $N_{(0,m)} = G/H$  where  $H \cong \text{SU}(2)$ ,  $G = N \rtimes H$ . We put

$$X = \sum_{k=1}^{4m} y_k F_k + \sum_{l=1}^3 z_l Z_l, \quad \xi(X) = \sum_{i=1}^{3m} \xi_i \widehat{D}_i$$

and the equation (1) gives  $4m + 3$  linear equations. For example the first quadruplet reduces to the matrix equation  $\widehat{A}_0 \xi = \widehat{\mathfrak{b}}_0$  for

$$\widehat{A}_0 = \begin{pmatrix} -y_2 & y_3 & -y_4 \\ y_1 & y_4 & y_3 \\ y_4 & -y_1 & -y_2 \end{pmatrix}, \quad \widehat{\mathfrak{b}}_0 = \begin{pmatrix} y_2 z_1 + y_3 z_2 + y_4 z_3 \\ -y_1 z_1 + y_4 z_2 - y_3 z_3 \\ -y_4 z_1 - y_1 z_2 + y_2 z_3 \end{pmatrix}.$$

The solution, which solves other equations too, is again linear:

$$\xi_1 = -z_1, \quad \xi_2 = z_2, \quad \xi_3 = -z_3.$$

We see, that all the H-type groups mentioned in this section so far are naturally reductive spaces.

Finally, we shall consider the general case of an H-type algebra with  $\dim \mathfrak{g} = 3$ . We have  $\mathfrak{n}_{(n,m)} = \mathfrak{g} + \mathfrak{v} + \bar{\mathfrak{v}}$  where  $\mathfrak{v} = \sum_{p=0}^{n-1} \mathfrak{v}_p$  and  $\bar{\mathfrak{v}} = \sum_{q=0}^{m-1} \bar{\mathfrak{v}}_q$ . Now, we put

$$\mathfrak{h} = \text{span}(D_1, \dots, D_{3n}, \bar{D}_1, \dots, \bar{D}_{3m}) \simeq \bigoplus_{p=0}^{n-1} \mathfrak{su}(2) \oplus \bigoplus_{q=0}^{m-1} \mathfrak{su}(2).$$

Each copy of  $\mathfrak{su}(2)$  acts effectively only on unique  $\mathfrak{v}_p$  and each copy of  $\mathfrak{su}(2)$  acts effectively on unique  $\bar{\mathfrak{v}}_q$  by the following skew-symmetric derivations. For  $p = 0, \dots, n-1$

$$\begin{aligned} D_{3p+1} &= -A_{(4p+1,4p+2)} + A_{(4p+3,4p+4)}, \\ D_{3p+2} &= +A_{(4p+1,4p+3)} + A_{(4p+2,4p+4)}, \\ D_{3p+3} &= +A_{(4p+1,4p+4)} - A_{(4p+2,4p+3)} \end{aligned}$$

and for  $q = 0, \dots, m - 1$

$$\begin{aligned}\bar{D}_{3q+1} &= +\bar{A}_{(4q+1,4q+2)} + \bar{A}_{(4q+3,4q+4)}, \\ \bar{D}_{3q+2} &= +\bar{A}_{(4q+1,4q+3)} - \bar{A}_{(4q+2,4q+4)}, \\ \bar{D}_{3q+3} &= +\bar{A}_{(4q+1,4q+4)} + \bar{A}_{(4q+2,4q+3)}.\end{aligned}$$

There are other skew-symmetric derivations of  $\mathfrak{n}_{(n,m)}$  involving the operators  $B_{(k,l)}$  but they are not needed here.

Now we put

$$X = \sum_{i=1}^{4n} x_i E_i + \sum_{j=1}^{4m} y_j F_j + \sum_{k=1}^3 z_k Z_k, \quad \xi(X) = \sum_{i=1}^{3n} \xi_i D_i + \sum_{j=1}^{3m} \bar{\xi}_j \bar{D}_j.$$

The equation (1) gives again the system of linear equations. It is equivalent to the matrix equation  $\mathbf{A}\xi = \mathbf{b}$  for the block square matrix  $\mathbf{A}$  (of rank  $3(n+m)$ ) with nonzero  $3 \times 3$  blocks along the diagonal. These blocks are

$$\mathbf{A}_p = \begin{pmatrix} -x_{4p+2} & x_{4p+3} & x_{4p+4} \\ x_{4p+1} & x_{4p+4} & -x_{4p+3} \\ x_{4p+4} & -x_{4p+1} & x_{4p+2} \end{pmatrix}, \quad \bar{\mathbf{A}}_q = \begin{pmatrix} y_{4q+2} & y_{4q+3} & y_{4q+4} \\ -y_{4q+1} & -y_{4q+4} & y_{4q+3} \\ y_{4q+4} & -y_{4q+1} & -y_{4q+2} \end{pmatrix}$$

for  $p = 0, \dots, n - 1$  and  $q = 0, \dots, m - 1$ .

The right-hand side vector  $\mathbf{b}$  (of  $3(n+m)$  entries) can be written in block form as  $\mathbf{b} = (\mathbf{b}_0, \dots, \mathbf{b}_{n-1}, \bar{\mathbf{b}}_0, \dots, \bar{\mathbf{b}}_{m-1})^t$ , where

$$\mathbf{b}_p = \begin{pmatrix} x_{4p+2}z_1 + x_{4p+3}z_2 + x_{4p+4}z_3 \\ -x_{4p+1}z_1 - x_{4p+4}z_2 + x_{4p+3}z_3 \\ x_{4p+4}z_1 - x_{4p+1}z_2 - x_{4p+2}z_3 \end{pmatrix} \quad \text{for } p = 0, \dots, n - 1,$$

$$\bar{\mathbf{b}}_q = \begin{pmatrix} y_{4q+2}z_1 + y_{4q+3}z_2 + y_{4q+4}z_3 \\ -y_{4q+1}z_1 + y_{4q+4}z_2 - y_{4q+3}z_3 \\ -y_{4q+4}z_1 - y_{4q+1}z_2 + y_{4q+2}z_3 \end{pmatrix} \quad \text{for } q = 0, \dots, m - 1.$$

By using the Cramer's rule we get (after cancelling the common factors out) the components of the canonical geodesic graph of degree 2. The components  $\xi_i$  and  $\bar{\xi}_j$

for  $1 \leq i, j \leq 3$  are

$$\begin{aligned} \xi_1 &= \frac{(-x_1^2 - x_2^2 + x_3^2 + x_4^2) z_1 - 2(x_3 x_2 + x_1 x_4) z_2 + 2(x_1 x_3 - x_2 x_4) z_3}{x_1^2 + x_2^2 + x_3^2 + x_4^2}, \\ \xi_2 &= \frac{2(x_2 x_3 - x_1 x_4) z_1 + (x_1^2 - x_2^2 + x_3^2 - x_4^2) z_2 + 2(x_1 x_2 + x_3 x_4) z_3}{x_1^2 + x_2^2 + x_3^2 + x_4^2}, \\ \xi_3 &= \frac{2(x_3 x_1 + x_4 x_2) z_1 + 2(x_3 x_4 - x_1 x_2) z_2 + (x_1^2 - x_2^2 - x_3^2 + x_4^2) z_3}{x_1^2 + x_2^2 + x_3^2 + x_4^2}, \\ \bar{\xi}_1 &= \frac{(y_1^2 + y_2^2 - y_3^2 - y_4^2) z_1 + 2(y_2 y_3 - y_1 y_4) z_2 + 2(y_3 y_1 + y_4 y_2) z_3}{y_1^2 + y_2^2 + y_3^2 + y_4^2}, \\ \bar{\xi}_2 &= \frac{2(y_3 y_2 + y_4 y_1) z_1 + (y_1^2 - y_2^2 + y_3^2 - y_4^2) z_2 + 2(y_3 y_4 - y_1 y_2) z_3}{y_1^2 + y_2^2 + y_3^2 + y_4^2}, \\ \bar{\xi}_3 &= \frac{2(y_2 y_4 - y_1 y_3) z_1 + 2(y_1 y_2 + y_4 y_3) z_2 + (y_1^2 - y_2^2 - y_3^2 + y_4^2) z_3}{y_1^2 + y_2^2 + y_3^2 + y_4^2} \end{aligned}$$

and the components  $\xi_{3p+i}$  and  $\bar{\xi}_{3q+j}$  for  $1 \leq p \leq n-1$  and  $1 \leq q \leq m-1$  are obtained after replacing all  $x_k$  by the corresponding  $x_{4p+k}$  and all  $y_l$  by the corresponding  $y_{4q+l}$  ( $k, l = 1, \dots, 4$ ).

In the general case  $N_{(n,m)}$  we can't use the similar construction as in the case  $N_{(1,0)}$  and construct linear geodesic graph. For example in  $N_{(1,1)}$  we have  $N_{(1,1)} = G/H$  where  $\mathfrak{h} = \text{span}(D_1, \dots, D_3, \bar{D}_1, \dots, \bar{D}_3)$ . The group  $G$  may be enlarged to the full connected isometry group  $\tilde{G}$  of  $N$  and we get  $N_{(1,1)} = \tilde{G}/\tilde{H}$  with  $\tilde{\mathfrak{h}} = \mathfrak{h} + \text{span}(\bar{D}_1, \dots, \bar{D}_3)$ . The action of additional elements of  $\tilde{\mathfrak{h}}$  on  $\mathfrak{n}$  is given by

$$\begin{aligned} \tilde{D}_1 &= 2B_{(2,3)} + A_{(1,2)} + A_{(3,4)} - \bar{A}_{(1,2)} + \bar{A}_{(3,4)}, \\ \tilde{D}_2 &= 2B_{(1,3)} - A_{(1,3)} + A_{(2,4)} + \bar{A}_{(1,3)} + \bar{A}_{(2,4)}, \\ \tilde{D}_3 &= 2B_{(1,2)} + A_{(1,4)} + A_{(2,3)} - \bar{A}_{(1,4)} + \bar{A}_{(2,3)}. \end{aligned}$$

If we compute the canonical geodesic graph with respect to  $\tilde{G}$  (here  $\dim \mathfrak{q}_X = 1$ ) we get the same map as with respect to  $G$ , the components of  $\{\tilde{D}_i\}_{i=1}^3$  are zero. It is not hard to show that the similar trick for decreasing the degree as in the case  $N_{(1,0)}$  doesn't work. It corresponds to the fact, which is known from the general theory, namely that the  $H$ -type groups with  $\dim \mathfrak{g} = 3$  and of general type are not naturally reductive.

**Acknowledgments.** This work was supported by the grant GAČR 201/99/0265. I wish to thank to my advisor Oldřich Kowalski for the aid with the final version of the paper.

REFERENCES

[1] Kowalski, O. and Nikčević, S. Ž., *On geodesic graphs of Riemannian g.o. spaces*, Arch. Math. 73 (1999), 223–234.  
 [2] Kowalski, O. and Vanhecke, L., *Riemannian manifolds with homogeneous geodesics*, Boll. Un. Mat. Ital. B(7)5 (1991), 189–246.

- [3] Berndt, J., Tricerri, F., Vanhecke, L., *Generalized Heisenberg groups and Damek-Ricci harmonic spaces*, Springer-Verlag 1995.
- [4] Szenthe, J., *Sur la connection naturelle à torsion nulle*, Acta Sci. Math. (Szeged), **38** (1976), 383–398.
- [5] Dušek, Z., *Structure of geodesics in a 13-dimensional group of Heisenberg type*, to appear (Proceedings of the Colloquium on Diff. Geom., Debrecen 2000).
- [6] Kaplan, A., *On the geometry of groups of Heisenberg type*, Bull. London Math. Soc. **15** (1983), 35–42.
- [7] Riehm, C., *Explicit spin representations and Lie algebras of Heisenberg type*, J. London Math. Soc. **29** (1984), 46–62.

MATHEMATICAL INSTITUTE, CHARLES UNIVERSITY  
SOKOLOVSKÁ 83, 186 75 PRAHA 8  
CZECH REPUBLIC  
E-mail: Zdenek.Dusek@mff.cuni.cz